

## PREVALENCE AND RISK FACTORS FOR OVERWEIGHT AND OBESITY AMONG SUBURBAN SEMI-NOMADIC FULANI'S OF NORTHWESTERN NIGERIA

Ramalan MA<sup>1</sup>, Gezawa ID<sup>1</sup>, Uloko AE<sup>1</sup>, Musa BM,<sup>1</sup>

<sup>1</sup>Endocrinology, Diabetes and Metabolism Unit, Department of Internal Medicine, Aminu Kano Teaching Hospital, Kano.

### ABSTRACT

#### Background

Obesity is a major public health problem that is associated with increased cardiovascular morbidity and mortality. It is an important threat that is fast becoming an emerging epidemic globally, Nigeria inclusive. Although there are numerous studies on the prevalence of obesity in Nigeria, the semi-nomadic Fulani's have not been extensively studied. This study will, therefore, help to assist policymakers to plan effective strategies for combating the epidemic among neglected ethnic populations.

#### Method

The study was a community-based cross-sectional study carried out in Kumbotso Local Government Area of Kano state, from June 2015 to December 2015. Anthropometric indices (Weight, height, waist and Hip circumferences) were measured for the subjects. Body mass index and waist hip ratio was also calculated. Serum blood glucose, lipid, and blood pressure were determined.

#### Results

The mean  $\pm$ SD age of the subjects was 38.97 $\pm$ 15/98 years, with a range of 18 – 90 years. There were 208 (39.1%) males and 324 (60.9%) females,  $p < 0.001$ . Obesity was found in 9.21% of the subjects (4.32% of males, 13.58% of females). Overweight was found in 20.86% of the subjects, 15.38% of males and 13.58% of females respectively. Factors associated with overweight include advancing age and female gender. The odds ratio for developing obesity is higher in subjects > 50 years (2.319, 95% CI, 1.203-3.540).

#### Conclusion

There is a relatively high prevalence of overweight and obesity among suburban semi-nomadic Fulani's. In view of this, there is the urgent need to initiate public health measures aimed at improving healthy lifestyle measures.

**Key words:** Prevalence, overweight, obesity, risk factors and sub urban.

NigerJmed 2019; 360-367  
© 2019. Nigerian Journal of Medicine

### INTRODUCTION

Obesity is a major public health threat that is fast becoming an emerging epidemic globally, Nigeria inclusive. The prevalence of overweight and obesity is expected to rise

over the next few decades. This increase in prevalence is related to the changing demographics of our population and the adoption of western lifestyles characterized by consumption of energy-dense foods and decreased physical activity.<sup>1-3</sup>

It is one of the major modifiable cardiovascular risk factors and is associated with increased risk of developing diabetes mellitus and other endocrine disorders, cardiovascular diseases; hypertension, atherosclerosis, as well as certain types of cancer.<sup>4,5</sup>

**Correspondence to:** Ramalan MA  
Endocrinology, Diabetes and Metabolism Unit,  
Department of Internal Medicine,  
Aminu Kano Teaching Hospital,  
Tel: +2348036783737,  
E-mail: mramalan@gmail.com

In the United States, for instance, more than two-thirds (68.8%) of the adult population is considered to be either overweight or obese and a third (35.7%) of adults are obese. In men almost, a three quarter (74%) are considered to be overweight or obese. Extreme obesity has been estimated to affect about one-tenth of the population (6.3%).<sup>6</sup>

In Nigeria, the exact prevalence of obesity is not known but rates ranging from 8.1% to 22.2 % have been reported, while the prevalence of overweight ranged from 20.3% - 35.1%, which has been noted to have increased from an earlier study in 1970 by Johnson<sup>8</sup> in Lagos, which reported a prevalence of 14.9% and 26.5% males and females respectively.<sup>7-9</sup>

The economic consequences of obesity are enormous with a direct cost of managing people with obesity expected to rise to about \$48-66 billion/year in the USA and by £1 9-2 billion/year in the UK by 2030<sup>4</sup>. Indirect costs relating to disability from the disease, absenteeism at places of work, premature mortality and workers' compensation related to the disease is also expected to rise exponentially globally.<sup>6,7</sup>

Most studies on the prevalence of overweight and obesity in Nigeria were conducted in urban populations. This study sets out to determine the prevalence of overweight and obesity among ethnic Fulani's of Northwestern Nigeria who are mostly nomads with an increased level of physical activity by virtue of their means of livelihood.<sup>10</sup>

### **Materials and method**

The study was a community-based cross-sectional study carried out in Kumbotso LGA, which is one of the 44 Local Government Areas of Kano state, and one of the eight local government councils that

make up Kano Metropolis from June 2015 to December 2015.

Study participants included adult males and females aged 18years and above resident within the study area who meet the inclusion criteria and consented to participate in the study.

We excluded participants who were ill, pregnant women and all persons who declined consent to participate in the study.

### **Sampling Method**

A multi-stage sampling technique was used to select the participants. In stage one, Kumbotso was randomly selected out of the eight local governments in Kano metropolis. In the second stage, Kumbotso ward was randomly selected out of eleven wards in Kumbotso local government. Three settlements (Kamfa, Rigafada, and Bayan Dala) were selected out of the nine settlements that make up Kumbotso ward using simple random sampling technique (Balloting). House and Household listing were done in the three selected settlements.

In the third stage, a systematic sampling technique was used to select a proportionate number of households in the settlement after calculating a sampling interval for each settlement. Three hundred and sixty households were selected.

Finally, eligible respondents in the sampled households were requested to participate after obtaining signed informed consent (about 3 respondents per household). The first household was selected using a table of random numbers while adding the sampling interval to preceding household number identified subsequent households until the required sample size was obtained.

### **Ethical Consideration**

Ethical approval was obtained from Aminu Kano Teaching Hospital Kano Ethical

review committee (NHREC/21/08/2008/AKTH/EC/1224, and the Kano state ministry of health (HMB/GEN/488/VOL1).

Participants were assured of the confidentiality of their information and those who declined consent to participate in the study were excluded from the study.

### Study Population

The study population consists of suburban semi-nomadic Fulani's residing in the study area who fulfil the inclusion criteria. The Fulani's are the most widely dispersed ethnic group in West Africa and the Sahel although they were originally thought to have originated from North Africa<sup>10</sup>. The number in excess of 20 – 25 million and they are the largest groups of nomadic pastoral community in the world.<sup>10</sup>

### Study Procedure

Permission and cooperation for the study were obtained from the village and ward heads of the respective communities where participants were recruited.

Research assistants who speak English, as well as the local languages, were recruited and trained to assist in data collection and anthropometric measurements.

An interviewer-administered questionnaire based on a modification of WHO STEPS Instrument was used for data collection.<sup>11</sup>

Anthropometric measurements were carried out on each subject as follows;

**Weight in kilograms (kg)** -Was measured to the nearest 0.1kg with subjects in light clothing and without shoes while standing on a calibrated bathroom scale, positioned on a flat even surface.<sup>12</sup>

**Height in metres (m)** -Was measured to the

nearest 0.1cm using a height stand with subjects barefooted and without headgear while standing erect.<sup>12</sup>

**Body mass index (BMI)** - The BMI was calculated as weight in kg divided by the square of height in metres (m<sup>2</sup>) i.e. kg/m using the Quetelet formula  $BMI = \text{weight (kg)} / \text{height (m)}^2$ ,<sup>12</sup>

**Waist circumference (WC)** - in centimetre (cm) - The WC was measured to the nearest 0.1cm using a non-stretchable dressmaker's tape at a point mid-way between the margin of the lowest rib and the iliac crest in the horizontal level. This was done with the subject standing erect at the expiratory phase of respiration<sup>12</sup>

**Hip circumference (HC)** - in centimetre (cm) - The HC was measured to the nearest 0.1cm at the horizontal level of the maximum circumference around the gluteal region (posteriorly) and the pubic symphysis (anteriorly) with the tape parallel to the floor. This was done with the subject standing erect at the end of a normal expiration<sup>12</sup>

**Waist to hip ratio (WHR)** - The WHR was calculated as the waist circumference in centimetre (cm) divided by hip circumference in centimetre (cm).<sup>12</sup>

### Statistical Methods

Data were analysed using SPSS version 20.0 (IBM Corporation, New York USA.) Presentation of the data was done in tables. Prevalence of overweight and obesity was expressed in percentages. The socio-demographic and lifestyle risk factors of the subjects were presented using descriptive statistics and the chi-squared test. Mean values were compared using the independent t-test.

P values <0.05 were considered statistically significant.

## Results

A total of five hundred and thirty subjects completed the study (532). The mean  $\pm$ SD age of the subjects was  $38.97 \pm 15/98$  years, with a range of 18 – 90 years. There were 208 (39.1%) males and 324 (60.9%) females.  $p < 0.001$ . The mean  $\pm$ SD age of the females and males was  $43.4 \pm 18.7$  and  $39.7 \pm 16.2$  respectively, with the latter being younger. The majority of the subjects were in their second and third decade of life.

The socio-demographic characteristics of the study population are as shown in Table 1.

## Anthropometric Indices

Table 2 shows the mean anthropometric indices of the study population by gender. The mean values of BMI and WC, were higher among females compared with males, while the reverse was the case with the WHR, which was noted to be higher among males although the difference was not statistically significant

## Prevalence of overweight and obesity

The prevalence rates of overweight and obesity among the study population based on BMI is shown in Table 3. The prevalence of overweight was 15.38% in males and 24.38% in females, while obesity was found in 4.32% of males and 13.58% of females, respectively.

About a third of the females was either overweight or obese, while about a quarter of the males was either overweight or obese. Class 1 obesity is the commonest pattern in both genders. None of the male subjects was morbidly obese (class 3 obesity).

## Clinical and laboratory variables

The values of anthropometric indices were higher in obese subjects than in non-obese subjects, however there is no statistically

significant difference in the systolic and diastolic blood pressures of the subjects as shown in table 4.

There was no difference in the mean values of blood glucose of the obese and non-obese subjects. However, there is a statistically significant difference in the mean values of the lipid profile of the obese and non-obese subjects. This is depicted in table 5.

## Risk factors for overweight and Obesity

The risk factors for developing overweight and obesity as established from this study obtained from multi variate analysis include advancing age (>50 years) and female gender.

This is shown in table 6.

## Discussion

There is the paucity of knowledge on the prevalence of overweight and obesity among the semi-nomadic Fulani ethnic group. To the best of our knowledge, this is the population-based study examining the overweight and obesity among the semi nomadic Fulani's of Northern Nigeria.

In this study, the prevalence of overweight and obesity is lower than reported from previous studies. Gezawa et al<sup>9</sup> reported prevalence rates of overweight and obesity of, 27.1% and 17.1%, respectively in a study among 1650 urban dwellers, while Puepet et al<sup>13</sup> reported obesity prevalence rates of, 19.4% and 23.5% among males and females respectively, in a study of 825 adults in Jos metropolis. An earlier study by Bakari et al reported a prevalence of 13.1% and 18.5% respectively for overweight and obesity in a suburban Northern Nigerian population consisting of 317 subjects. The finding in our study is also lower than the 6.0% reported by Iloh<sup>14</sup> et al in a study of 2156 adult Nigerians seeking medical care at a mission General Hospital in Imo state, Southeastern Nigeria. The prevalence of overweight and

obesity found in this study is however similar to the 21% and 4% respectively, reported from a study among a rural community of Benue State North-central Nigeria by Adediran<sup>16</sup> et al in 2011. The lower rates of overweight and obesity reported in this study may be related to the study population, with urban dwellers having higher rates compared with rural residents and the increased level of physical activity among the nomads, however much lower rates of overweight and obesity have been reported from Northern Cameroon by Kufe et al (6.1and 2.2% respectively).<sup>18</sup> Diet and lifestyle may also contribute to the lower rates of obesity and overweight in the study population.

More women than men were found to be overweight and obese in our study which is consistent with reports from previous studies.<sup>9,13</sup> The higher prevalence of obesity among females in this study may be due to lower levels of physical activity among women compared with the men who travel long distances to rear their animals and add to their engagement in farming activities.

### Conclusion and Recommendation

This study depicts the effects of culture and environment in modulating the development, cause and progression of the disease.

The strength of this study includes the fact that it is a community-based study of an ethnic group hitherto not extensively studied with respect to the prevalence of obesity. The use of the standardized WHO questionnaire enabled us to compare this study with studies done elsewhere.

More studies are therefore needed to be done in order to develop intervention programs and strategies that will help in the long run to curb the rising tide of non-communicable disease in the suburban population.

### Limitations of the study

Because it is a cross-sectional study, it is limited by the fact that the temporal profile of the disease occurrence cannot be determined.

Inability to take diet history and food intake pattern of the participants.

### Conflicts of Interest

We declare that there is no potential conflict of interest.

### Acknowledgments

The study was self-funded.

### Authors Contribution

All the authors contributed equally to the study.

### Tables

**Table 1. Socio demographic characteristic of the study population by Age and Gender**

Variable	Male		Female		Total	
	N	%	N	%	N	%
Age (Years)						
<30	74	35.58	89	27.47	163	30.64
30-39	66	31.73	80	24.69	146	27.44
40-49	27	12.98	61	18.83	88	16.54
50-59	20	9.62	43	13.27	63	11.84
60-69	13	6.25	20	6.17	33	6.20
70	8	3.85	31	9.57	39	7.33
Total	208	100.00	324	100.00	532	100.00
<b>Educational Status</b>						
Able to read or write						
Yes	143	68.75	93	28.70	236	44.36
No	65	31.25	231	71.30	296	55.64

Total	208	100.00	324	100.00	532	100.00
<b>Level of Formal Education</b>						
Quranic Education	77	53.85	48	51.61	125	52.97
Primary Education	55	38.46	41	44.09	96	40.68
Secondary Education	11	7.69	4	4.30	15	6.36
Total	143	100.00	93	100.00	236	100.00
<b>Marital Status</b>						
Single	47	22.60	25	7.72	72	13.53
Married	159	76.44	256	79.01	415	78.01
Separated	0	0.00	41	12.65	41	7.71
Widowed	2	0.96	2	0.62	4	0.75
Total	208	100.00	324	100.00	532	100.00

**Table 2. Mean Anthropometric indices of the study population by Gender**

Anthropometric Index	Gender	N	Mean	<i>p</i> Value
<b>BMI</b>	Male	208	22.56	0.001*
	Female	324	24.35	
<b>WC</b>	Male	208	81.41	0.001*
	Female	324	84.78	
<b>HC</b>	Male	208	91.84	0.003*
	Female	324	96.54	
<b>WHR</b>	Male	208	0.89	0.117
	Female	324	0.88	

BMI= Body Mass Index; WC = Waist circumference; HC= Hip Circumference; WHR= Waist Hip Ratio; \*  $p < 0.05$

**Table 3. Distribution of the study population by BMI**

BMI category	Male n =208		Female n= 324		Total N= 532		<i>P</i> Value	
	N	%	N	%	N	%		
<b>Underweight</b>	15	7.21	35	10.80	50	9.40	0.001*	
<b>Normal</b>	152	73.08	166	51.23	318	59.77	0.001*	
<b>Overweight</b>	32	15.38	79	24.38	111	20.86	0.001*	
<b>Obesity</b>	<b>Class 1</b>	6	2.88	30	9.26	36	6.77	0.001*
	<b>Class 2</b>	3	1.44	10	3.09	13	2.44	0.001*
	<b>Class 3</b>	0	0.00	4	1.23	4	0.75	0.001*
Total	208	100.00	324	100.00	532	100.00		

BMI= Body Mass Index; \*  $p < 0.05$

**Table 4: Mean values of anthropometric indices and Clinical Parameters of the study subjects**

Mean Parameter	Obesity Present	Obesity Absent	<i>P</i> Value	
Height	Total	1.59	1.62	0.003*
	Male	1.68	1.68	0.887
	Female	1.57	1.58	0.224
Weight	Total	86.42	59.40	0.001*
	Male	94.00	62.50	0.001*
	Female	84.06	57.17	0.001*
BMI	Total	34.08	22.94	0.001*
	Male	33.06	22.09	0.001*
	Female	34.29	22.79	0.001*
WC	Total	104.50	81.14	0.001*
	Male	104.56	80.36	0.001*
	Female	104.49	81.68	0.001*

HC	Total	112.38	92.75	0.001*
	Male	108.33	91.10	0.001*
	Female	113.20	93.92	0.001*
WHR	Total	0.93	0.87	0.001*
	Male	0.97	0.88	0.001*
	Female	0.92	0.87	0.001*
BPS	Total	137.70	131.86	0.001*
	Male	129.56	127.17	0.644
	Female	139.36	135.19	0.394
BPD	Total	89.87	82.57	0.062
	Male	86.22	79.14	0.062
	Female	90.16	85.01	0.043*

BMI= Body Mass Index; WC = Waist circumference; HC= Hip Circumference; WHR= Waist to Hip Ratio; BP= Blood Pressure; BPS= Systolic Blood Pressure; BPD= Diastolic Blood Pressure;\*  $p < 0.05$

**Table 5: Mean values of biochemical parameters among obese and non-obese subjects**

Mean Parameter	Obesity Present	Obesity Absent	P Value	
FPG	Total	4.29	4.39	0.469
	Male	4.35	4.56	0.589
	Female	4.29	4.31	0.549
CPG	Total	5.27	5.81	0.857
	Male	5.22	5.85	0.501
	Female	5.87	5.78	0.172
TChol	Total	4.75	3.67	0.640
	Male	4.40	3.50	0.001*
	Female	4.82	3.50	0.020
HDL	Total	1.10	0.95	0.001*
	Male	1.19	0.94	0.001*
	Female	1.09	0.96	0.030*
LDL	Total	2.71	2.05	0.150
	Male	2.20	1.17	0.001*
	Female	2.80	2.15	0.001*
Trig	Total	1.53	1.21	0.015*
	Male	1.83	1.17	0.001*
	Female	1.47	1.24	0.010*

T Chol = Total Cholesterol; HDL = High Density Lipoprotein Trig = Triglyceride; HDL = High Density Lipoprotein; LDL = Low Density Lipoprotein; FPG= Fasting Plasma glucose; CPG= Casual Plasma Glucose;\*  $p < 0.05$

**Table 6. Odds ratio for Obesity**

Parameter	Odds Ratio	95 % CI	pValue
Males	0.621	0.351-1.317	
Females	2.273	1.176-2.729	0.01*
≥ 50 Years	2.319	1.203-3.540	0.01*
Hypertension	0.928	0.569-1.182	0.23

\*  $p < 0.05$

## References

- Sani MU, Wahab KW, Yusuf BO, Gbadamosi M, Johnson OV, Gbadamosi A. Modifiable cardiovascular risk factors among apparently healthy adult Nigerian population – a cross-sectional study. BMC Res Notes. 2010;3:11.
- World Health Organization. Physical Status: The Use and Interpretation of Anthropometry. Geneva, Switzerland:

- World Health Organization; 1995. Technical Report Series; p. 854. 1-1-9950.
3. Obesity: preventing and managing the global epidemic. Report of a WHO consultation. World Health Organ Tech Rep Ser. 2000;894: i–xii. 1–253.
  4. Bray GA. Complications of obesity. *Ann Intern Med.* 1985;103(6 (Pt 2)):1052-62.
  5. Heraief E. The contribution of epidemiology to the definition of obesity and its risk factors. *Ther Umsch* 1989 May;46(5):275-80.
  6. Trogdon JG, Finkelstein EA, Hylands T, Dellea PS, Kamal-Bahl SJ. Indirect costs of obesity: a review of the current literature. *Obes Rev.* 2008; 9:489-500
  7. Chukwuonye, I. I., Chuku, A., John, C., Ohagwu, K. A., Imoh, M. E., Isa, S. E. et al. Prevalence of overweight and obesity in adult Nigerians – a systematic review. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*; 2013: 6, 43–47.
  8. Johnson T.O. Prevalence of overweight and obesity among adult subjects of an urban African population sample. *British Journal Prev. Soc. Med.* 1970; 24: 105-109.
  9. Gezawa ID, Puepet FH, Mubi BM, Uloko AE, Bakki B, Talle MA. Prevalence of overweight and obesity in Maiduguri, North-Eastern Nigeria. *Niger J Med.* 2013; 22(3): 171-4.
  10. David L. Fulani. *Encyclopedia of world cultures: Africa and the Middle East.* Vol. 9. 1996
  11. World Health Organization WHO STEPwise approach to chronic disease risk factor surveillance-instrument v2.0 Department of Chronic Diseases and Health Promotion. World Health Organization 20 Avenue Appia, 1211 Geneva 27, Switzerland. (Available at <http://www.who.int/chp/steps/en/>)
  12. World Health Organization Expert Committee: Physical status; The use and interpretation of anthropometry. Report of a WHO expert committee. Technical report series 854, WHO Geneva, 1995.
  13. National Institute of Health. The Practical guide to the identification, evaluation and treatment of overweight and obesity in adults, the evidence report. *Obes Res* 1998; 6 (2). 515-2095.
  14. Puepet FH, Zoakah AJ, Chuhwak EK. Prevalence of overweight and obesity among urban Nigerian adults in Jos. *Highland Medical Research Journal* 2002; 1: 13-1.
  15. Iloh GUP, Amadi AN, Nwankwo BO, Ugwu VC. Obesity in adult Nigerians: A study of its patterns and common primary co-morbidities in a rural Mission General Hospital in Imo State, Southeastern Nigeria. *Niger J Clin Pract.* 2011;14(2) 212-218.
  16. Bakari AG, Onyemelukwe GC, Sani BG, Aliyu IS, Hassan SS, Aliyu TM. Obesity, overweight and underweight in suburban northern Nigeria. *Int J Diabetes Metab* 2007; 15: 68-69.
  17. Adediran SO, Adeniyi SO, Jimoh KA, Alao OO. Underweight, Overweight and Obesity in Adult Nigerians living in rural and urban communities of Benue State. *Ann Afr Med* 2011; 10(2):139-143.
  18. Kufe NC, Ngufur G, Mbeh G, Mbanya JC. Distribution and patterning of non-communicable disease risk factors in indigenous Mbororo and non-autochthonous populations in Cameroon: a cross-sectional study. *BMC Public Health.* 2016; 16:1188