

Daniyan OW
Ezeanosike OB
Ogbonna-Nwosu C
Iloduba UC

CC-BY



Thanatophoric dysplasia type 1 as seen in a tertiary institution in South-East Nigeria: A case report

DOI:<http://dx.doi.org/10.4314/njp.v47i3.14>

Accepted: 9th April 2019

Daniyan OW (✉)
Ezeanosike OB
Ogbonna-Nwosu C, Iloduba UC
Department of Paediatrics,
Alex-Ekwueme Federal University
Teaching Hospital, Abakaliki
Email: daniyanolapeju@gmail.com

Abstract: Thanatophoric dysplasia is a lethal form of skeletal dysplasia seen in neonates. The word ‘thanatophoric’ is derived from the Greek word thanatophorus meaning death bringing. Thanatophoric dysplasia results from mutations within the Fibroblast Growth Factor Receptor 3 (FGFR3) gene which is located on chromosome 4p16.3. A female neonate with dysmorphic features such as macrocephaly, frontal bossing, periorbital swelling and depressed nasal bridge was deliv-

ered to a 35year old woman. The upper and lower limbs were short with excessive skin folds. A case of female neonate with thanatophoric dysplasia is hereby reported to raise awareness of this condition and to describe the features of thanatophoric dysplasia seen in this patient .

Key words: Thanatophoric Dysplasia (TD), Fibroblast Growth Factor Receptor 3 Gene (FGFR3), dysmorphic, macrocephaly

Introduction

Thanatophoric dysplasia (TD) is a lethal form of skeletal dysplasia seen in neonates. The word ‘thanatophoric’ is derived from the Greek word thanatophorus meaning death bringing as described by Maroteaux et al.¹ Thanatophoric dysplasia results from mutations within the Fibroblast Growth Factor Receptor 3 (FGFR3) gene which is located on chromosome 4p16.3.² The condition is characterized by limb shortening, macrocephaly, folds of redundant skin, narrow chest and pulmonary hypoplasia.^{3,4} Thanatophoric dysplasia has two major subtypes TD I and TD 2. Type 1 has a curved limb while type 2 has straight limb with cloverleaf shaped head. A case of female neonate with thanatophoric dysplasia is hereby reported to raise awareness of this condition and to describe the features of thanatophoric dysplasia seen in this patient .

Case report

A case of a female neonate with dysmorphic features who was delivered at the Alex Ekwueme Federal University Teaching Hospital Abakaliki in May 2019 to a booked 35 year-old para 9 woman at a gestational age of 40 weeks through emergency caesarean section due to prolonged labour. Mother was not regular with antenatal visit and had taken herbal mixtures consisting of leaves mixed with alcohol at various times during the period of pregnancy. She had no history of hypertension or diabetes and no family history of congenital abnormalities. Had only one prenatal scan done during early trimester, which showed no abnormality. Father was a 60 year old man.

At delivery APGAR score was 3¹, 5⁵ and 6¹⁰ Resuscitation with intermittent positive pressure ventilation (IPPV) with bag and mask was done at birth. She had respiratory distress and was admitted into to the newborn intensive care unit (NICU) where she was commenced on intranasal oxygen. The baby had dysmorphic features such as macrocephaly with head circumference of 41cm (>97th percentile), frontal bossing, periorbital swelling and depressed nasal bridge. The upper and lower limbs were short with excessive skin folds . The chest appeared small with protuberant abdomen. The weight was 3.25kg with a length of 38cm. diagnosis of thanatophoric dysplasia type 1 was made. The baby died 3 hours after delivery.

Newborn with features suggestive of thanatophoric dysplasia type 1



Discussion

Thanatophoric dysplasia is a lethal skeletal dysplasia and is classified in group 1 of the FGFR3 chondrodysplasia group on molecular bases.⁵ It is inherited in an autosomal dominant fashion.

The prevalence of thanatophoric dysplasia has been estimated to be 1.1 per 100,000 births in Japan⁶ and a prevalence of 0.21 to 0.30 per 10,000 live births in US.⁷ There are no available data on the prevalence of thanatophoric dysplasia in Nigeria, this may be due to under reporting. Few case of TD has been reported in Nigeria, Joel-Medewase *et al*⁸ and Komolafe⁹ reported cases of TD in South-West Nigeria.

FGFR3 is a negative regulator of bone growth and its mutation is responsible for TD. The mutation results in activation of FGFR3 tyrosine kinase which sends negative signals to the chondrocytes which result in inhibition of cell division, stimulation of cell maturation and differentiation giving rise to abnormal bone development.¹⁰ An R248C mutation has been found in the FGFR3 gene.¹¹ Also missense mutation of Y373C, S371C and S249C has been reported in some cases of TD.¹²

Similar features seen in this neonate like shortening of the limbs, macrocephaly, short neck, depressed nasal bridge, narrow thorax with distended abdomen has been reported by other authors.^{13,14} The long bones in TD 1 are curved and appears like telephone receiver on radiographs.¹⁵ Thanatophoric dysplasia type 2 is characterized by clover leaf shaped skull due to premature closure of coronal and lambdoid sutures and the long bones are not bent or curved.^{16,17}

Most cases of TD are delivered still birth and those born alive rarely survive due to pulmonary insufficiency from

narrow chest and hypoplastic lungs commonly seen in TD.¹⁸ Thanatophoric dysplasia should be differentiated from other causes of skeletal dysplasia like achondroplasia, achondrogenesis, osteogenesis imperfecta, camptomelic dysplasia. Achondroplasia is a non lethal form of chondrodysplasia characterized by big head, short limb that is marked in the proximal segment and fingers that assume trident position.¹⁹

TD can be diagnosed based on clinical features or use of radiographic studies either at the prenatal period by use of ultrasonography or in the immediate newborn period. On the index case the parents refused further investigation on the baby. The mother had only one prenatal ultrasound scan. A late scan in pregnancy may have detected the clinical condition and this would have prepared the parents for the birth of the baby.

An autopsy could also be done to determine the severity of the disease condition in babies with TD, autopsy was not done in this case because the parents did not give consent.

Conclusion

This is a case of a female neonate with clinical features of TD 1. The report underscores the importance of ultrasonography in pregnancy for early detection of fetal anomalies to enable appropriate counseling of the parents, preparation for birth of the baby and early intervention where indicated.

Conflict of Interest: None

Funding: None

References

1. Maroteaux P, Lamy M, Robert JM. Thanatophoric dwarfism. *Presse Med* 1967;75: 2519-2524
2. Cohen JR. Achondroplasia, hypochondroplasia and thanatophoric dysplasia: clinically related skeletal dysplasia that are also related in molecular level. *Int J Oral Maxillofac* 1998; 27: 451-5
3. Thanatophoric dysplasia. Genetic home reference. [Cited 2019 July 17]. Available from: <https://ghr.nlm.nih.gov>
4. Kocheria K, Kocheria V. Antenatal diagnosis of thanatophoric dysplasia: a case report and review of the literature. *Int J Res Med Sci* 2014; 2: 1176-9.
5. Warman MJ, Cormier-daire V, Hall C, Krakow D, Lachman R, LeMerrer M *et al*. Nosology and classification of genetic skeletal disorder: 2010 revision. *Am J Med Genet A* 2011; 155: 943-68.
6. Sawai H, Oka K, Ushioda M, Nishimura G, Omori T, Numabe H *et al*. National survey of prevalence and prognosis of thanatophoric dysplasia in Japan. *Pediatr Int* 2019;61. DOI: <https://doi.org/10.1111/ped.13927>
7. Waller DK, Correa A, Vo TM, Wang Y, Hobbs C, Langlois PH *et al*. The population based prevalence of achondroplasia and thanatophoric dysplasia in selected regions of the US. *Am J Med Genet Part A* 2008; 146A: 2385-9
8. Joel-Medewase V, Aworinde O, Alabi A, Akinremi O, Lekan-Kehinde O. Lethal Neonatal Rhizomelic Dwarfism: A Report of Two Cases. *Int J Health Sci Res* 2018; 8:178-83
9. Komolafe F, Adetoro OO, Adetoro OO. Thanatophoric dwarfism in a Nigerian child. *Afr J Med Med Sci* 1986; 15:109-12
10. Search articles by 'Adetoro OO' Germaine GL. Thanatophoric dysplasia. Medscape (updated September 2018). Cited [2019 July 17]. Available from: <https://emedicine.medscape.com/article/949591>

11. Noe EJ, Yoo HW, Kim KN, Lee SY. A case of thanatophoric dysplasia type I with an R248C mutation in the FGFR3 gene. *Korean J Pediatr* 2010; 53: 1022-5
12. Jung M, Park SH. Genetically confirmed thanatophoric dysplasia with fibroblast growth factor receptor 3 mutation. *Exp Mol Pathol* 2017; 102: 290-5.
13. Davanageri RS, Shokeen PD, Bannur HB, Patil KP. Thanatophoric dysplasia type 1: A rare case report at Fetal autopsy. Reshma S Davanageri. *J Lab Physicians* 2014; 6: 121-3.
14. Naveen Ns, Murlimanju BV, kumar V, Pulakunta T, Jeeyar H. Thanatophoric dysplasia: a rare entity. *Oman Med J* 2011; 26: 196-7.
15. Chavadi C, Mahale N, Jasthi G, Prabhu S, Kory S. Thanatophoric dysplasia: Antenatal to postterm. *J Sci Soc* 2015; 42: 210-2.
16. Miller E, Blaser S, Shannon P, Widjaja E. Brain and bone abnormalities of thanatophoric dwarfism. *A J Roentgenol* 2009; 192: 48-51.
17. Gulasi S, Atici A, Celik Y. A case of thanatophoric dysplasia type 2: A novel mutation. *J Clin Res Pediatr Endocrinol* 2015; 7: 73-6.
18. Jahan U, Sharma A, Gupta N, Gupta S, Usmani F, Rajput A. Thanatophoric dysplasia: a case report. *Int J Reprod Contracept Obstet Gynecol* 2019; 8: 758-61.
19. Sethi RS, Kumar L, Chaurasiya OS, Gorakh R. Case report : Achondroplasia. *Curr Pediatr Res* 2011; 15: 137-40