



Paediatric Emergency Admissions, Mortalities, and Unmet Intensive Care Needs at a Tertiary Hospital in Southern Nigeria

Eki-Udoko Fidelis E¹, Ani Chidiebere¹, Osagie Ekienabor G,¹
Atimati Anthony O^{1,2}

¹Department of Child Health, University of Benin Teaching Hospital, Benin City, Edo State, Nigeria.

²Department of Child Health, University of Benin, Benin City, Edo State, Nigeria.

Correspondence

Dr Fidelis E. Eki-Udoko, Department of Child Health, University of Benin Teaching Hospital, Benin City, Edo State, Nigeria. E-mail: f.ekiudoko@ubth.org ; ORCID – <https://orcid.org/0000-0002-9672-6150>.

Abstract

Background: The essence of seeking medical services is to be provided with essential medical care to prevent complications and possibly death from the illness.

Objective: To examine the pattern of admissions into the Children's Emergency Room and unmet intensive care needs in a tertiary health facility.

Methods: This prospective, cross-sectional study was conducted at the University of Benin Teaching Hospital (UBTH), Benin, Nigeria, over a 24-month period (2018 – 2019).

Results: A total of 10 138 children presented to the children's emergency room, and 2 914 children (28.6%) were admitted. There were a total of 144 mortalities (4.94%) of the total admissions. Infectious diseases accounted for the Majority of the deaths. The commonest morbidity necessitating admission was meningitis, while the sickle cell crisis and oncologic pathologies contributed the least - most children presented with multiple morbidities. The under-5s made up 61.1% of deaths recorded. More deaths amongst males compared to female children (1.3:1.1). Majority of the deaths occurred within 12 hours of presentation. Of the 144 mortalities recorded, 140 (97.2%) required intensive care services, while four did not qualify for ICU care. Of the 140 children who qualified for ICU care, 17 (12%) were admitted into the ICU for further care, of which only 2 (12.3%) survived and were discharged home. The others died.

Conclusion: The persistently high contribution of infectious disease to infant and child mortality, coupled with an inability to offer intensive care services, should be an important consideration for health planners and administrators.

Keywords: Admissions, Intensive Care Services, Paediatric Emergency, Tertiary Hospital, Unmet needs.

Introduction

The essence of seeking medical services is to be provided with essential medical care to prevent complications and possibly death from the illness. The death of a child at the emergency ward is one of the most difficult problems that the clinicians attached to these wards have to deal with. The Sustainable Development Goals (SDGs) represent the transition from the Millennium Development Goals, and goal five

has universal coverage of quality healthcare as a target, including the prevention and treatment of communicable and non-communicable diseases.² To achieve this goal, accurate information about the causes of morbidity and mortality within various populations is essential.³

Moreover, there are reports on the mortality and morbidity patterns at the paediatric emergency

units locally,^{4,5} nationally⁶⁻¹⁶ and internationally.¹⁷⁻³⁰ Still, there is no record of any study that examined the met or unmet intensive care needs in children hospitalized as emergencies, and their contributions to the mortality rate at the emergency rooms. This study examined the admission pattern into the paediatric emergency room and the unmet intensive care needs at the University of Benin Teaching Hospital in Southern Nigeria. Information from this study may provide insight into paediatric mortality rates, and baseline data of unmet intensive care needs essential to health care providers and administrators. This will help them in allocating resources, especially to paediatric intensive care units, and the development of interventions to reduce preventable childhood deaths in Nigeria.

Methods

This was a cross-sectional study conducted at the Children's Emergency Division [Children Emergency Room (CHER) and Children's Emergency Ward (CHEW)] of the University of Benin Teaching Hospital (UBTH), Benin City, Edo State, Nigeria. This is a tertiary health facility in the South-South region of Nigeria. It provides all health care services to children from under 24 hours to 18 years old and is open for 24-hour service delivery. This facility has 28 beds, comprising the 20-bed CHEW and 8-bed CHER.

This unit provides care for children who require prompt life-saving measures, after which they are discharged as soon as possible from the unit, preferably within 24 to 48 hours of admission, either by transfer to the main paediatric wards or home to continue treatment as outpatients. The attending paediatrician and nursing staff prioritize all patients according to the severity of their condition (Emergency Triage Assessment and Treatment; ETAT),³¹ ensuring that the more critically ill child is attended to first. The unit is staffed by two consultants, seven resident doctors, eight medical house officers, four nursing interns and eighteen nurses.

The unit caters for children who present directly from home and referred cases from UBTH Consultant Outpatient Departments and General Practice Clinics. It also attends to referrals from other primary, secondary, and tertiary health care facilities in all parts of Edo State and the neighbouring States of Delta, Bayelsa, Kogi, Anambra, and Ondo, respectively.

All children brought into the CHER of the UBTH were recruited into this study over two years, from January 2018 to December 2019. Ethical approval was obtained from the Ethics Review Committee of UBTH (Protocol number: ADM/E 22/A/VOL.VII/148301230) and the parental/caregivers' consent and patient's assent were obtained where applicable.

Data was collected from the patient's case notes only and the information recruited included age, gender, date of presentation, clinical diagnosis, duration of admission, outcome and time of death, where applicable. The diagnoses represent the final diagnoses reached mainly after review by the team of doctors working in the CHER and CHEW. The diagnosis was based on clinical features and relevant laboratory investigations, where possible. Diagnosis of sepsis and acute gastroenteritis were mainly clinical and supported by laboratory investigations, while malaria was diagnosed with microscopy or a rapid diagnostic test. In a few cases, no diagnosis was reached before the patient's demise. All children with a severe cardiopulmonary compromise for any reason qualified for paediatric intensive care services. All children brought-in-died (BID) were excluded from this study.

At the time of this study, there was no designated PICU at the study locale. The main hospital intensive care unit (ICU) was the only one available to offer intensive care services. Children needing intensive care were offered care subject to the availability of spaces and the

Paediatric Emergency Admissions, Mortalities, and Unmet Intensive Care Needs at a Tertiary Hospital in Southern Nigeria

ability to settle the financial implications required for admission.

Data analysis was done with Statistical Package for the Social Sciences (SPSS), version 20.0 (IBM Corporation, Chicago, USA, 2011). Categorical data were summarized using proportions and percentages, while continuous data were summarized using mean and standard deviation for normally distributed variables and median for non-normally distributed variables. Statistical differences between the categorical variables were determined using the Pearson Chi-Square test of association, and $p < 0.05$ at

95% confidence interval was considered as being statistically significant.

Results

As shown in Figure 1, a total of 10,138 children presented to the children's emergency room over the 24 months of the study. The highest numbers of children were seen in September (1046; 10.3%) and May (1000; 9.9%), while January and December contributed the lowest proportions (3.7% and 6.4% respectively).




Figure 1: Monthly distribution of children presenting to the Emergency Unit

As shown in Figure 2, of the 10,138 children seen, 2,914 (28.7%) were admitted; the last three months of the year contributed nearly half (40.1%) of the total number of admissions, while May and June contributed the lowest proportions, with 2.5% and 2.3% of the total admissions respectively. The admission rate was low at the start of the year and gradually rose through the course of the year.

Figure 3 shows the distribution of mortalities per month. There were 144 mortalities, which comprised 4.94% of the total admissions. The months of June and September had the highest mortality rates at 12.5% each, while the lowest rates were reported in July and February at 2.8% and 5.6%, respectively. The mortality rate gradually increased towards the middle of the year and then declined again towards the end of the year.




Figure 2: Monthly distribution of children admitted from CHER into CHEW




Figure 3: Monthly distribution of the mortalities recorded at the Emergency Unit

Mortality was highest in patients with meningitis, closely followed by those with sepsis/septic shock, severe malaria and anaemia with heart failure, as shown in Table I. The commonest morbidity necessitating admission was meningitis, while sickle cell crisis and oncologic pathologies constituted the least. The case fatality rate was highest amongst children

with pneumonia compared to other illnesses. It is important to note that most children presented with multiple morbidities. Children within the first year of life contributed over a third of the mortalities within the two-year study period. As shown in Table II, the under-5s made up 61.1% of the total deaths reported. There were more male deaths, giving a male-to-female ratio of 1.3:1.

Table I: Frequencies of admission diagnoses, mortality rates and case fatality rates (CFR)

	<i>Number of cases</i>	<i>Number of deaths</i>	<i>CFR (%)</i>
Meningitis	707 (24.3)	68	9.6
Sepsis/Shock	603 (20.7)	58	9.6
Severe malaria	542 (18.6)	52	9.6
Anaemia/Heart Failure	395 (13.5)	38	9.6
Pneumonia	369 (12.7)	36	9.7
SCA	164 (5.6)	12	7.3
Cancer	134 (4.6)	7	5.2
Total	2914 (100)	253*	

*Some cases had more than one diagnosis *Sickle Cell Anaemia (SCA)

Numbers in parentheses are percentages of the total

Table II: Sex and age distribution of mortalities

	<i>Sex</i>			
	Male	Female	Total	Male-to-Female ratio
1-12months	30 (58.8)	21 (41.2)	51 (35.4)	1.4:1
>12months – 59 months	20 (54.1)	17 (45.9)	37 (25.7)	1.2:1
>59months	31 (55.0)	25 (45.0)	56 (38.9)	1.2:1
Total	81 (56.3)	63 (43.7)	144 (100.0)	1.3:1

A quarter died within 6 hours of presentation, while 75% were early mortalities (between the 7th and 48th hour of presentation). Of the 144 critically ill children admitted, 140 (97%) required intensive care (ICU) services. In contrast, 4 (3%) did not qualify for intensive care as they were brought in actively dying (resuscitated but died within an hour of presentation).

Intensive Care Services Unmet, Offered and Outcome

Of the 140 children who qualified for ICU care, only 17 (12%) were admitted into the ICU, while the remaining 123 (88%) did not receive the needed intensive care, and all of them died. Of the 17 (12%) children that were admitted into the ICU, only 2 (12.3%) survived and were discharged home, while the remaining 15 children died (87.7%).

Duration of illness, use of native herbs, over-the-counter drugs and treatments received at referring hospital/chemist shops

The duration of illness among the mortalities ranged from a few hours to months before presentation at our facility. The majority (97%) of the mortalities presented late with complications of the illnesses. Of all the mortalities recorded in this study, 98% of the children received medications bought from patent medicine stores or native herbs administered by caregivers. In contrast, others had intravenous or oral drugs at the referral facilities prior to presentation at UBTH. All these interventions outside the hospital setting delayed the time of proper medical evaluation and prompt treatment. The researcher could not ascertain the method of care cost payment.

Discussion

The admission rate in this study is much higher than that reported from similar studies like

those from tertiary facilities within the south-south and southwestern regions of Nigeria^{7,8} and the report by Cui-ping Zhu in China.³⁰ The high admission rate may be due to the failure of the primary and secondary levels of health services in the study locale and environs resulting in over patronage of the tertiary health care system.

The number of participants and the admission rate in this study are much higher than the values reported from similar studies in other parts of the country^{7-9,16,30} The higher number obtained in this study is reflective of the fact that UBTH is one of the busiest tertiary facilities in the country and serves as a referral centre for several neighbouring states. Edo State, being a confluence state, also has a heavy population of people going through and residing transiently for business and pleasure.

The months that reported the highest mortalities had conversely low admission rates. Of note is that these months witnessed industrial actions by various healthcare worker unions. These periods would have adversely affected the quality of care given to the patients, resulting in poorer outcomes. It is also reflective of the fact that periods of reduced staff strength per patient have a significant negative impact on the quality of clinical outcomes. Similar findings were noted by Agbesanwa *et al.*⁸ in southwestern Nigeria. A mortality rate of 4.9% of admissions observed in this study compares favourably with findings from similar studies in this region by Ibeziako and co-workers³² and the previous report from the study locale.⁵ Other researchers have reported higher mortality rates; Agbesanwa *et al.*⁸ in Ekiti State, an earlier study in Benin City,⁴ the study conducted at the children's emergency room of the Abia State University Teaching Hospital (9.6%)³³ and previous Nigeria studies, showed mortality rates of 11.1%, 12.6%, 9.5%, 10.0%, 14.3%, and 15.1%^{11, 34-38} Conversely, Enyuma and colleagues⁹ reported a mortality rate of 0.5% which is especially low in comparison to findings from other facilities in the region. The finding in this study suggests that mortality

rates have not significantly changed over the last two decades.

The spread of morbidities and mortalities by age, as recorded in this study, is consistent with the high mortality rate in under-5s reported in other studies^{32, 39-41} and especially within the first year of life. Most of the children admitted were under-5, with children within the first 12 months of life having the largest contribution. This finding further highlights the vulnerability of children in these age groups to communicable diseases. Similar findings were noted by other researchers reporting admission patterns from children's emergency rooms in Nigeria.^{5,7,9,16,42}

As observed in this study, males presented more in CHER, in keeping with previous studies.^{7, 43-45} This observation could be buttressed by the family's male sex preference attitude in Nigeria^{34,35} and certain parts of Africa and Asia, whereby families seek health care earlier and more frequently for their male children than for their daughters. Families in Nigeria and certain parts of Africa and Asia preferring males over female children, both in education and health care seeking behaviours' is a documented observation in literature.⁴⁴⁻⁴⁶ Generally, the male-to-female ratio of the study participants was skewed in favour of the males; mortality was slightly higher among males compared to females. This is in tandem with other studies in Nigeria, as reported by Agbesanwa and colleagues⁸ in Ekiti state, Duru *et al.*⁷ in Bayelsa, and Njoku *et al.*¹⁶ from the Southeastern part of Nigeria for reasons earlier stated above.

Case fatality was highest in patients who were diagnosed with pneumonia, closely followed by those with meningitis, sepsis and septic shock, severe malaria and anaemia with heart failure. The pattern of infectious diseases among the mortalities is similar to the findings in other tertiary facilities within the country.^{4,5,7,8,16} Although these conditions were treatable, late presentation and indiscriminate use of over-the-

Paediatric Emergency Admissions, Mortalities, and Unmet Intensive Care Needs at a Tertiary Hospital in Southern Nigeria

counter antibiotics have contributed to delayed presentation and treatment with worsening prognosis at the time of presentation. Other Nigerian researchers have reported similar trends.^{7,8,16} It would be safe to imply that late presentation with more severe illness may be a significant contributor to the high case fatality reported, especially with cases with respiratory difficulties, requiring mechanical ventilation.

Only a small proportion of the children in need of intensive care were admitted into the intensive care unit. This was mainly due to the unavailability of space in the ICU and caregivers' inability to pay the admission deposit. These reflect the healthcare finance method in the study locale and the country in general, where out-of-pocket payment accounts for most of the payments made for healthcare services. Similarly, the mortality recorded in this study of patients admitted into ICU is markedly higher than reported from another centre in China.³⁰ Other Nigerian studies on the pattern of admission in the children's emergency room did not report any met or unmet ICU admissions and outcomes. The abysmally small number that survived in the ICU in this study may be due to late presentation into the unit and possibly sub-optimal medical care received prior to presentation at our facility. All these may have also negatively impacted survival. The absence of a designated Paediatric Intensive Care Unit (PICU) at the study locale may have greatly impacted the ease of admission compared to sharing an ICU with adult patients. Also, the lack of staff with paediatric critical care training at the main ICU may have adversely affected the quality of care and in turn, the clinical outcomes.

Conclusion

Infectious diseases are the leading causes of paediatric emergencies and the most typical causes of death in CHER, especially among under-five children in our facility. The high contribution of infectious disease to infant and

childhood mortality would suggest that health promotion and disease prevention programs must be further strengthened. The majority of the children requiring ICU care did not get it, but amongst those opportune, the majority eventually died. The need for a functional, well-manned and equipped Paediatric Intensive Care Unit (PICU) cannot be over-emphasized. More research is needed on the extent of met and unmet intensive care services at paediatric emergency rooms in tertiary hospitals in Nigeria. Healthcare policymakers should also consider the provision and efficient operation of Paediatric Intensive Care Units (PICUs) at tertiary health facilities in Nigeria.

Acknowledgement: The authors appreciate the support of the resident doctors and house officers in the unit in collecting data.

Authors' Contributions: E-UFE conceived the study and did the literature review with AC, OEG and AAO. All the authors analysed and interpreted the data, drafted the manuscript, revised it for sound intellectual contents and approved the final version of the manuscript.

Conflicts of Interest: None declared.

Funding Supports: The authors received no funding for the research and publication of this article.

Accepted for publication: 28th May 2024.

References

1. O'Malley PJ, Barata IA, Snow SK. Death of a child in the emergency department. *Pediatrics*. 2014;134(1):198–201. Doi: [10.1186/s12873-020-00352-9](https://doi.org/10.1186/s12873-020-00352-9)
2. Goal 3: Good health and well-being | Sustainable Development Goals | United Nations Development Programme [Internet]. [cited 2023 12th December]. Available from: <https://www.undp.org/sustainable-development-goals/good-health>
3. Bryce J, Boschi-Pinto C, Shibuya K, Black RE. WHO estimates of the causes of death in children. *Lancet* 2005; 26:365(9465):1147–52. DOI: [10.1016/S0140-6736\(05\)71877-8](https://doi.org/10.1016/S0140-6736(05)71877-8)
4. Diakparome MA, Obi JO. The Pattern of Paediatric Emergencies in the University of

- Benin Teaching Hospital. *Niger J Paediatr* 1980; 7:43-5.
5. Abhulimhen-Iyoha BI and Okolo AA. Morbidity and mortality of childhood illnesses at the emergency paediatric unit of the University of Benin Teaching Hospital, Benin City. *Niger J Paediatr* 2012;39(2):71–74.
<http://dx.doi.org/10.4314/njp.v39i2.7>
 6. Bilkisu GI, Aminu MS, Sunday OO, Bassey E, Smart A, Muyideen AB. Pattern of medical childhood morbidity and mortality in a new specialist hospital in Gusau, Nigeria. *Ann Nigerian Med* 2014;8:15-19. DOI:[10.4103/0331-3131.141024](https://doi.org/10.4103/0331-3131.141024)
 7. Duru C, Peterside O, Akimbami F. Pattern and outcome of admissions as seen in the Paediatric Emergency Ward of the Niger Delta University Teaching Hospital Bayelsa State, Nigeria. *Niger J Paediatr* 2013; 40:232-237.
<http://dx.doi.org/10.4314/njp.v40i3.6>
 8. Agbesanwa TA, Babatola AO, Fatunla OA, Ibrahim A, Aina FO, Ogundare EO, et al. Pattern of admissions and outcome in the children emergency department of a tertiary health institution in Southwestern Nigeria: A four-year review. *Afr J Emerg Med* 2023;13(2):45–51. DOI:[10.4314/ahs.v19i2.14](https://doi.org/10.4314/ahs.v19i2.14)
 9. Enyuma COA, Anah MU, Pousson A, Olorunfemi G, Ibisomi L, Abang BE, et al. Patterns of paediatric emergency admissions and predictors of prolonged hospital stay at the children emergency room, University of Calabar Teaching Hospital, Calabar, Nigeria. *Afr Health Sci*. 2019;19(2):1910–1923. DOI:[10.4314/ahs.v19i2.14](https://doi.org/10.4314/ahs.v19i2.14)
 10. Ezeonwu BU, Chima OU, Oguonu T, Ikefuna AN, Nwafor I. Morbidity and Mortality Pattern of Childhood Illnesses Seen at the Children Emergency Unit of Federal Medical Center, Asaba, Nigeria. *Ann Med Health Sci Res* 2014;4(3):239-244. DOI:[10.4103/2141-9248.141966](https://doi.org/10.4103/2141-9248.141966)
 11. Anyanwu OU, Ezeanosike OB, Ezeonu CT. Pattern and outcome of admissions at the children's emergency room at the Federal Teaching Hospital Abakaliki. *Afr J Med Health Sci* 2014;13:6-10. DOI: [10.4103/2384-5589.139435](https://doi.org/10.4103/2384-5589.139435).
 12. Frank-Briggs AI, Grange A. Disease pattern and childhood mortality. In: Azubuike JC, Nkanginieme KEO (2nd Ed.). *Paediatrics and Child Health in a Tropical Region*. African Educational Services, Owerri, 2007: 12-16.
 13. Ntekim BE, Ikpeme E. Common paediatric emergencies. In Azubike JC, Nkanginieme KEO (2nd Eds). *Paediatrics and Child Health in a Tropical Region*. African Educational Services, Owerri, 2007: 111-115
 14. Bassey EU, Ijezie E. Pediatric emergencies seen in a tertiary Hospital in Uyo, Akwa Ibom state of Nigeria: a two-year review. *Int J Sci Stud*. 2016;4(4):42–45. DOI: [10.4236/ojim.2024.142016](https://doi.org/10.4236/ojim.2024.142016)
 15. Onubogu UC, West BA, Azubogu US. The pattern of mortality among children hospitalized in the children's emergency ward of a single tertiary hospital in Nigeria. *Pediatr Emerg Med J* 2023;10(1):3-10. DOI: [10.22470/pemj.2022.00598](https://doi.org/10.22470/pemj.2022.00598)
 16. Njoku PU, Iloh GI, Dienye PO, Korie FC, Ahaiwe VC. Pattern of paediatric conditions seen in the children emergency room of an urban hospital in South-Eastern Nigeria. *GSC Bio Pharm Sci* 2020;30:13(3):174–180.
<https://doi.org/10.30574/gscbps.2020.13.3.0246>
 17. de Silva S. Enhancing child survival by bridging the gaps in emergency paediatrics. *Sri Lan J Child Health* 2013;42(3):115-128
<http://dx.doi.org/10.4038/sljch.v42i3.6013>
 18. UNICEF: 2013 statistical snapshot child mortality. www.childinfo.org. Accessed 28th March, 2024.
 19. CDC Childhood Injury Report: Patterns of Unintentional Injuries among 0-19-Year-Olds in the United States, 2000-2006. stacks.cdc.gov/view/cdc/5155. Accessed on 18th February 2015)
 20. Jofiro G, Jemal K, Beza L, Bacha HT. Prevalence and associated factors of pediatric emergency mortality at Tikur Anbessa specialized tertiary hospital: A 5-year retrospective case review study. *BMC*

Paediatric Emergency Admissions, Mortalities, and Unmet Intensive Care Needs at a Tertiary Hospital in Southern Nigeria

- Pediatr* 2018;18:316.
DOI: [10.1186/s12887-018-1287-4](https://doi.org/10.1186/s12887-018-1287-4)
21. Santhanam I, Pai M, Kasturi K, Radhamani MP. Mortality after admission in the pediatric emergency department: a prospective study from a referral children's hospital in southern India. *Pediatr Crit Care Med* 2002;3(4):358–363. Doi: [10.1097/00130478-200210000-00006](https://doi.org/10.1097/00130478-200210000-00006).
 22. Zhu CP, Wu XH, Liang YT, Ma WC, Ren L. The mortality of patients in a pediatric emergency department at a tertiary medical centre in China: An observational study. *World J Emerg Med*. 2015;6(3):212. <https://doi.org/10.5847/wjem.j.1920-8642.2015.03.009>
 23. Whitfill T, Auerbach M, Scherzer DJ, Shi J, Xiang H, Stanley RM. Emergency Care for Children in the United States: Epidemiology and trends over time. *J Emerg Med*. 2018;55(3):423–434. DOI: [10.1016/j.jemermed.2018.04.019](https://doi.org/10.1016/j.jemermed.2018.04.019)
 24. O'Meara M, Trethewey S. Managing paediatric death in the emergency department. *J Paediatr Child Health* 2016;52(2):164–167. DOI: [10.1111/jpc.12957](https://doi.org/10.1111/jpc.12957)
 25. Maniktala A, Pruitt C, Poirier M. Deaths in a non-trauma center pediatric emergency department: a ten-year experience. *J Emerg Med* 2002; 1(2)
 26. Lopez E, Udaondo J, Olabarri M, Martinez-Indart L, Benito J, Mintegi S. Mortality in Spanish pediatric emergency departments: a 5-year multicenter survey. *Eur J Emerg Med*. 2017;24(6):392–397. DOI: [10.1097/MEJ.0000000000000365](https://doi.org/10.1097/MEJ.0000000000000365)
 27. Menge I, Essamai F, Van Renken D, Anabwani G. Pediatric morbidity and mortality at Eldoret Hospital, Kenya. *East Afr Med J* 1995;72:165-169.
 28. Lahmini W, Bourrous M. Mortality at the pediatric emergency unit of the Mohammed VI Teaching Hospital of Marrakech. *BMC Emerg Med* 2020;20:57;2-6. DOI: [10.1186/s12873-020-00352-9](https://doi.org/10.1186/s12873-020-00352-9).
 29. Robison JA, Ahmad ZP, Nosek CA, Durand C, Namathanga A, Milazi R, et al. Decreased pediatric hospital mortality after an intervention to improve emergency care in Lilongwe, Malawi. *Pediatrics* 2012;130:676–82. DOI: [10.1542/peds.2012-0026](https://doi.org/10.1542/peds.2012-0026).
 30. Cui-ping Z, Xiao-hui W, Yu-ting L, Wen-cheng M, Lu R. The mortality of patients in a pediatric emergency department at a tertiary medical centre in China: An observational study. *World J Emerg Med* 2015;6(3): 212–16. DOI: [10.5847/wjem.j.1920-8642.2015.03.009](https://doi.org/10.5847/wjem.j.1920-8642.2015.03.009).
 31. WHO/ Emergency Triage Assessment and Treatment; ETAT. [http:// www.who.int/child_adolescent_health/documents/9241546875/en/](http://www.who.int/child_adolescent_health/documents/9241546875/en/) Accessed 9th April 2024
 32. Ibeziako SN, Ibekwe RC. Pattern and Outcome of Admissions in the Children's Emergency Room of the University of Nigeria Teaching Hospital, Enugu. *Niger J Paediatr*. 2022;29(4):103–7. DOI: [10.4314/njp.v29i4.12018](https://doi.org/10.4314/njp.v29i4.12018).
 33. Okoronkwo NC, Chapp-jumbo AUN. Pattern of Morbidity and Mortality of Childhood Illnesses at the Emergency Paediatric Unit of Abia State University Teaching Hospital, Aba, Nigeria. *East Cent Afr Med J* 2015;2:70-73.
 34. Fajolu IB, Egri-Okwaji MTC. Childhood Mortality in The Children Emergency Centre of the Lagos University Teaching Hospital. *Niger J Paediatr*. 2011;38(3):131-135. DOI: [10.4314/njp.v38i3.72270](https://doi.org/10.4314/njp.v38i3.72270)
 35. Fetuga B, Ogunlesi T, Adekanbi F, Olanrewaju D, Olowu A. Comparative analyses of childhood deaths in Shagamu. Implications for the fourth MDG. *South Afr J Child Health* 2007;1(3):106 -111.
 36. Ayoola OO, Orimadegun AE, Akinsola A.K, Osinusi k. A five-year review of childhood mortality at UCH Ibadan. *West Afr J Med*. 2005;24(2):175-179. Doi: [10.4314/wajm.v24i2.28192](https://doi.org/10.4314/wajm.v24i2.28192).
 37. Lesi FEA, Temiye EO, Epelle TGS. The changing pattern of childhood mortality in the children's emergency room of Lagos University Teaching Hospital after 20 years. *Niger Med J*. 2000;38(3):38-40.
 38. Wammanda RU, Alli FU. Conditions associated with risk of death within 24 hours of admission in Zaria, Nigeria. *Ann Afr Med* 2004;3(3):134-137.

39. Fagbule D, Joiner KT. Pattern of childhood mortality at the University of Ilorin Teaching Hospital. *Niger J Paediatr* 1987;14:1-5.
40. Adeyokunnu AA, Taiwo O, Antia AU. Childhood mortality among 22,255 consecutive admissions in the University College Hospital, Ibadan. *Niger J Paediatr* 1980;7:7-15.
41. Elusiyan JBE, Obiajunwa PO, Adejuyigbe EA, Olowu WA, Adeodu OO, Owa JA, *et al.* Pattern of Morbidity and Mortality among Children hospitalized at the Obafemi Awolowo University Teaching Hospital, Ile-Ife. *Niger J Paediatr* 2009;36: 22-28.
42. Ndu IK, Uleanya ND, Nwokoye IC, Edelu BO, Asinobi IN, Ekwochi U, *et al.* Pattern of Morbidity and Mortality at the Children Emergency Unit of Enugu State Teaching Hospital, Park Lane, Enugu. *J Exp Res* 2016;4(1):48–54.
43. Sa'ad YM, Hayatu A, Al-Mustapha II, Orahachi YM, Hauwa MU. Morbidity and mortality of childhood illnesses at the emergency pediatric unit of a tertiary hospital, north-eastern Nigeria. *Sahel Med J.* 2015; 18:1-3. DOI:10.4103/1118-8561.15215.
44. Kam-lun EH, Edmund ASN. Gender disparity in paediatric hospital admissions. *Ann Acad Med Singapore* 2006;35:882-888.
45. George IO, Alex-Hart BA, Frank-Briggs AI. Mortality Pattern in Children: A Hospital Based Study in Nigeria. *Int J Biomed Sci.* 2009; 5(4): 369–372.
46. Obi CC. Gender Differences in Educational Opportunities: The case of girl-child education in Nigeria. *Afr Econs Bus Rev* 2009;7:1-12.