

THE EFFECTS OF MENOPAUSE ON THE SERUM LIPID PROFILE OF NORMAL FEMALES OF SOUTH EAST NIGERIA

J. C. IGWEH¹, I. U. NWAGHA¹, J. M. OKARO²

¹*Department of Physiology, Faculty of Medicine, College of Medicine, University of Nigeria Enugu Campus, Enugu Nigeria.*

²*Department of Obstetrics and Gynaecology, Faculty of Medicine, University of Nigeria, Enugu Campus, Enugu Nigeria.
. Email: jcigweh@yahoo.com, Tel: +234 8033 247 168*

Summary: Dyslipidemia in menopause is a known feature in women, whether it leads to significant increase in the development of coronary heart disease (CHD) is still controversial, more so in our environment where little work has been done. The present study is aimed at comparing the level of total serum cholesterol, including its subunits in premenopausal with that of their postmenopausal counterparts. This is to enable us ascertain the relative risk of developing cardiovascular disease in postmenopausal women in our environment. One hundred and twenty six (126) apparently healthy, non-pregnant females (74 premenopausal and 52 postmenopausal) were recruited for the study. Serum total cholesterol and their subfractions- high-density lipoproteins (HDL), low-density lipoproteins (LDL), very low-density lipoproteins (VLDL) and triglycerides (TG) were estimated using enzymatic and established mathematical methods. There was no significant difference in the total serum cholesterol and triglyceride between the two groups. There was however, a significant reduction of HDL and VLDL in the postmenopausal group ($P < 0.005$) and a significant increase in the level of LDL in the postmenopausal group ($P < 0.005$). The elevated LDL and the reduction of cardio protective HDL and VLDL is an indication that menopause is an independent risk factor for developing cardiovascular disease in our environment.

Key Words: Menopause, Dyslipidemia, Cholesterol

Introduction

Cardiovascular disease is a leading cause of death among women in the developed world. In the United States, more than 500,000 women die of cardiovascular disease and about half are due to coronary artery disease (CAD) (Ariyo *et al*, 2002). Multiple risk factors have been identified as contributory to the development of CAD. These risk factors are important in both men and women and are present in both Caucasians and Africans. They include cigarette smoking, hypertension, Diabetes mellitus and hypercholesterolemia.

Hypercholesterolemia is a key factor in the pathophysiology of atherosclerosis (Igweh *et al*, 2003). Thus dyslipidemia can be modified by efforts aimed at cholesterol reduction (Fick *et al*, 1987, Kane *et al*, 1990, Davidson 1993). Studies have shown that women are at less risk of developing CAD than their male counterparts, but this is abolished after 60 years of age (Rich-Edward 1995 *et al*, Courderc *et al*, 1999). High levels of LDL and low levels of HDL are strongly associated with the risk of CAD (McNamara

et al 1992). Smaller LDL particles (LDL-III) are considered more atherogenic than larger more buoyant species because of their increased susceptibility to oxidation (Dejagar *et al*, 1993) and their increased residence time in plasma (Rainwater, 2000). Plasma triglycerides concentration also has a determinative influence on the concentration of small dense LDL particles in normal population (Dejagar *et al*, 1993, Mendelsohn, 1999).

After menopause, there is loss of ovarian function. This results in adverse changes in glucose and insulin metabolism, body fat distribution, coagulation, fibrinolysis and vascular endothelial dysfunction (Spencer *et al*, 1977). There is also derangement of lipoprotein profile independent of age (Bales, 2000).

A number of changes that occur in the lipid profile after menopause are associated with increased cardiovascular disease risk. Lack of estrogen is an essential factor in this mechanism. Apart from maintaining friendly lipid profile, estrogen changes the vascular

tone by increasing nitrous oxide production. It stabilizes the endothelial cells, enhances antioxidant effects and alters fibrinolytic protein (Taddec *et al*, 1996). All these are cardioprotective mechanisms, which are lost in menopause.

Dyslipidemia at menopause is not restricted to Caucasians. It is also evident in women of South Eastern Nigeria. These studies were aimed at determining the degree of dyslipidemia in the women of this region and relate it to the risk of developing cardiovascular diseases. It also attempted to suggest possible methods of reverting to lipid friendly status.

Materials and Methods

A group of 126 women, 74 premenopausal aged between 25-45 years and 52 postmenopausal aged between 55-70 years were studied. They were of the same social class and selected from workers and students of the University of Nigeria and the University of Nigeria Teaching Hospital (UNTH) community. Some of the postmenopausal women were relatives of the workers. They were randomly selected by a lucky dip of yes or no after an informed consent and ethical clearance from the relevant ethical committee of UNTH Enugu was obtained. Exclusion criteria include obesity, pregnancy, diabetes mellitus, hypertension, hormonal contraception and heavy exercise.

Fasting venous samples (10ml) were collected in heparinized bottles. This however, was done on the 7th day of the L.M.P. for the premenopausal group. Sample was centrifuged and plasma separated and stored in plastic tubes at 4°C.

Total cholesterol was measured using established enzymatic methods of Allain *et al* (1974) with the Randox cholesterol kit (Randox England). HDL-C was isolated by HDL-C precipitant method (Lopes-virella *et al*, 1977). LDL was calculated with a formula from total cholesterol and triglycerides (Friedwald, 1972). Triglyceride was isolated enzymatically as described by Henry (1991) and the expert panel on detection, evaluation and treatment of high blood cholesterol in adults (Expert Panel, 1988). VLDL was calculated using the formula $VLDL-C = TG/2.825$ mmol/L (Henry, 1991).

Statistical analysis was done using Excel. Showing the mean and standard deviation. Comparison of mean was by student t-test. Permission was obtained from the Faculty Ethics committee to carry out this experiment.

Results

The mean, Standard deviation and the P-values for total cholesterol, HDL-C, LDL-C VLDL-C and Triglycerides are shown in Table 1. There was no significant difference in the total cholesterol and the triglycerides between the two groups ($P>0.005$). However there was significant reduction in HDL and VLDL-C fraction in the postmenopausal group and a significant increase in the value of LDL-C ($P<0.005$).

Table 2 shows the HDL/LDL ratio between the two groups. Figs 1,2,3,4, and 5 represent comparisons of means between the two groups. The mean age for premenopausal women is 36.7 ± 4.74 and that for post menopausal women is 60.9 ± 9.9

Table 1: Showing Total plasma cholesterol and its subfractions in pre and postmenopausal women.

	Total Cholesterol	HDL	LDL	VLDL	TG
Premenopausal women(n-74)	4.30±0.68	1.37±0.020	2.7±0.74	0.28±0.04	1.12±0.43
Post menopausal women(n-56)	4.51±0.74	1.23±0.21	3.03±0.69	0.25±0.04	1.3±0.73
P.Values	0.117	0.005	0.039	0.004	0.137
Significance	$P>0.05$	$P<0.05$	$P<0.05$	$P<0.05$	$P>0.05$

Table 2: HDL/LDL Ratio in pre and postmenopausal women

	HDL	LDL	HDL/LDL Ratio
Premenopausal women	1.37	2.7	1:2
Post menopausal women	1.23	3.03	1:3

Fig. I: Mean Serum Cholesterol (Pre and post menopausal)




Fig III: Mean LDL-C (Pre and post menopausal)




Fig. II: Mean HDL-C (Pre and post menopausal)




Fig IV: Mean VLDL-C (Pre and post menopausal)





Fig. V: Mean TG (Pre and post menopausal)



Discussion

There are variations in lipid levels obtained in different individuals based on race, age, sex, obesity, exercise, smoking, alcohol, diet, diseases like hypertension, chronic liver and renal diseases (Gordon *et al*, 1987, Jarikre *et al*, 2000). However, in this study we tried to exclude the co founding variables and results obtained can be taken as the standard normal in our environment. The premenopausal values are similar to values obtained in other studies in different parts of the country (Ajose *et al*, 2002, Igweh *et al*, 2003).

However, there is paucity of literature in postmenopausal values in Nigeria, but studies on total cholesterol alone done in Ibadan showed no significant change (Otolorin *et al*, 1989) as was the case in our study.

Studies in some centers in developed countries showed higher values with dyslipidemia being taken as cholesterol > 240mg% (6.21mmol/l) and occurs in 25-33% of women (Grundy, 1994, Gordon *et al*, 1989). As a result of racial, genetic and technical differences, it has been suggested that our own cut off level should be adjusted to suit our peculiar circumstances. In determining the risk of cardiovascular disease, the absolute cholesterol value is not the most important factor; rather the concentrations of the various subclasses of cholesterol. In the present study there was no significant differences in the total cholesterol level but there was significant reduction in the cardio protective HDL-C and VLDL-C and significant increase in the arterosclerotic LDL-C. This is in agreement with findings in other studies (Jenson *et al*, 1990, Eder *et al*, 1982, Igweh *et al*, 2003). It has also been estimated that for any 1mg/dl (0.026mmol/ml) increase in HDL-C, there is a 3% decrease in risk of coronary artery disease and a 4.7% decrease in the risk of mortality from cardiovascular disease (Okonofua 1990). In this study this protective effect was lost by five fold. Again the HDL/LDL ratio was increased in the post menopausal group and it has been shown that HDL/LDL ratio is a significant predictor for the development of arterosclerosis both in Caucasians and Nigerians (Aina 1992, Park *et al*, 2000).

There is no doubt from this study that the changes that occur in the lipid profile after menopause is not friendly for the cardiovascular health of our women. It is generally believed that postmenopausal symptoms are less in our women than their Caucasian counterparts. This may not be true because several studies in the past have shown psychological, physical, biochemical,

hormonal and vasomotor parameters that are not of great variance with what is obtained elsewhere (Grundy, 1994, Mandel *et al*, 2002, Pirway *et al*, 2002). The problem is that the harsh climatic and poor socioeconomic environment overwhelms our women that they hardly ever complain about menopausal symptoms unless it is severe.

The elevated LDL and the reduction in the cardio protective HDL and VLDL is an indication that menopause is an independent risk factor for developing cardiovascular disease in our environment.

Several studies have shown the beneficial effects of hormonal replacement therapy on the lipid profile of menopausal women (Abbot *et al*, 1988, Stampfer *et al*, 1991, Stampfer, 1991). However, controversy exists as to whether these changes culminate to reduced risk of cardiovascular heart disease.

Observational studies over the years have touted the beneficial effects of hormone replacement therapy (HRT) in preventing coronary heart disease in post menopausal women (Hulley *et al*, 1998). More recent studies have however, cast some doubts on the beneficial effects of HRT especially in patients with established cardiovascular disease (Cheng, 2000). Further studies are needed in this area.

It should be noted that our post menopausal women have unfriendly lipid profile; it is thus important to note this and device means of correcting the dyslipidemia since the use of HRT and lipid lowering drug is still controversial. It is important to counsel on proper dietary, social and physical habits.

References

- Abbot, R. D., Wilson, PWF, Kannel, WB and Castelli, WP: (1988). High-density lipoprotein cholesterol, total cholesterol screening and myocardial infarction. The Framingham study. *Arterosclerosis* 8:207
- Aina, A.O. (1992). An investigation into climacterics in Nigerian women. *J.Med.Assoc. Thai.* 75(3) 168-72.
- Ajose, O. A., Fasubaa, O. B., Thomas, K. D., Bolodeoku, J. O. (2002). Serum lipids and lipoprotein cholesterol profile in pregnant Nigerian women. *J. Clin. Sci.* 2(1-2) 9-13.
- Allain, C. C., Poon, L. S., Chan, C. S. G., Richmond, W., Fu, P. C. (1974). Enzymatic determination of total serum cholesterol. *Clin. Chem.* 20: 470-475.
- Ariyo, A. A., Villablanca, A. C. Estrogen and Lipid: (2002). Can HRT, designer

- estrogens and phytoestrogens reduce cardiovascular risk markers after menopause? *Post grad. Med.* 111(1) 23-30.
- Bales, A. C. (2000). In search of lipid balance in older women; New studies raise questions about what works best. *Postgrad. Med.* 108 (7) : 57-72.
- Cheng, G. S. (2000). Cardiac events increased in the first 2 years of HRT. *Intern. Med. News* 33(9) 1-2.
- Couderc, R, Machi, M. (1999). Lipoprotein (a): risk factor for atherosclerotic vascular Disease important to take into account in practice. *Ann-Biol-Clin* 57(2):157-67.
- Davidson, M. H., (1993). Implications for the present and Direction for the future. *Am . J. Cardiol.* 71: 32B-36B.
- Dejagar, S. Bruckert, L., Chapman, M. J. (1993). Dense low density lipoprotein subspecies with diminished oxidative resistance predominate in combined hyperlipidemia. *J. Lipid Res.* 34; 295-308.
- Eder, H. A., Gidez, L. L. (1982). The Clinical significance of the plasma high density lipoprotein. *Med. Clin. North Am.* 66: 431-434.
- Expert panel on Detection, Evaluation and Treatment of high blood cholesterol in Adults: (1988). Report of the National Cholesterol Education Program Expert Panel on detection, evaluation, and treatment of high blood cholesterol in adults. *Arch. Intern. Med.* 148: 36.
- Fick, M. H., Elo, O., Haapa, K., Hheinonen, O. P., Heinsalmi, P., Helo, P., Huttunen, J. K., Kaitaniemi, P. Koshinen, V., Maenpaa, H., Malkonen, M., Manttari, M., Norola, S., Pasternack, A., Pikkarame, J., Roo, M., Sjoblom, T. and Nikkila, E. A. (1987). Helsinki heart study: Primary-Prevention trial with gemfibrozil in middle-aged men with dyslipidemia. Safety treatment changes in risk factors and incidence of coronary heart disease *N. Engl. J. Med.* 317- 1245.
- Friedwald, W. T., Levy, R. L., Fredrickson, D.S. (1972). Estimation of the concentration of low density lipoprotein cholesterol in plasma without use of the preparative ultracentrifuge *Clin., Chem.* 18: 499.
- Gordon, D. J., Trost, D. C., Hyde, J., Whaley, F. S., Hannan, P. J., Jacobs, D. R. and Ekelund, L. G: (1987). Seasonal cholesterol cycles: The lipid Research Clinics Coronary Primary Prevention Trial Placebo group. *Circulation.* 76:1224
- Gordon, D. J., Probstfield, J. L., Gramson, R. J. (1989). High density lipoprotein cholesterol and cardiovascular disease: Four prospective American Studies. *Circulation* 79(1)8-15.
- Grundy, S. M., (1994). Guidelines for cholesterol Management: Recommendations for the natural cholesterol Education Programs. Adult treatment Panel II. *Heart Dis. Stroke* 3(3) 123-7.
- Henry, J. B. (1991). Clinical Chemistry. In Clinical and Diagnosis Management by Laboratory Methods. 18th Edition, W.B. Saunders Company London 189-214.
- Hulley, S., Grady, D., Bush, T. (1998). The randomized trial of estrogen plus progestin for secondary prevention of coronary heart disease in postmenopausal women: Heart and Estrogen/progestin replacement study (HERS) Research group. *JAMA* (7) 605-13.
- Igweh, J. C., Aloamaka, C.P. (2003). HDL/LDL Ratio- A Significant Predisposition to the Onset of Atherosclerosis. *njhbs.* Vol. 2: (2) 78-82.
- Igweh, J. C., Nwafia, W.C., Ejezie, F. E. (2003.) Low Density Lipoprotein as a vehicle of Atherosclerosis. *Njhb.* 2(1) 7-11.
- Igweh, J.C., Aloamaka, C.P. (2003). Cholesterol Profile of Adults Resident in Eastern Nigeria. *O.J.Med.* 15(3&4) 46-50.
- Jarikre, A. E., Ola, F., Abbiyesuku, M. F., Oluwafomoju, I.O. (2000). Cross sectional study of plasma triglycerides and cholesterol levels in first and second trimesters of pregnancy in Lagos. *Nig. J. Int. Med.* 3(1): 1-3.
- Jensen, J, Nilas, L., Christiansen, C. (1990). Influence of menopause on serum lipid and lipoprotein. *Matruita* 12: 321-31.
- Kane, J. P., Malloy, M. J., Ports, T. A., Philips, N. R., Diolu, J. C., Havel, R. J. (1990). Regression of coronary atherosclerosis during treatment of familial hypercholesterolemia with combined drugs regimens . *JAMA* 264:3007-3012.
- Lopes-Virella, M. F., Stone, P., Ellis, S. (1977). Cholesterol determination in high density lipoprotein separated by three different methods. *Clin. Chem.* 23: 882
- Mandel, T., Purakayastha, S., Allauddin, M., Biswas, S., Nandi, J. (2002). Effects of hormonal replacement therapy on the lipid profile of postmenopausal women. *J. obstet. Gynae of India.* 52(2) 102-104.
- McNamara, J. R., Jenner, J. L., Li Z., Wilson, P. W., Schaefer, E. J. (1992). Changes in

- LDL Particle size is associated with change in plasma triglyceride concentration. *Arterioscler. Thromb.* 12: 1284-1290.
- Mendelsohn, M. E., Karas, R. H. (1999). The protective effects of estrogen on the cardiovascular system. *N Engl. J. med.* 340: 1801-1811.
- Okonofua, E. E., Lawal, A., Bamgbose, J. K.. (1990). Features of Menopause and Menopausal age in Nigerian women. *Int. J. Gynaecol. Obstet.* 31(4) 341-5.
- Otolorin, E. O., Adeyefa, I., Osotimehin, B. O. (1989). Clinical, hormonal and biochemical features of menopausal women in Ibadan Nigeria. *Afri. J. Med. Sci.* 18 (4) 251-255.
- Park, J. S., Jung, H. H. , Yang, W. S., Kim, S. B., Min, W. K., Chi, H. S. (2000). Effects of hormonal replacement therapy on lipid and homeostatic factors on postmenopausal ESRD patients. *Nephrol Dial Transplant.* 15: 1835-1840.
- Pirwany, R. I, Satter, N., Greer, A. I., Packard, C. J. (2002). Fleming R. Supraphysiological concentrations of estradiol in menopausal women given repeated implant therapy so no adversely affect lipid profile. *Human Reproduction.* 17(3) 825-829.
- Rainwater, D. L. (2000). Lipoprotein correlate of LDL particle size. *Atherosclerosis.* 148: 151-158.
- Rich-Edward, J. W., Manson, J. E., Hennekens, C. H. (1995). The Primary prevention of coronary heart disease in women. *N. Engl. J Med.* 332(20); 1758-1766.
- Spencer, C.P., Godsland, H., Stevenson, J. C. (1977). Is there a menopausal metabolic syndrome? *Gynecol. Endocrinol.* 11: 341-355.
- Stampfer, M. J., Colditz, G. A. (1991). Estrogen replacement therapy, and coronary heart disease: A quantitative assessment of the epidemiological evidence. *Prev. Med.* 20(1) 47-63.
- Stampfer, M. J., Colditz, G. A., Willet, W. C. (1991). Postmenopausal estrogen therapy and cardiovascular disease. Ten year follow up from Nurses Health Study. *N. Engl. J. Med.* 325(11) 956-62.
- Taddec, S., Virdis, A., Ghiadoni, L., Mattec, P., Sudano, I., Bernini, G. (1996). Menopause is associated with endothelial dysfunction in women. *Hypertension* 28: 576-582.

Received:22/3/2005

Accepted: 21/10/2005