

The tables and figures in this article were omitted when the article was published in the vol. 52, issue 1, 2011 of Nigerian Medical Journal. The error is highly regretted. The omissions are hereby corrected.

Editor-In-Chief, NMJ

Prevalence of Intestinal Parasites among Pupils in Rural North Eastern, Nigeria

Damen, J.G., ²Luka, J., ¹Biwan, E.I., ¹Lugos, M

ABSTRACT

Background: The study determined the prevalence of intestinal parasitism among pupils in rural schools (Almajiris) in Konduga local Government Area of Borno state. **Materials and Methods:** A total of 257 stool specimens were collected at random among pupils (Almajiris) in rural quranic schools; the stools were processed and examined both macroscopically and microscopically by concentration techniques. **Results:** The prevalence of intestinal parasitism among the Almajiris was 80.9%. The highest prevalence rate was 97.8% while the least prevalence was 67.4%. The 6-8 years age group had the highest prevalence of 85.7% while the least prevalence of 77.7% in the 13-16years age bracket. *Ascaris lumbricoides* had the highest prevalence of (19.1%) while *Trichuris trichiura* had the least prevalence of (3.5%). Thirteen pupils in the 5-8 years had multiple parasites; multiple parasitism also occurred in 22 pupils aged 9-12 years and in 11 pupils aged 13-16 years. **Conclusion:** There is a high prevalence rate of intestinal parasites with attendant risk of intestinal obstruction among the Almajiris in rural north eastern Nigeria.

KEYWORDS: Almajiris, intestinal parasites, intestinal obstruction

INTRODUCTION

Intestinal protozoa and faeco-oral transmitted helminths (STH) constitute major health problems, especially in the tropical and sub-tropical regions¹. They can be the cause of a wide spectrum of clinical problems ranging from apparently symptomless infections to life-threatening conditions such as intestinal obstruction as in *Ascaris* infestation,¹ failure to thrive as well as anaemia if not properly treated.²

The 'Almajiris' are children drawn from different parts of the northern Nigeria to learn and be trained in Islamic religion under the leadership of Mallams (Islamic instructors-/scholars). The study aimed to determine the prevalence of intestinal parasites among the Almajiris in a local government area in north eastern Nigeria.

MATERIALS AND METHODS

Approval and permission for this study was

¹Department of Medical laboratory Science University of Jos

²Federal School of Medical Laboratory, Jos

Corresponding Author

J.G. Damen. Department of Medical laboratory science. Faculty of Medical sciences University of Jos. P.M.B.2084 Jos damenjgd@yahoo.com

granted by the Chief Imam of Konduga Central Mosque and head (Mallam) of the Quranic schools and/or children's guardians before the commencement of the study.

Stool specimens were collected in a clean, wide mouth, screw capped, transparent, dry and disinfectant-free containers from 257 Almajiris who were chosen randomly in each of the Quranic schools studied between February 2006 to May 2006. The stool specimens were immediately examined in the laboratory of General Hospital located in the local government area.

Each of the stool specimens was examined both macroscopically and microscopically using saline preparation, Lugol's Iodine and Formal ether concentration techniques as described by Cheesbrough.³


The Almajiris that had intestinal parasites were treated with single dose of Albendazole or Praziquantel and or metronidazole for 5 days depending on the type of intestinal parasites seen in their stools.

RESULTS

Of the total 257 stool specimens examined, 208(80.9%) had intestinal parasites. The

distribution among the schools showed a prevalence range of 97.8-67.4% (figure 1). The age group of 6-8 years had the highest prevalence (85.7%) while the 13-16 years bracket had the least (77.7%) [Table 1].

Figure 1: Distribution of Intestinal parasites among Qu'ranic Schools in Konduga



KEY

- MGMF - Mallam Gani Modu Fewa
- H. Goni - Hassan Goni
- MGS - Mallam Garba Sabon gari
- MSG - Modu Selim Madarari.
- K-K - Kauri-Kaigamari

TABLE 1: Prevalence of intestinal parasites among Almajiris in Konduga in relation to age groups

Age groups (Years)	No Examined	No Positive	Prevalence (%)
5-8	56	48	85.7
9-12	107	87	81.3
13-16	94	73	77.7
Total	257	208	80.9

TABLE 2: Prevalence of individual intestinal parasites among the Almajiris in the Konduga; n=257

Parasites	No Positive	Prevalence (%)
Ascaris lumbricoides	49	19.1
Entamoeba histolytica	38	14.8
Entamoeba coli	33	12.8
Enterobius vermicularis	17	6.6
Giardia lamblia	23	8.9
Hookworm	37	14.4
Schistosoma mansoni	48	18.7
Trichuris trichiura	9	3.5
Total	254	98.8

Table 2 showed the distribution of intestinal parasites. *Ascaris lumbricoides* had the highest prevalence of (19.1%) while *Trichuris trichiura*

had the least prevalence of (3.5%). Thirteen pupils in the 5-8 years had multiple parasites; multiple parasitism also occurred in 22 pupils aged 9-12 years and in 11 pupils aged 13-16 years (See Table 3).

TABLE 3: Prevalence of Multiple intestinal parasites in Almajiris

Age Groups (YRS)	No Examined	No. Multiple parasitism
5-8	61	13
9-12	107	22
13-16	89	11
TOTAL	257	46.0

DISCUSSION

The overall prevalence rate of intestinal parasitism among 'Almajiris' was 80.9%. This prevalence is higher when compared with previous findings in south-west Nigeria (28%),⁴ south-eastern Nigeria (55.2%)⁵ and south- south Nigeria (67.2%)⁶. High prevalence of intestinal parasitic infestation is apt to occur in low socio economic condition, characterised by inadequate water supply and poor sanitary disposal of faeces⁶⁻⁸. The higher prevalence rate of intestinal parasitism in this study might be attributed to the poor hygienic practice and sanitary environment in the Almajiris live. The indiscriminate disposal of human wastes and unhygienic way of life might have been predisposing factors. The schools in this study were located in rural areas lacking portable water, proper system of refuse and human waste disposal. These may lead to a vicious cycle leading to worm infestation endemicity among these Almajiris as they come in contact with these wastes while playing on the playing ground.

The predominance of *Ascaris lumbricoides* in this study is in consonance with other reports.¹³ The danger with ascariasis is that it is intimately related with intestinal obstruction and malnutrition in children. The prevalence of *Entamoeba histolytica* and *Schistosoma mansoni* was higher in this study compared with that of adult population¹⁰⁻¹³.

One consequence of intestinal parasite infestation is risk of mechanical intestinal obstruction, especially in times of massive infestation. Mechanical Intestinal obstruction may be caused by a bolus (ball) of parasites and

volvulus due to parasitic load and/or neurotoxins they excrete (Ascaris) causing contractions of the bowel.^{14,15} Massive de-worming of children with anthelmintic medications especially carries this risk of intestinal obstruction with a poorer outcome.^{16,17}

The study also showed that children within the 6-10 years age group had the highest prevalence of intestinal parasites, closely followed by the 11-15 years age groups. A similar trend was reported by Higgs¹⁸, and Jombo¹⁹ in which the bulk of parasitic infestation occurred in the 8-15years age group. Again the prevalence of intestinal parasites decreased with age in this study. This observation has been alluded to in another report²⁰. This inverse relationship between the age and the prevalence of intestinal parasites might be due to higher level of awareness and good hygienic practice in the older age groups.

The 46% polyparasitism/multiple parasitism rate in this study was higher compared with 20.5% rate reported elsewhere²⁰, but agrees with others^{18, 21}. Therefore, while treating children with intestinal parasites, it is advisable to use broad spectrum or multi-agent drug combinations because of this multiple parasitism susceptibility in children.

This study showed a high prevalence of intestinal parasitism among the Almajiris in the rural north eastern Nigeria. It is suggested that children with intestinal parasites should be treated periodically using broad spectrum or multi-agent drug combinations because of the multiple parasitism susceptibility in children. Public enlightenment and emphasis on personal hygiene and clean environment may be necessary in the prevention and control of parasitic infections among children in rural areas.

ACKNOWLEDGMENT

The authors are grateful to the Chief Imam of Konduga Central Mosque for given us permission for the study. The cooperation of the Mallams of the Quranic schools is highly appreciated. We appreciate the financial support of Lafaith Medical and Diagnostics laboratories, Jos, for treatment of infested subjects.

REFERENCE

1. Savioli L, Bundy DAP, Tomkins A. Intestinal parasitic

- infections: a soluble public health problem. *Trans R Soc Trop Med Hyg* 1992; 86: 353-354.
2. Glickman,LT, Camara,AO, Glickman,NW, Macabe, GP Nematodes intestinal parasites in children in rural Guinea, Africa: Prevalence and relationship to Geophagia *International Journal of Epidemiology* 1999 28:169-174
3. Cheesbrough M. *Medical Parasitology. Medical Laboratory Manual or Tropical Countries* Sheck Wah Tong Printing press ltd Hong Kong low price edition (1991).
4. Salako A.A. Effects of portable water availability on intestinal parasitism among rural school children with sewage disposal facilities in the Mjidum and Owutu sub- urban community of Lagos state. *Nigeria Medical practioner* 2001; 39: 3-4.
5. Agwu N. A., Incidence of intestinal helminthiasis in school children in Aba urban city, Abia state, Nigeria, *International Journal, Environmental Health and Human Development*. 2001; 1: 47-51.
6. Meremikwu M.M., Antia-Obong O.E., Asindi A.A., Ejezie G.C. Prevalence and Intensity of intestinal Helminthiasis in Pre-School children of peasant farmers in Calabar, Nigeria. *Nigerian Journal of Medicine*. 1995; 2: 40-44
7. Rajeswari B., Sinnaih B., Husseini H. Socioeconomic factors associated with intestinal parasites among children living in Gombak Malaysia Asia Pac. Health 1994; 7 (1): 21-25.
8. Al-Agha R., Teodorescu I. Intestinal parasites infections and anaemia in primary schools children in G29 Gonerorates Palestine *Roum- Arch-Microbiol- Immunol* 2000; 59 (1-2): 131-143
9. Nobel and Nobel. The prevalence and intensity of *Ascaris lumbricoides* infection in Muslim Children from Northern Bangladesh. *Trans. Royal Soc. Trop. Med. Hygiene* 1994; 77 (5): 702-706.
10. Okpala I. A survey of the incidence of intestinal parasites among Government Workers in Lagos, Nigeria. *West Africa Medical Journal* 1961; 10: 148-157.
11. Roche J. and Bento A. Prevalence of intestinal parasites infection with special reference to *Entamoeba histolytica* on the Island of Bioko (Equatorial Guinea) *Am. Trop. Med. Hygiene* 1999; 60 (2): 257-262.
12. Luka S.A., Ajogi I., Umoh J.U. Helminthosis among primary schools in Lere local Government Area Kaduna state, Nigeria. *The Nigerian Journal of parasitology*. 2000; 21: 109-116.
13. Eneanya C.I., Anikwue C. A school based intestinal helminthiasis programme in Nigeria: Perceptions, attitude and acceptability to community members. *Nigerian Journal of parasitology*. 2005; 26: 55-60.
14. Correa Antúnez M.I., Serrano Calleja Pimentel Leo, J.J., Sanjuán Rodríguez S. "Bezoar Gastrico" *Cir peditr* 2001; 14: 82-84
15. Montiel-Jarquín A., Carrillo-Ríos C., Flores-Flores J. *Ascaridiasis vesicular a sociada a hepatitis aguda. Manejo conservador. Cir Ciruj* 2003; 71: 314-318
16. Kamiya T., Justiano M., Durán A., Uechi C.

Prevalence of Intestinal Parasites among Pupils in Rural North Eastern, Nigeria

- Biliopancreatic ascariasis: endoscopic approach. *J Gastroenterol* 2002; 37(Suppl 13): 97-99
17. Chiarpenello J. Actualización: Infecciones por helmintos. *Evid. actual. práct. ambul.* 2004; 7: 178-181
 18. Higgs D. A., Jenkwis P, Human intestinal parasites in areas of Indonesia. *Annals of Tropical medicine and parasitology* 1984; 78: 637-641.
 19. Jombo G.T.A., Egah D.Z., Akuson J.T., Mbaawuga E.M. Human intestinal parasitism in a rural settlement of Northern Nigeria, A Survey. *Nigeria Medicalpractioner* 2007; 1 / 2: 11-15.
 20. Person, V; Ahmed, F. Gebre, Medhim, M. Relationship between Vitamin A, Iron status and Helminthiasis in Bangladeshi school children. *Public health nutrition* 2000; 3: 83-89.
 21. Ashford R. W., Craig P.S., Oppehiam S.T. Polyparasitism on the Kenya Coast spatral heterogeneity in parasitic distribution. *Ann. Trop. Med. and parasitol* 1993; 87 (3): 283-293.