

Psychosocial Impact of disclosure of HIV Serostatus in heterosexual relationship at the Lagos University Teaching Hospital, Nigeria

Adeyemo Titilope A,¹ Adediran A.¹ Umeh C.², Akinbami. A.³ Unigwe O.⁴ Akanmu A. S.¹

ABSTRACT

Objectives: To determine the rate and pattern of disclosure and non disclosure of HIV serostatus among people living with HIV and the psychosocial impact of disclosure. **Methods:** Participants were drawn from the adult HIV clinic at the Lagos University Teaching Hospital, Lagos, Nigeria. An interviewer-administered questionnaire was used to collect data from consenting participants that included socio-demographic information, pattern and reason for non-disclosure and the possible consequences of disclosure of HIV serostatus to their sexual partners. **Results:** Four hundred and ninety nine respondents with a mean age of 37.3 ± 9.6 years were recruited into the study. There were 157 males and 342 females. Majority of the participants were married (62%) and belonged to the low socio-economic class. Overall 61.5% (307 of 499) had disclosed their status to sex partner(s). Gender, social class and length of year of diagnosis were not associated with disclosure but number of sexual partners was strongly associated with non disclosure. $P=0.0063$. The most common reason for non-disclosure was fear of rejection (65%). Majority (96.7%) of those who disclosed their status had no regret and majority (81.1%) of those who had not disclosed had protected sex. After counseling, only 18.8% (36 of 192) of those that had not disclosed thought that the counseling had helped them overcome the fear of disclosure and were willing to disclose. **Conclusion:** Many people would disclose their HIV serostatus to sex partner(s). Protected sex (through the use of condom) is widely accepted in our setting.

Keywords: HIV serostatus disclosure, rejection, stigmatization, 'confidant assisted counseling'

INTRODUCTION

HIV/AIDS is ravaging sub-Saharan Africa, with greater than 60% of all HIV infections in the world occurring in this region.^{1,2} To date, few empirical studies have examined the sexual relationships and sexual behaviors of people living with HIV/AIDS (PLWHA) in Africa. In a study from Uganda, 47% of HIV-positive men and 21% of HIV-positive women were sexually active before initiating ARV treatment, with 45% of these people reporting unprotected inter-

course in a 3-month period.³

Not disclosing HIV status to sex partners is probably the product of multiple factors. [4, 5]

Some people living with HIV infection may generally conceal their HIV status from people in their lives, not just their sex partners. In addition, people who do not disclose their HIV status may have had adverse experiences related to previous disclosures, including loss of social support, rejection, loss of employment, violent reactions and other forms of discrimination. Finally, people who fail to disclose their HIV status may merely lack a sense of being able to effectively disclose their HIV status, especially to their sex partners.

Disclosure of HIV status is a planned and selective behavior which responds to the person's balance of potential risks and benefits of secrecy and disclosure¹. It might also be considered to be an expression of responsibility towards a spouse or sex partners⁶. Disclosure to others, lovers, family or friends, has been shown

¹. Department of Haematology and Blood Transfusion,

². Department of Psychiatry, Faculty of Clinical Sciences, College of Medicine, University of Lagos, Nigeria.

³. Department of Haematology and Blood Transfusion, Lagos State University College of Medicine, Ikeja, Nigeria.

⁴. Counseling Unit, AIDS Prevention Initiative in Nigeria (APIN), Lagos University Teaching Hospital Site, Lagos, Nigeria.

Correspondence: Akinbami Akinsegun.

Department of Haematology and Blood Transfusion, Lagos State University, College of Medicine, Ikeja
Email: ajoke_clinic@yahoo.co.uk Tel: +234 802 306-4925

Psychosocial Impact of disclosure of HIV Serostatus in heterosexual relationship at the L.U.T.H.

to be a potent stressor, as individuals living with HIV/AIDS might fear negative reactions such as blame, rejection or violence^{7,8,9}.

Obviously, progression of HIV might precipitate disclosure due to visible signs of disease or an urgent need for emotional support¹⁰. However, as HAART has caused HIV infection to become a chronic condition, this situation might have become less frequent, at least in countries with accessible treatment.

Among the various determinants of disclosure behavior, cultural factors have been identified, both regarding level and pattern of disclosure. Friends appear to be closer confidants than immediate families among gay men, while relatives might be chosen as confidants over spouses among HIV+ African women¹¹.

Fear of being a burden to the immediate family, or stigma associated with HIV infection, might force persons living with HIV/AIDS (PLWHA) to keep their disease secret from their social network, as has been shown in Asian populations^{12,13}.

Since the advent of HIV pandemics, disclosure of patient serostatus to sexual partners has been the concern of HIV caregivers. Overall rates of disclosure vary between studies, but many studies indicate that disclosure does not occur in a substantial majority of sex partnerships, perhaps 20% to 50% in developed countries [14, 15]. This is however different in developing countries where rate of disclosure is said to be lower. McNiel et al reported a disclosure of 24.7% among spouses and 18.8% among sexual partners¹⁶.

Though married spouses are more likely to disclose than casual sexual partners but for fear of rejection, non-disclosure is expected to be common amongst stable relationships that may lead to marriage. The most common reason for non-disclosure among married couples was fear of separation/divorce.¹⁷

Previous studies have shown that, one, not disclosing positive HIV serostatus to at least some sexual partners is common, [18-20]. Two, HIV-positive persons are more likely to disclose their status to steady partners than to non steady partners,²¹⁻²³ and three, unprotected sex without disclosure occurs within both casual and steady type of partnerships.¹⁸⁻²²

The purpose of this study was to determine the rate, pattern and factors associated with non

disclosure of HIV serostatus among people living with HIV.

MATERIALS AND METHODS

A cross-sectional survey was carried out at the HIV clinic of the Lagos University Teaching Hospital after obtaining ethical clearance from the institution review board. Eligibility criteria were; being 18 or older and in a heterosexual relationship(s). Patients with severe physical or mental impairment (as assessed by their attending physician), and those with other sexual orientation (eg men sex men, women sex women and bisexuals) were excluded from the study.

Subjects were randomly selected during their regular scheduled appointment. Questionnaire administrations took place at the hospital clinics and were administered in English by trained interviewers. The questionnaire covered a range of social issues including history of HIV testing and treatment, employment, income and living conditions, parenthood, sexuality and sexual relationships, social network and disclosure.

After the administration of the questionnaire, those that indicated they had not disclosed had individual counseling sessions on the benefits of disclosure and thereafter asked about the willingness to disclose.

Disclosure was defined as the patient him/herself telling his/her sexual partner/-spouse in heterosexual relationships about positive HIV serostatus.

Four hundred and ninety nine participants were included in the statistical analysis and data was subjected to descriptive analysis using Epi Info version 3.5.1 (2008). Test of significance was set at $P < 0.05$.

RESULTS

Four hundred and ninety nine participants were enrolled into the study consisting of 342 females and 157 males with a mean and median age of 37.3 ± 9.6 and 36 years respectively.

Though all the participants were African, there was a tribal diversity with about 44.4% being Igbos, 28.8% Yoruba, 4.2 % Hausa and 22.6% were from other Nigerian tribes. Majority (62%) of participants were married, 19.8% were single, 12% were widow, 3.2% were divorced and the rest 3% were separated. Many of the participants had formal education (secondary

Adeyemo Titilope A,¹ Adediran A,¹ Umeh C,² Akinbami A,³ Unigwe O,⁴ Akanmu A. S.¹

42.2%, tertiary 33.8% and primary 20.4%) while very few (3.6%) had no formal education. Of the 499 participants, 307 (61.5%) had disclosed their HIV status to their spouses/sexual partners while 192 had not. Many of the married participants (87.4%) had disclosed their HIV status to their partners while only a few (19%, 36 of 189) of the singles in sexual relationships had disclosed their status.

Table 1: Age group, marital status and HIV serostatus disclosure

Age group (years)	Disclosure (%)	Non disclosure (%)	Total
	<i>Married</i>		
20	0 (0.0)	0 (0.0)	0
21 – 30	63 (23.2)	11 (28.2)	74
31 – 40	122 (45.0)	17 (43.6)	139
41 – 50	65 (24.0)	9(23.1)	74
51 – 60	17 (6.3)	1 (2.6)	18
>60	4 (1.5)	1 (2.6)	5
	<i>Single</i>		
20	0 (0.0)	4 (2.6)	4
21 -30	15 (41.7)	41 (29.4)	56
31 -40	14 (38.9)	53 (34.6)	67
41 -50	7 (19.4)	33 (21.6)	40
51 -60	0 (0.0)	16 (10.5)	16
>60	0 (0.0)	6 (3.9)	6
Total	307	192	499

Majority of both married and singles that had disclosed are in the same age range 31-40 years (44.8% and 35.4% respectively). (Table 1)

Many of those that had disclosed (married 44.5%; and single 38.6%) had completed secondary school education and a few in both instances (married 2.9%; single 4.8%) had no formal education. The probability that a male participant will disclose his HIV serostatus, whether married or singles (married 89.1%; single 19.1%) is not significantly higher than that of a female participant disclosing her status (married 86.5%; single 19%). $P = 0.63$

Though majority of those that are single whether they had disclose or not had single sex partners, those with multiple sex partners are more likely not to disclose. $P = 0.0063$ (Table 2).

Majority, 96.7% (297 of 307) of those that had disclosed their status had no regret for disclosing and had not suffered rejection or any other adverse event for disclosing.

Reasons given for not disclosing include fear of rejection (65%); loss of intimacy (32.5%); stigmatization 21.3%); loss of economic support

(15%) and threat to personal well being (12.5%).

Protected sex is found to be common among the sexually active participants. Of the 192 who had not disclosed their status, only 53 (27.6%) claimed they were sexually active. Out of this, 43 (81.1%) claimed they use condom regularly and consistently while 10 (18.9%) do not use condom or any form of protection.

After counseling on the benefit of disclosure, 59 (30.7%) of 192 who had not disclosed their status thought that counseling had helped them overcome the fear of not wanting to disclose but only 18.8% (36 of 192) was still willing to disclose.

DISCUSSIONS

The belief that disclosure of HIV status to sexual partners will lead to rejection and stigmatization appears not to be true as majority of participants in the study expressed no regret for disclosing their status.

Although declining somewhat over the past few years, the levels of stigma and discrimination against people with HIV/AIDS still remains high in Nigeria especially in some communities least affected by HIV/AIDS. However, the outcome of disclosure to sexual partners, relations and friends is expected to be different from that of disclosure to private employers and colleagues in Nigeria where enforcements of labour rights of individuals are poorly coordinated.

Disclosure of HIV status to sex partners is therefore less likely to occur when a person has experienced adverse outcomes from previous disclosures for non sexual partners.

For couples, spouse/steady partner appears to be the first and the most reliable confidant as many of the couples interviewed confessed to disclosing to their spouses before friends and family members. This finding differs from observations among pregnant Tanzanian HIV+ women who were more likely to tell their status to a female relative when it was concealed from their spouse²⁴.

Sexual transmission risk behaviours were not found in this study in people who had not disclosed their HIV status to sex partners. This may suggest a successful campaign on the use condom at our centre. That this finding does not agree with what was found by many authors in other parts of the world may however suggest

Psychosocial Impact of disclosure of HIV Serostatus in heterosexual relationship at the L.U.T.H.

insincerity among our respondents. For example, Simbayi et al²⁵ found a close association between having not disclosed HIV status to sex partners and engaging in practices with high risk of HIV transmission in PLWHA in South Africa. And in another study conducted in two United States cities, Stein MD et al²⁶ found a high rates of non disclosure and the low rates of condom use, and concluded that sexual partners of HIV-infected persons continue to be at high risk for HIV transmission.

In this study, we found that HIV-related fear of rejection, loss of intimacy and stigmatization that are associated with not disclosing HIV status to sexual partners may not be overcome by conventional counseling techniques. We also found that unprotected intercourse and not disclosing HIV status to partners was found in multiple partnership relationship.

Even though counseling on the benefit of disclosure seems to help, 30.7% overcame the fear of not wanting to disclose their status, that only 18.8% of those that agreed that it has helped were willing to disclose after counseling has highlighted the fact that a new approach is necessary to complement the present counseling methods. An approach we recommend is 'confidant assisted counseling' in which a trusted family member or a friend is present with the patient in the counseling room and this confidant continues with the counseling after the patient has left the clinic. The counselor is able to monitor progress made through this trusted confidant.

In conclusion, a higher rate of disclosure when compared with similar studies was found in our centre. We also found a low level of sexual transmission risk among our non-disclosure group and that nondisclosure of HIV status is common among those with multiple sexual partners. Lastly, we recommend 'confidant assisted counseling' as a new counseling approach to support present counseling methods.

ACKNOWLEDGEMENT

Grateful to Mrs .F. N. Saliu for coordinating the secretarial duties.

REFERENCES

1. UNAIDS/WHO. AIDS epidemic update, December 2005. Joint United Nations Programme on HIV/AIDS 2005.

2. Shisana O, Rehle T, Simbayi L, et al. South African National HIV Prevalence, Incidence, Behaviour and Communication Survey. Cape Town: Human Sciences Research Council Press, 2005.
3. Bunnell R, Ekwaru J, Solberg P, et al. Changes in sexual behavior and risk of HIV transmission after antiretroviral therapy and prevention interventions in rural Uganda. *AIDS* 2006;20:85–92.
4. Sullivan KM. Male self-disclosure of HIV positive serostatus to sex partners: a review of the literature. *J Assoc Nurses AIDS Care* 2005;16:33–65.
5. L C Simbayi, S C Kalichman, A Strebel, A Cloete, N Henda, A Mqeketo Disclosure of HIV status to sex partners and sexual risk behaviours among HIV-positive men and women, Cape Town, South Africa. *Sex Transm Infect* 2007; 83:29-34 doi:10.1136/sti.2006.019893
6. Kim Bouillon, France Lert, Rémi Sitta, Annie Schmaus, Bruno Spire, and Rosemary Dray-Spira. Factors correlated with disclosure of HIV infection in the French Antilles and French Guiana: results from the ANRS-EN13-VESPA-DFA Study. *AIDS*. 2007 January; 21 Suppl 1: 889-894
7. Holt R, Court P, Vedhara K, Nott KH, Holmes J, Snow MH. The role of disclosure in coping with HIV infection. *AIDS Care*. 1998;10(1):49–60.
8. Kalichman SC, DiMarco M, Austin J, Luke W, DiFonzo K. Stress, social support, and HIV-status disclosure to family and friends among HIV-positive men and women. *J Behav Med*. 2003;26(4):315–332.
9. Medley A, Garcia-Moreno C, McGill S, Maman S. Rates, barriers and outcomes of HIV serostatus disclosure among women in developing countries: implications for prevention of mother-to-child transmission programmes. *Bull World Health Organ*. 2004;82(4):299–307. Review.
10. Serovich JM. A test of two HIV disclosure theories. *AIDS Education and Prevention*. 2001;13(4): 355–64.
11. Antelman G, Smith Fawzi MC, Kaaya S, Mbwambo J, Msamanga GI, Hunter DJ, et al. Predictors of HIV-1 serostatus disclosure: a prospective study among HIV-infected pregnant women in Dar es Salaam, Tanzania. *AIDS*. 2001;28(15):1865–74.
12. Yoshioka MR, Schustack A. Disclosure of HIV status: cultural issues of Asian patients. *AIDS Patient Care and STDs*. 2001;15(2):77–82.
13. Chandra PS, Deepthivarma S, Manjula V. Disclosure of HIV infection in South India: patterns, reasons, reactions. *AIDS Care*. 2003;15(2):207–215.
14. Marks G, Burris S, Peterman TA. Reducing sexual transmission of HIV from those who know they are infected: The need for personal and collective responsibility. *AIDS* 1999; 13:297-306.
15. Niccolai LM, Dorst D, Myers L, Kissinger PJ. Disclosure of HIV status to sexual partners: Predictors and temporal patterns. *Sex Transm Dis* 1999; 26:281-285.
16. McNeil JM, Mberesero F, Kilonzo G. Is care and support associated with preventive behaviour

Adeyemo Titilope A,¹ Adediran A,¹ Umeh C,² Akinbami A,³ Unigwe O,⁴ Akanmu A. S.¹

- among people with HIV? *AIDS Care* 1999; 11: 539-546
17. Daniel OJ, Falola RL, Ogundahunsi OA, Ogun SA, Odusoga OL, Salako AA, Oladapo OT, Ogunfowora OB, Osho OB, Adenekan AA, Oluwole FA, Kolapo KO, Boyle BA; Self-disclosure of HIV status among PLWHA on HAART in Sagamu, Nigeria. *Int Conf AIDS*. 2004 Jul 11-16; 15: abstract no. ThPeD7777.
 18. Marks G, Richardson JL, Maldonado N. Self-disclosure of HIV infection to sexual partners. *Am J Public Health*. 1991;81:1321-1322.
 19. Mansergh G, Marks G, Simoni JM. Self-disclosure of HIV infection among men who vary in time since seropositive diagnosis and symptomatic status. *AIDS*. 1995;9:639-644.
 20. Simoni JM, Mason HR, Marks G, Ruiz MS, Reed D, Richardson JL. Women's self-disclosure of HIV infection: rates, reasons, and reactions. *J Consult Clin Psychol*. 1995;63:474-478.
 21. Stein MD, Freedberg KA, Sullivan LM, et al. Sexual ethics. Disclosure of HIV-positive status to partners. *Arch Intern Med*. 1998;158:253-257.
 22. Wolitski RJ, Rietmeijer CA, Goldbaum GM, Wilson RM. HIV serostatus disclosure among gay and bisexual men in four American cities: general patterns and relation to sexual practices. *AIDS Care*. 1998;10:599-610.
 23. Niccolai LM, Dorst D, Myers L, Kissinger PJ. Disclosure of HIV status to sexual partners: predictors and temporal patterns. *Sex Transm Dis*. 1999;26:281-285.
 24. Antelman G, Smith Fawzi MC, Kaaya S, Mbwambo J, Msamanga GI, Hunter DJ, et al. Predictors of HIV-1 serostatus disclosure: a prospective study among HIV-infected pregnant women in Dar es Salaam, Tanzania. *AIDS*. 2001;28(15):1865-74.
 25. Simbayi LC, Kalichman SC, Strebel A, Cloete A, Henda N, Mqeketo A. *Sex Transm Infect* 2007;83:29-34 doi:10.1136/sti.2006.019893
 26. Stein MD, Freedberg KA; Sullivan LM; Savetsky J; Levenson SM; Hingson R; Samet JH. Disclosure of HIV-positive status to partners. *Arch Intern Med* 1998 Feb 9 158 253-257.

