



Molecular And Serological Detection Of Newcastle Disease Virus In Live-Bird Markets, Jos, Plateau State In Nigeria.

Chinyere C.N¹., Mpkuma N¹., Meseko C.A¹., Shittu I¹., Okwor E.C.², EzemaW.S²., Nwosuh C¹.

¹National Veterinary Research Institute, Vom Plateau State, ²University of Nigeria, Nsukka. *Corresponding author: Email: chinoscnc@gmail.com; Tel No: +2348030726007

ABSTRACT

Newcastle disease (ND) is a highly infectious viral disease of birds caused by the Newcastle disease virus (NDV) and it has been reported in domestic birds in Nigeria. Waterfowls and village poultry in live bird markets (LBM) acts as reservoirs, potentially reintroducing the virus to commercial poultry. This study aims at molecular and serological detection of NDV at live bird markets in Plateau State, Nigeria. A cross sectional analysis involved 309 pooled cloacal and tracheal swabs over three months were analyzed by reverse transcriptase polymerase chain reaction (RT-PCR), RT-qPCR, virus isolation and haemagglutination and haemagglutination-inhibition test (HA &HI). Virus isolation was attempted in 9 to 11 days old specific antibody negative (SAN) embryonated chicken eggs and fifteen samples showed haemagglutination. Subsequent tests confirmed nine of these as NDV positive through haemagglutination and haemagglutination inhibition (HI) tests. Conventional RT-PCR and RT-qPCR further validated five of the nine NDV positive isolates. The phylogenetic analysis of partial F gene nucleotide sequences revealed that all three isolates belonged to class II genotype XIV.2_XIVb. This finding underscores the persistent threat of NDV to local poultry, necessitating comprehensive virological surveillance to understand, isolate and characterize the virus in Nigeria. Therefore, monitoring for emerging lineages and sub-lineages in Nigeria birds is crucial for safeguarding commercial poultry production.

Keywords: Local chicken, Live bird market, Newcastle disease virus, Phylogenetic tree, Nigeria.

INTRODUCTION

Newcastle disease (ND) is a highly contagious and lethal viral disease that affects all bird

species. The virus responsible is Newcastle disease virus (NDV), an enveloped virus with a single stranded negative sense RNA genome that

is linear and non-segmented. It belongs to the family *Paramyxoviridae* in the subfamily *Avulavirinae*, genus *Orthoavulavirus* (ICTV, 2018, Dimitrov *et al.*, 2019; Ferreira *et al.*, 2021). According to the "rule of six" by Calain and Roux, 1993 which says that the viral RNA replication will be efficient only when the RNA nucleotide length is divisible by six, the RNA genome of NDV is approximately 15 kb and codes for six major proteins: nucleoprotein (NP), matrix protein (M), fusion protein (F), haemagglutinin neuraminidase (HN), phosphoprotein (P), and large RNA polymerase (L), in the order 3'-NP-P-M-F-HN-L-5' (Kolakofsky *et al.*, 2005). The disease affects a large number of hosts, about 241 species belonging to 27 orders, out of the 50 known orders of birds (Madadger *et al.*, 2013). Several wild birds of various ages as well as chickens, turkeys, ducks, pigeons, guinea fowl, and Japanese quail are among the more often impacted species (Nanthakumar *et al.*, 2000; Zhang *et al.*, 2011). Ducks are generally considered as natural NDV reservoirs, and even for NDV strains that are fatal to chickens, they exhibit little to no clinical symptoms following infection (Liu *et al.*, 2010; Stanislawek *et al.*, 2022). A virulent NDV infection can be fatal to chickens, leading to a large reduction in egg production and a high death rate. One of the main issues still plaguing established or emerging poultry industries in many nations, including Nigeria, is the disease.

Newcastle disease virus is classified into three major pathotypes based on the clinical signs induced in the infected chicken: Velogenic (highly virulent), mesogenic (intermediate virulent) and avirulent (lentogenic strain) (Lamb and Kolakofsky, 2001). Studies comparing the

deduced amino-acid sequence at the cleavage site of the NDV, showed that virulent viruses usually have the motif ¹¹²R/K-R-Q-K/R-R-F¹¹⁷ and avirulent viruses have ¹¹²G/E-K/R-Q-G/E-R-L¹¹⁷ (Dortmans *et al.*, 2009). Clinical signs of this disease in affected birds may vary widely and are dependent on factors like the virus strain, host species, age, immune status, environmental stress and concurrent infection. In chickens, the disease may vary from sudden death with 100% mortality to subclinical disease. Due to the enormous financial losses linked to the disease, particularly in chickens and turkeys, it is a serious threat to the poultry industries and has a global distribution (Miller *et al.*, 2010).

NDVs can be divided into classes I and II according to the phylogenetic analysis of the F gene, which has been done in a number of studies (Diel *et al.*, 2012; Samal *et al.*, 2012). Class II has at least 18 genotypes identified, compared to Class I which has nine (Snoeck *et al.*, 2013; Dimitrov *et al.*, 2016). Major factors influencing NDV pathogenicity include the cleavability of the fusion protein precursor (F0) and the presence of several basic residues in the fusion protein cleavage site (Martin-Garcia *et al.*, 2012). In Africa, velogenic NDVs have been isolated from sick and seemingly healthy poultry (Damena *et al.*, 2016; Molini *et al.*, 2017). Because of the poultry industry's explosive growth, high stocking densities, and inadequate biosecurity protocols, which encouraged the disease's endemicity and spread, NDV has been reported to be endemic in Nigeria (Okwor and Eze, 2010). Even though Nigeria experiences annual outbreaks of the NDV, little is known about the current spread of the virus in some of the regions in the country.

In Nigeria, NDV was first recorded in the 1950's (Hill *et al.*, 1953) but the molecular characterization of the Nigerian isolate was just recently available. Seven NDV genotype I, II, IV, VI, XIV, XVII, and XVIII members have all been found in the country (Snoeck *et al.*, 2013; Shittu *et al.*, 2016; Bello *et al.*, 2018; Welch *et al.*, 2019). The mainstays of NDV prevention and control are the stringent implementation of biosecurity protocols and extensive immunization campaigns, both of which have been effectively implemented for many years across the globe (Alexander, 2000). Even in vaccinated birds, the high frequency of NDV infection is not only associated with inadequate immunization or immunological suppression; it may also result from viral mutation that modifies the virus genomic sequence, changing its biological characteristics and virulence (Kattenbelt *et al.*, 2006). According to the growing number of reports of ND outbreaks in Nigerian flocks vaccinated against the virus, birds may not be entirely protected by the conventional vaccinations or vaccine failure may have resulted in more virulent new NDV variants. Despite research on NDV in Nigeria, little is known about the genetic makeup of the viruses that causes the disease to be endemic. Thus, the purpose of our research is to assess the characteristics of ND viruses that were isolated from local poultry in the northern and central regions of Plateau State, Nigeria.

MATERIALS AND METHOD

Study Area

Plateau State can be found in between Lat. 08° 24'N and longitude 008° 32' and 010° 38' E. It is situated in the tropical zone, with a temperature of 22°C and mean minimum low temperature of

18°C. The range of the mean yearly rainfall is 131.75 to 146 cm and the highest rainfall is recorded during the wet season months of July and August. Temperatures appear highest between the months of March and April (National Bureau of Statistics, 2009). It has a population of approximately 3.5 million people (NADIS, 2006a). Jos is divided into three LGAs namely Jos North, Jos South and Jos East (National Bureau of Statistics, 2009) and also into three senatorial districts or zones that comprises of Northern, Central and Southern zones. The people of Plateau State are predominantly farmers. The estimated number of poultry in Plateau State is eight million (Livestock population figures, FMARD, 2018, Animal husbandry Department, Federal Ministry of Agriculture). Because of the comparatively cool climate in Plateau State, poultry farming flourishes there. This research was conducted in major live-bird markets (LBMs) in the Northern and Central zones, specifically Yankaji old and Mangu LBMs. The State was chosen because of the fact that some farms in the state had experienced repeated outbreaks of Newcastle disease.

Figure. 1: Map of Plateau State displaying the research area (Chinyere C.N).

Sample Collection and Transportation

A multi-stage random sampling of local birds including chicken, ducks, turkey and geese brought to the LBM in Jos metropolis were sampled for this study. A total of 309 pooled cloaca and trachea swabs samples were collected. Each bird was sampled by inserting a polyester tipped swab stick on the cloaca and oropharyngeal cavities. Both samples were combined and placed in a 1.8ml cryovial containing 1 ml of viral

transport medium (VTM) containing penicillin, streptomycin, gentamicin and amphotericin B. The samples collected were transported in a cool box with ice packs to the Influenza Laboratory of National Veterinary Research Institute, Vom, Plateau State, Nigeria. The swabs were kept at -80°C until analyzed.

Virus isolation

Virus isolation was performed by inoculating 9-11 day old embryonated chicken eggs that are specific-antibody-negative (SAN) with a supernatant fluid of the swab specimens and incubated for five days for a period of 24hours interval. Egg inoculation, incubation, candling and virus harvesting were conducted in accordance with the OIE Manual of Standards for Diagnostic Tests and Vaccines (OIE, 2012). At the end of the incubation period, allantoic fluid (ALF) was harvested from the eggs and screened for Haemagglutination (HA) test. The HA positive ALF were further subjected to heamagglutination-inhibition (HI) test and RT-PCR for confirmation. Virus stocks were stored at -70°C until further used.

Extraction of viral RNA

The genomic viral RNA was extracted from the swabs using the QIAamp viral RNA mini kit (QIAGEN, Hilden, Germany) according to the manufacture's protocol and was stored at -80°C until use.

RT-qPCR

The samples were screened for L gene using RT-qPCR for the detection of avian paramyxovirus type 1 using Qiagen One-step RT-qPCR kit according to the manufacturer's instructions. The Reverse and forward primers sequence for the

NDV-L gene ND-F - 5'-GAG CTA ATG AAC ATT CTT TC -3', ND-R- 5'-AAT AGG CGG ACC ACA TC TG - 3', Probe L Pro MGB 6FAM 5'-CCA ATC AAC TTC CC -3', Probe L Pro MGB2 VIC 5'-AAT AGT GTA TGA CAA CAC -3' respectively were used. The L gene segment was reverse transcribed and amplified using applied biosystem (AB) thermocycler rotogen PCR. Reverse transcription was achieved by incubation with reverse transcriptase at 50°C for 20min and initial PCR activation 95°C for 15min. RT-qPCR was performed by denaturing cDNA at 94°C for 45sec, annealing at 50°C for 45sec. The program was repeated for 40 cycles.

RT-PCR for FUSION (F) GENE

Samples that were positive for NDV L gene were further amplified for F gene using a Qiagen One-step RT-PCR kit according to the manufacturer's instructions. The reverse and forward primers sequence for the NDV-F 5296F- 5'-ATTGGTAGCGGCTTGATCACTG-3' and NDV-F 6295R 5'-CGTTCTACCCGTGTACTGCTCTTT-3' respectively was used. The F gene segment was reverse transcribed and amplified using Applied Biosystem thermocycler PCR platform which amplified a 1000bp fragment of the NDV F gene. Reverse transcription was achieved by incubation with reverse transcriptase at 50°C for 30min and initial PCR activation 94°C for 15min. RT-PCR was performed by denaturing cDNA at 94°C for 30sec, annealing at 55°C for 1min and extending the fragment at 68°C for 90sec. The above program was repeated for 40 cycles with a final extension at 68°C for 10 min and PCR products maintained at 4°C.

Agarose gel electrophoresis

The PCR products were separated in 1.5% agarose gel in TBE buffer stained with ethidium bromide and compared with molecular marker (100bp). The gel was visualized on UV trans-illuminator a Biostep dark hood DH-40/50 imaging system (Germany).

SEQUENCING AND PHYLOGENETIC ANALYSIS

Positive PCR products were purified with QIAquick PCR purification kit (Qiagen, Germany) and sent to a commercial sequencing company Macrogen Inc, Netherlands for sequencing using ABI3730XL sequencer (Applied Biosystems) in both forward and reverse directions. Thereafter, consensus sequences were generated through CAP3 alignment editor and a phylogenetic tree was constructed using Molecular Evolutionary Genetic Analysis MEGA X. Genotype nomenclature according to Dimitrov *et al.*, 2019 were used to identify the genotype XIV.2_XIVb.

GenBank accession number

NDV Sequences obtained from this study were added to the GenBank database under accession numbers OK072691-OK072693.

Phylogenetic analysis

The partial F gene nucleotide sequences from the NDV isolates in this study were subjected to phylogenetic analysis and compared with those from classes I and II that were retrieved from the NCBI database. The results revealed that the three isolates all belong to class II genotype XIV.2_XIVb (**Figure 2**).

RESULTS

The conventional RT-PCR and RT-qPCR confirmed that 5 out of 15 (33%) viruses isolated from 309 pooled tracheal and cloacal swabs were NDV. Nine out of the 15 (60%) harvested were positive by HI. Three positive isolates (two from the Yankaji live-bird market and one from Mangu live-birds market) were sequenced to confirm their clade.

Fig. 2: Phylogenetic tree

The maximum likelihood phylogenetic tree of the NDV isolates from this study, along with other published sequences obtained from GenBank, is displayed in Figure 2. The Tamura-Nei model and the Maximum Likelihood method were used to infer the evolutionary history (Tamura and Nei, 1993). The displayed tree has the highest log likelihood (-32194.90). There were 39 nucleotide sequences in this analysis. Codon positions included were 1st+2nd+3rd+Noncoding. The final dataset contained 15232 positions in total. Evolutionary analyses were carried out in MEGA X (Kumar *et al.*, 2018).

DISCUSSION

The endemicity of Newcastle Disease Virus (NDV) in Nigeria is highlighted by the discovery of the virus in seemingly healthy local poultry in this study conducted at the live-bird market in Plateau State. It may also buttress the assertion that local poultry are relatively more resistant to NDV with a lower mortality rate than commercial poultry.

Phylogenetic analysis indicated that the NDV isolates in this study belong to the class II genotype XIV.2_XIVb. Members of this genotype are widely distributed in West and Central African countries, of which Nigeria is a part, with high virulence and mainly isolated from domestic birds. Similar findings have been reported in the

North-Central States of Benue and Kogi, as well as in the North-West states of Nigeria, including Sokoto, Kaduna, and Jigawa. Others are Taraba and Yobe (North- East), Lagos State (South-West) (Bello *et al.*, 2018), (Owoya *et al.*, 2020). Because of the local diversity of the genotype XIV subgenotype XIV.1_XIVa and XIV.2_XIVb, they are also known as regional NDV genotype. Similar results was obtained by Catherine *et al.*, 2019 where she reported a wild distribution XIV.2_XIVb in domestic birds such as pigeon, chicken, duck, turkey etc from 2002 -2015 from various parts of the country.

Phylogenetically, the genotype XIV.2_XIVb from this study clusters around other Nigerian XIV.2_XIVb genotype isolated in 2008 and 2011 with GenBank accession number HF969187, HF969145, HF969149, HF969210 and KT948996 (Fig.2). Elsewhere, genotype XIV has been reported in Burkina Faso with Burkina Faso/2415-580/2008 (BF/08), South Africa with Chicken/South Africa/08100426/2008 (SA/08), and Niger with Niger/1377-7/2006 (Niger/06) (Susta *et al.*, 2015).

Strains from LBMs and outbreaks from commercial chicken farms shared sequence similarities (genotype XIVb) (Welch *et al.*, 2019). Previous reports have noted some observed sequence similarity between strains recovered from outbreaks in commercial and backyard poultry farms in Jos, Plateau State, and those from LBMs in Sokoto and Kano states (Solomon *et al.*, 2012). This may be due to horizontal spread by live poultry vendors from rural settlements and live bird markets to commercial and backyard poultry farms. Because different species of birds from different places interact uncontrollably, live-bird markets have remained a major factor in the

spread of NDV (Choi *et al.*, 2005; Abah *et al.*, 2017). However, given the mindset of farmers who replace their LBM stock with existing poultry on their farms, the possibility of LBM spillover to backyard and commercial farms cannot be completely ruled out.

CONCLUSION

As part of an early warning system, Nigeria must maintain active, flock-based serological, virological, and molecular surveillance for NDV in order to detect the virus quickly and establish long-term control. The combination of these different methods ensures prompt detection of the virus and needs a proactive measures for the prevention of major outbreaks.

ACKNOWLEDGEMENT

Many thanks to Dr. David Shamaki, former Director/Chief Executive, National Veterinary Research Institute, Vom; Dr. Maryam Muhammad, Director/Chief Executive, National Veterinary Research Institute; Dr. T.M Joannis, Former H.O.D, Regional Laboratory for Animal influenza and other TAD's, NVRI, Dr. Clement A. Meseko, Assistant Director/HoD, Infectious and TAD's, NVRI and all the entire staff of Regional Laboratory for Animal influenza and other TAD's.

REFERENCES

- Abah H.O., Assam A., Abdu P.A. (2017). Newcastle disease and biosecurity practices in live bird markets in Benue State, Nigeria. *Nigerian Veterinary Journal* 38(1):13-25.
- Aldous E. W., and Alexander, D. J. (2001) Detection and differentiation of Newcastle

- disease virus (avian paramyxovirus type 1. *Avian Pathol.* 30:117–128.
- Alexander D.J. (2000). Newcastle disease and other avian paramyxo viruses. *Revue scientifique et technique International Office of Epizootics.* 19 (2): 443-462. <https://doi.org/10.20506/rst.19.2.1231>.
- Bello M.B., Yusoff K.M., Ideris A., Hair-Bejo M., Peeters B.P., Jibril A.H., Tambuwal F.M., Omar A.R (2018) Genotype diversity of Newcastle disease virus in Nigeria: disease control challenges and future outlook. *Advances in Virology.* <https://doi.org/10.1155/2018/6097291>.
- Calain P. and Roux L. (1993). The rule of six, a basic feature for efficient replication of Sendai virus defective interfering RNA. *J.Virol.* 67 (1993), pp. 4822-4830.
- Catherine N.W., Ismaila S., Celia A., Ponman S., Dimitrov K.M., Tonya L.T., Dawn W.C., Iryna V.G., Mesoko C.A., Ibu J.O., Dorcas A.G., Joannis T.M., and Claudio I.A. (2019). Genomic comparison of Newcastle disease viruses isolated in Nigeria between 2002 and 2015 reveals circulation of highly diverse genotypes and spillover into wild birds. *Archives of Virology* 164:2031-2047.
- Cattoli G., Fusaro A., and Monne I. (2010). Emergence of a new genetic lineage of Newcastle disease virus in West and Central Africa: implications for diagnosis and control. *Veterinary Microbiology* 142:168–176.
- Choi K.S., Lee, E.K., Jeon, W.J. and Kwon, J.H. (2010). Antigenic and immunogenic investigation of the virulence motif of the Newcastle disease virus fusion protein. *J. Vet. Sci.*, 11 (3): 205–211.
- Damena D., Fusaro A., and Sombo M. (2016). Characterization of Newcastle disease virus isolates obtained from outbreak cases in commercial chickens and wild pigeons in Ethiopia. *Springer Plus* 5(1):476.
- Diel D.G., Susta L., Cardenas G.S., Killian M. L., Brown C. C. and Miller P. J. (2012). Complete genome and clinicopathological characterization of a virulent Newcastle disease virus isolate from South America. *Journal of Clinical Microbiology* 50:378-387.
- Dimitrov K.M., Afonso C.L., Yu Q., and Miller P.J. (2016). Newcastle disease vaccines - A solved problem or a continuous challenge? *Veterinary Microbiology* 206:126-136.
- Dimitrov K.M., Abolnik C., Afonso C.L., Albina E., Bahl J., Berg M., Briand F.X., Brown I.H., Choi K.S., Chvala I., Diel D.G., Durr P.A., Ferreira H.L., Fusaro A., Gil P., Goujgoulova G.V., Grund C., Hicks J.T., Joannis T.M., Torchetti M.K., Kolosov S., Lambrecht B., Lewis N.S., Liu H, Liu H, McCullough S, Miller PJ, Monne I, Muller CP, Munir M, Reischak D, Sabra M, Samal S.K., Servan de Almeida R., Shittu I., Snoeck C.J., Suarez D.L., Van Borm S., Wang Z., Wong F.Y.K. (2019). Updated unified phylogenetic classification system and revised nomenclature for Newcastle disease virus. *Infect Genet Evol.*;

- 74:103917. doi: 10.1016/j.meegid.2019.103917. Epub 2019 Jun 11. PMID: 31200111; PMCID: PMC6876278.
- Dortmans, J.C., Koch, G., Rottier, P.J. and Peeters, B.P. (2009). Alternative method for rapid diagnosis and pathotyping. Virulence of pigeon paramyxovirus type 1 does not always correlate with the cleavability of its fusion protein. *J. Gen. Virol.*, 90 (Pt 11): 2746–275.
- Ferreira H.L., Miller P.J., Suarez D.L. (2021). Protection against different genotypes of Newcastle Disease Viruses (NDV) afforded by an adenovirus-vectored fusion protein and live NDV vaccines in chickens. *Vaccines*, 9 (2021), p. 182.
- Owoya, H.A., Ismaila, S., Paul, A.A., and Chinwe, J.A. (2020). Molecular characterization and phylogenetic analysis of Newcastle disease virus isolated from poultry in North Central States of Nigeria. *Journal of Veterinary Medicine and Animal Health*, 12(2), 20-26. <https://doi.org/10.5897/jvmah2019.0825>.
- Hill D.H., Davis O.S., Wilde J.K.H. (1953) Newcastle disease in Nigeria. *British Veterinary Journal* 109(9):381-385.
- International Committee on Taxonomy of Viruses (ICTV) (2018). Order Mononegavirales; family Paramyxoviridae; genus Avulavirus. *Virus taxonomy: 2018a release. EC 50*, Washington, DC, July 2018; email ratification February 2019.
- Kattenbelt J. A., Steven M.P., and Gould A.R. (2006). Sequence variation in the Newcastle disease virus genome. *Virus Research* 116:168-184.
- Kolakofsky, D., Roux, L., Garcin, D., and Ruigrok, R.W. (2005) Paramyxovirus mRNA editing, the “rule of six” and error catastrophe: a hypothesis. *J Gen Virol* 86:1869-1877.
- Kumar S., Stecher G., Li M., Knyaz C., and Tamura K. (2018). MEGA X: Molecular Evolutionary Genetics Analysis across computing platforms. *Molecular Biology and Evolution* 35:1547-1549.
- Lamb R.A., and Kolakofsky, D. (2001) *Paramyxoviridae: The Viruses and Their Replication*. 4th edition. Hagerstown: Lippincott Williams and Wilkins.
- Liu H., Chen F., and Zhao Y. (2010). Genomic characterization of the first class I Newcastle disease virus isolated from the mainland of China. *Virus Genes* 40(3):365-371.
- Madadger O., Karimi V., Nazaktabar A., Kazemimanesh M., Ghafari M.M., Dezfouli S.M.A., and Hojjati P. (2013). A study of Newcastle disease virus obtained from exotic caged birds in Tehran between 2009 and 2010. *Avian Pathology* 42(1):27-31.
- Martin-Garcia F., Mendieta-Moreno J.I., Mendieta J., and Gomez-Puertas P (2012). Molecular dynamics analysis of conformational change of paramyxovirus F protein during the initial steps of membrane fusion. *Biochemical and Biophysical Research Communications* 420(1):42-47.
- Miller P.J., Afonso C.L., Spackman E., Scott M.A., Pedersen J.C., Senne D.A., Brown

- J.D., Fuller, C.M., Uhart M.M., Karesh W.B., Brown I.H., Alexander D.J. and Swayne D.E. (2010). Evidence for a New Avian Paramyxovirus Serotype-10 Detected in Rockhopper Penguins from the Falkland Islands. *J. Virol.*, 84 (21): 11496-11504.
- Molini U., Aikukutu G., Khaiseb S., Cattoli G., and Dundon W. G. (2017). First genetic characterization of Newcastle disease viruses from Namibia: identification of a novel VIIk subgenotype. *Archives of Virology* 162(8):2427-2431.
- NADIS (National Animal Disease Information and Surveillance (NADIS) (2006a). Avian Influenza: A *bulletin of The Federal Department of Livestock and Pest Control Services. Pan African Program for the Control of Epizootics*. Bulletin No 1. 15th October 2006.
- Nanthakumar T., Kataria R.S., Tiwari A.K., Butchiah G., and Kataria J.M. (2000). Pathotyping of Newcastle disease viruses by RT-PCR and restriction enzyme analysis. *Journal of Veterinary Research and Communication* 24:275-286.
- OIE. (2012). Manual of diagnostic tests and vaccines for terrestrial animals mammals, birds and bees. *Newcastle Dis.* 1:555–574
- Okwor E.C and Eze D.C. (2010). The Annual prevalence of Newcastle disease in commercial chickens reared in south eastern savannah zone of Nigeria. *Research Journal of Poultry Science* 3:23-26.
- Samal S., Khattar S. K., Kumar S., Collins P. L., and Samal S. K. (2012). Coordinate deletion of N-glycans from the heptad repeats of the fusion F protein of Newcastle disease virus yields a hyperfusogenic virus with increased replication, virulence, and immunogenicity. *Journal of Virology* 86:2501-2511.
- Shittu I., Joannis T.M., Odaibo G.N., Olaleye O.D. (2016). Newcastle disease in Nigeria: epizootiology and current knowledge of circulating genotypes. *Virus disease*. 27(4):329-339. doi: 10.1007/s13337-016-0344-6. Epub 2016 Sep 8. PMID: 28004012; PMCID: PMC5142595.
- Snoeck C.J., Owoade A.A., Couacy-Hymann E., Alkali B.R., Okwen M.P., and Adeyanju A.T. (2013). High genetic diversity of Newcastle disease virus in poultry in West and Central Africa: co-circulation of genotype XIV and newly defined genotypes XVII and XVIII. *Journal of Clinical Microbiology* 51(7): 2250-2260.
- Stanislawek W.L., Wilks V.R., Meers J., Horner G.W., and Alexander D.J. (2002) Avianparamyxoviruses and influenza viruses isolated from mallard ducks (*Anas platyrhynchos*) in New Zealand. *Archives of Virology* 147:1287-1302.
- Susta, L., Jones, M.E.B., Cattoli, G., Cardenas-Garcia, S., Miller, P.J., Brown, C.C., and Afonso, C.L. (2015). Pathologic characterization of Genotypes XIV and XVII Newcastle Disease Viruses and Efficacy of classical vaccination on specific pathogen-free Birds. *Veterinary Pathology*, 52(1), 120-131. <http://doi.org/10.1177/0300985814521247>.

