

Submitted: 11/07/2015

Accepted: 02/11/2015

Published: 20/11/2015

Cadmium affects the mitochondrial viability and the acid soluble thiols concentration in liver, kidney, heart and gills of *Ancistrus brevifilis* (Eigenmann, 1920)

P. Velasquez-Vottelerd*, Y. Anton and R. Salazar-Lugo

Laboratorio de proteínas e inmunotoxicidad, Postgrado de Biología Aplicada, Núcleo de Sucre, Universidad de Oriente, Cumaná, Venezuela

Abstract

The freshwater fish *Ancistrus brevifilis*, which is found in Venezuelan rivers, is considered a potential sentinel fish in ecotoxicological studies. The cadmium (Cd) effect on the mitochondrial viability (MV) and acid soluble thiols levels (AST) in *A. brevifilis* tissues (liver, kidney, heart, and gill) was evaluated. Forty-two fish with similar sizes and weights were randomly selected, of which 7 fish (with their respective replicate) were exposed for 7 and 30 days to a Cd sublethal concentration (0.1 mg.l⁻¹). We determined the MV through a Janus Green B colorimetric assay and we obtained the concentration of AST by Ellman's method. Mitochondrial viability decreased in fish exposed to Cd for 30 days with the liver being the most affected tissue. We also detected a significant decrease in AST levels was in fishes exposed to Cd for 7 days in liver and kidney tissues; these results suggests that AST levels are elevated in some tissues may act as cytoprotective and adaptive alternative mechanism related to the ROS detoxification, maintenance redox status and mitochondrial viability. Organ-specific variations were observed in both assays. We conclude that the Cd exposure effect on AST levels and MV, vary across fish tissues and is related to the exposure duration, the molecule dynamics in different tissues, the organism and environmental conditions.

Keywords: *Ancistrus brevifilis*, Cadmium, Soluble thiols, Janus Green B, Mitochondrial viability.

Introduction

In Venezuela, there has been a disturbing increase in the ambient concentrations of heavy metals in watersheds (Bifano and Mogollón, 1995; Hermoso and Márquez, 2005; Corona, 2013; Mora *et al.*, 2013). In particular, cadmium (Cd) has been reported in moderate concentrations (Hermoso and Márquez, 2005; Salazar, 2009). Cd is known as cytotoxic and immunotoxic metal and is potentially carcinogenic (Salazar *et al.*, 2009). As a result of the increased levels of Cd in the aquatic environment, the bioaccumulation of the metal in organisms has enhanced, especially in fishes (Souid *et al.*, 2013; Salazar-Lugo *et al.*, 2014; Perera *et al.*, 2015). The accumulation of Cd in tissues has been reported from: *Aglyptodactylus laticeps* (Marcano and Troconis, 2001), *Hoplias malabaricus*, *Prochilodus reticulatus* (Vanegas, 2003), *Colossoma macropomum* (Hernández, 2005), *Centropomus decimalis* (Márquez *et al.*, 2008). The liver, kidney and gills are the target organs of metal uptake (Perera *et al.*, 2015; Sabullah *et al.*, 2015). The heart, is an aerobic organ rich in mitochondria and several studies have suggested that the heart is very sensitive to Cd toxicity (Wang *et al.*, 2004; Soares *et al.*, 2008; Akpakpan and Akpanyung, 2014).

In cells, the mitochondria is a target for Cd. There are numerous reports in fish documenting the effect of Cd

on these organelles including ATP synthesis depression, free radicals generation, lipid peroxidation, and mitochondrial membrane depolarization (Sokolova, 2004; Atli and Canli, 2008; Padmini and Usha, 2011). To counter the oxidative damage derived from Cd accumulation in different tissues, organisms increase their antioxidant molecules, such as metallothioneins (MT), glutathione (GSH), and other molecules rich in thiol groups. These molecules constitute the first defense line of the cell against oxidative damage (Salazar *et al.*, 2009; Sevcikova *et al.*, 2011; Dorts *et al.*, 2012; Hatem *et al.*, 2014).

The Guaraguara (*Ancistrus* sp.), is a native fish widely distributed in freshwater ecosystems of northeastern Venezuela, the Amazon River and other rivers in South America. It is the fourth most common fish species, in relative abundance, in the Manzanares River, Venezuela (Ruiz *et al.*, 2005). It is a nocturnal benthic fish, which lives in the bottom of rivers and feeds mainly on algae and invertebrates. These traits lead to an increased uptake of pollutants and make *A. brevifilis* a sentinel organism to detect pollution in these ecosystems (Lárez, 2011).

The biomonitoring of freshwater fish species can be used to assess the consequences of chemical pollution by heavy metals in aquatic environments. These ecotoxicological studies are needed to increase

*Corresponding Author: Patricia Velásquez Vottelerd. Laboratorio de Proteínas e inmunotoxicidad, Postgrado de Biología aplicada, Cerro del Medio, Universidad de Oriente, Cumaná, Venezuela 6101. E-mail: patriciavelasquezv@gmail.com

our understanding of the impact humans have on the environment. The objective of this study was to evaluate the effects of Cd exposure on the mitochondrial viability (MV) and acid soluble thiols levels (AST) in *A. brevifilis*.

Materials and Methods

Fish maintenance

We collected *A. brevifilis* specimens (n=84) from the Manzanares River, Yaque sector (10°12'17.58" N and 63°53'21.08" W). (11.39±1.69 m and 12.21±3.01 g) were used. All fish handling and maintenance procedures were done according to bioethics code of FONACIT (2011). We transported fish in black bags with aerated water; to the Immunotoxicity Protein Laboratory, Universidad de Oriente, Sucre State. We maintained the fish for 15 days in holding tanks with continuously aerated and flowing de-chlorinated tap water at a constant temperature of 27±3 °C. We fed fish *ad libitum* with fish food, protein content 45%. In order to simulate the natural conditions these fish live in, we reduced artificial illumination by surrounding the aquariums with black bags. We maintained the pH at 7.7±0.25, temperatures (27±3 °C), oxygen level at 3.8 mg.l⁻¹, and total hardness at 106 mg.l⁻¹ throughout all experiments.

Lethal mean concentration (LC₅₀) determinations

We determined the LC₅₀ of Cd at 96 using a static test with water exchange every 24 h and daily feeding as previously described (Peltier and Weber, 1985), but with a few modifications. We exposed fishes, in groups of four (with replicates), to 0.01, 0.5, 5, 10, 20, and 30 mg.l⁻¹ Cd for 96 h in 12 l plastic aquariums. In addition, we included a control group, which was kept in water free of Cd. We recorded mortality was at 12, 24, 36, and 96 h post exposure to Cd. We calculated the LC₅₀ using the Logic method (Weber, 1993) using LC₅₀ software. *A. brevifilis* LC₅₀ was 11.81 mg.l⁻¹, 0.1 mg.l⁻¹ Cd from CdCl₂ was selected for sublethal assay (<1 % LC₅₀ 96 h).

Bioassay with 0.1 mg.l⁻¹ Cd

We tested the Cd levels in *A. brevifilis* tissues prior to any sub lethal experiments. The Cd concentrations were 0.19 (±0.005 mg.g⁻¹) which are under the permitted limits for fish (Osman and Kloas, 2010).

We randomly selected 42 fishes for the assay. We made sure all fish were a similar weight and size. We exposed seven groups of fishes to sublethal 0.1 mg.l⁻¹ Cd for 7 and 30 days. We included 14 fishes as a control unexposed to Cd. We continuously aerated each 40 l aquarium and we maintained the same physical, and chemical characteristics of the water as those for the laboratory acclimation. The aquarium water was renewed daily to maintain the sublethal Cd concentrations. Control fish were maintained under the same conditions in water devoid of detectable Cd. Fishes were fed before adding Cd to the water (every 24 h). We anesthetized

the fishes with cool water and euthanized them by ventral dissection 7 and 30 days post Cd exposure. We collected the liver, kidney, heart and gills were and stored them at -20 °C for biochemical test.

Mitochondrial viability with Janus Green B (JG-B) assay

We used a previously described protocol to measure mitochondrial fraction (Saz and Lescure, 1969). Briefly, we minced 0.05 g of tissue with small scissors in 1 ml of cold mitochondrial buffer (0.24 mg.l⁻¹ sucrose, 0.0005 mg.l⁻¹ EDTA and 0.15 % BSA pH 7.4). After mincing, we gently homogenized the tissue in a glass homogenizer and then centrifuged at 100 x g for 10 min at 4 °C to remove nuclei, unbroken cells, and other non-subcellular tissue. We then filtered the supernatant through glass wool and centrifuged it at 7800 x g for 30 min at 4 °C. We then re-suspended the dark packed lower layer (heavy mitochondrial fraction) in mitochondrial buffer and again centrifuged at 7800 x g for 30 min. We verified the mitochondrial integrity fraction by NADH extinction at 340 nm.

We used specialized mitochondrial staining, JG-B, for the mitochondrial viability assay (Mohammadi and Ghazi, 2007). We suspended the mitochondrial suspension (1 mg protein) with JG-B (1 mg.l⁻¹) in 1:1 proportion. We prepared the blank solution the same way, but without the addition of mitochondrial suspension. Each sample was then measured via spectrophotometry at 607 nm. Mitochondrial viability (% MV) was calculated using the formula:

$$\% \text{ mitochondrial toxicity} = 1 - \frac{\text{Mean absorbance of toxicant}}{\text{Mean absorbance of negative control}} \times 100\%$$

$$\% \text{ VM} = 100 \% - \% \text{ mitochondrial toxicity}$$

Acid soluble thiols (AST) determination

We determined the level of acid soluble thiols (AST) as previously described (Sedlak and Lindsay 1968), with the proper modifications. We homogenized a 0.1 g of tissue with 0.9 ml Tris EDTA buffer (30 mmol.l⁻¹ Tris HCl, 3 mmol.l⁻¹ EDTA, pH 8.9). We then centrifuged the homogenate at 1500 rpm for 5 min at 4 °C. We then mixed 200 µl of supernatant with 0.5 g sulfosalicylic acid, placed in the freezer for 15 min, and then centrifuged at 7000 rpm for 10 min to precipitate proteins. We measure the absorbance of the reaction mixture (200 µl supernatant, 800 µl Tris HCl buffer pH 8.9 and 80 µl 5.5'-dithiobis-2-nitrobenzoic acid - DTNB) at a wavelength of 412 nm. We performed the calibration curve from 100 µmol.l⁻¹ glutathione reduced (GSH) as standard and expressed the results in µmol⁻¹–SH/ml.

Statistical analysis

All values were expressed as mean±standard error (SEM). Statistical difference in % MV and AST levels (for each tissue evaluated during 7 and 30 days) were determined by Student's t-test (Ts). If the assumptions required for this test were not met, we used the nonparametric Mann-Whitney test. Additionally, we performed a Pearson Correlation analysis to associate % MV and AST levels in each *A. brevifilis* tissue tested. We analyzed all data using SPSS v15 (IBM Corporation, Armonk, New York, USA).

Results

During the last days of the Cd bioassay (7-30 days post exposure), we observed *A. brevifilis* displaying hyperactive aggressive behaviour in conjunction with damage on the skin and scales loss. This behavior was not observed in the control fish.

Mitochondrial viability and AST levels in *A. brevifilis* tissues

Liver

We detected a decrease in MV of hepatocytes by Cd exposure ($P \leq 0.05$), and the lowest values were observed in fish exposed for 30 d (Fig. 1a). We also observed a significant difference ($P \leq 0.01$) in AST distribution in the liver between exposed and unexposed fishes in both treatment groups. We detected lower AST concentrations in fishes in the 7d Cd exposed group

compared to the controls and higher concentrations in the 30d Cd exposed group when compared to the controls (Fig. 1b).

Kidney

In the kidney we observed a decrease in MV in the fishes exposed to Cd. The lowest values were observed in fishes exposed to Cd for 30 d (Fig. 2a). We also detected a significant decrease in AST concentrations in 7d Cd exposed fishes ($P \leq 0.001$); however we did not detect any significant differences in the fishes exposed to Cd for 30 d (Fig. 2b).

Heart

We did not detect any difference in heart MV as a result of Cd exposure (Fig. 3a); however AST levels in this tissue decreased significantly ($P \leq 0.001$) in Cd exposed fish both at in both exposure groups (Fig. 3b).

Gills

We detected an increase in MV in the gills of *A. brevifilis* in the fishes in the 30 d exposure group ($P \leq 0.01$) (Fig. 4a). We also detected a significant decrease of AST concentration in the 30 d exposure group. ($P \leq 0.001$) (Fig. 4b).

Correlation between MV and AST levels in *A. brevifilis* tissues

We detected an inverse association between MV and AST levels ($r = -0.968^{**}$) in the heart. However, in the gills we observed a positive association between %




Fig. 1. (a): Mitochondrial viability (%) of *A. brevifilis* liver 7 and 30 days of Cd exposure. (b): AST levels of *A. brevifilis* liver control, and 7 and 30 days Cd exposed. *: significant differences ($P < 0.05$); **: Very significant differences ($P < 0.01$); ***: Highly significant differences ($P < 0.001$).




Fig. 2. (a): Mitochondrial viability (%) of *A. brevifilis* kidney 7 and 30 days of Cd exposure. (b): AST levels of *A. brevifilis* kidney control, and 7 and 30 days Cd exposed. NS: Not significant; ***: Highly significant differences ($P < 0.001$).




Fig. 3. (a): Mitochondrial viability (%) of *A. brevifilis* heart 7 and 30 days of Cd exposure. (b): AST levels of *A. brevifilis* heart control, and 7 and 30 days Cd exposed. NS: Not significant; ***: Highly significant differences (P<0.001).




Fig. 4. (a): Mitochondrial viability (%) of *A. brevifilis* gills 7 and 30 days of Cd exposure. (b): AST levels of *A. brevifilis* gills control, and 7 and 30 days Cd exposed. NS: not significant; **: Very significant differences (P<0.01); ***: Highly significant differences (P<0.001).

Table 1. Correlation analysis between mitochondrial viability (%) and AST levels (µmol.l⁻¹-SH/ml) in *A. brevifilis* tissues exposed to Cd.

Tissue	Pearson correlation (r)	p value
Liver	-0.395 ^{NS}	0.439
Kidney	0.013 ^{NS}	0.981
Heart	-0.968 ^{**}	0.001
Gills	0.933 ^{**}	0.007

P: Probability; NS: Not significant; **: Very significant p<0.01.

MV and AST levels (r = 0.933^{**}). We did not detect any significant trends with MV and AST in the liver (Table 1).

Discussion

We showed that MV is able to be used as a marker for chronic Cd chronic toxicity in *A. brevifilis*. The liver and kidney are the most sensitive organs to mitochondrial damage generated by chronic metal toxicity, which correlates with their primary function as Cd bioaccumulator tissues (Perera *et al.*, 2015). Similar results in mitochondrial Cd toxicity have been reported in a variety of organisms from mammals (Nguyen *et al.*, 2015), *Crassostrea virginica* oyster (Sokolova, 2004), fish (Adiele *et al.*, 2010, Padmini and Usha, 2011) to plants (Miller *et al.*, 1973), suggesting that metal site

specific target are conserved throughout distant taxa.

The mitochondria are an essential organelle to ATP generation, involved in other processes such as ROS natural generation, cell death and the ageing processes (Chaiyakit and Thongboonkerd, 2009). Reports suggest that mitochondrial function is highly tuned and prone to heavy metals damage (Adiele *et al.*, 2010). The presence of metals such as Cd substantially increases ROS production resulting in lipid peroxidation, mtDNA cleavage and ATP synthesis inhibition which results in mitochondrial damage and apoptosis induction (Cuypers *et al.*, 2010). Cadmium may inhibit Wang *et al.* (2004) inhibit complex III CTE (ubiquinone: cytochrome c oxidoreductase) in the liver, heart, and brain mitochondria; in addition, to stimulating ROS production in this complex (Wang *et al.* 2004). This accumulation is probably mediated by Cd binding between semiubiquinone and cytochrome b₅₆₆ of the Q₀ site of cytochrome b complex III, resulting in accumulation of semiubiquinones at the Q₀ site. The semiubiquinones, being unstable, are prone to transfer one electron to molecular oxygen to form superoxide. In *Oncorhynchus mykiss*, Cd accumulation in mitochondria of liver was related to complex III inhibition, causing mitochondrial damage (Adiele *et al.*, 2010). It is possible that similar processes could be occurring in *A. brevifilis* liver mitochondria.

Gobe and Crane (2010) reported that kidney mitochondrial toxicity is mediated by similar mechanisms to those reported for liver, including ROS production, ATP altered levels and membrane potential alteration that stimulates apoptosis activation causing nephrotoxicity. Cell morphology studies in different organs of the fish *Dicentrarchus labrax*, showed that the mitochondria in liver, kidney, and gills exposed to Cd were damaged. This damaged was marked by signs of swelling, disappearance of ridges, vacuolation and myeloid bodies formation, classic characteristics of cellular processes caused by high metals action (Giari et al., 2007).

In the present study, cardiac mitochondrial viability of Guaraguara was unaffected by Cd exposure. Some studies suggest that heart mitochondria are vulnerable to metal poisoning (Wang et al., 2004); however, we found a negative association between MV and AST levels in heart. This result suggests that increment of AST concentration emerges as a compensatory mechanism conferring long-term protection to mitochondria against the Cd oxidative effect. Hatem et al. (2014) reported the exist of a protection mechanism against oxidative stress conditions that stimulates GSH accumulation in nucleus and mitochondria.

On the other hand, the variation of AST content in *A. brevifilis* could be an adaptive and antioxidative effect to organ-specific Cd accumulation. Zirong and Shijun (2007) showed that GSH content may increase or decrease in different tissues exposed to Cd, attributable to organ-specific responses also varies among species. In liver tissues from *Labeo rohita*, the GSH content decreased in Cd chronic assay clear that heavy metals cause of oxidative stress (Khalid et al., 2015).

Cao et al. (2012) reported the long-term GSH increase (20 days exposure) with increase of γ -glutamylcysteine synthetase (γ -GCS) enzyme activity. This is believed to constitute an adaptive compensatory mechanism in the liver to synthesize GSH *de novo* as a way to counteract metal oxidative effect, as observed for *A. brevifilis* liver and the other tissues evaluated. Other studies in *Oreochromis niloticus* report decrease in liver of GSH associated to Cd oxidative effect and its use as a cofactor in glutathione peroxidase and glutathione transferase enzymatic systems (Zirong and Shijun, 2007).

GSH is synthesized in the liver and transported in the blood to organs such as kidney and muscle (Atli and Canli, 2008). In the kidney tissues of *Anguilla anguilla*, a decrease in GSH levels was evidence to prevent ROS regeneration (Ahmad et al., 2006). In the kidney tissue of *Paralichthys olivaceus*, it was found that GSH levels decrease in fish exposed to Cd, but was associated with high metal accumulation in these cells, which causes complexes Cd-GSH formation that results on GSH reduction (Cao et al., 2010, 2012). While, in *Cyprinus carpio* an increase in GSH levels and antioxidant enzymes

in liver and kidney were observed due to their function as high metal accumulation and detoxification organs (Dugmonits et al., 2013; Paritha and Deepak, 2015).

We could establish that the AST elevated levels in some tissues act as cytoprotective and adaptive alternative mechanism related to the ROS detoxification, maintenance redox status and mitochondrial viability. Additionally, it is also important to note that the Cd exposure effect on AST levels, vary across fish tissues and is related to the exposure duration, the molecule dynamics in different tissues, the organism, and environmental conditions. Future research should include histopathology studies and an evaluation with other biochemical markers (e.g. antioxidant enzymes) to help establish tolerance mechanisms of *A. brevifilis* to heavy metal contamination.

Conflict of interest

The Authors declare that there is no conflict of interest.

Acknowledgments

This study was supported by funding from FONACIT (Project No. G2005000775) and Research Council-Universidad de Oriente. We are grateful to Postgraduate Applied Biology at the University of Oriente, Venezuela and Yaque community (providing the fish used in this study).

References

- Adiele, R., Stevens, D. and Kamunde, C. 2010. Reciprocal enhancement of uptake and toxicity of cadmium and calcium in rainbow trout (*Oncorhynchus mykiss*) liver mitochondria. *Aquat. Toxicol.* 96, 319-327.
- Ahmad, I., Maria, V., Oliveira, M., Pacheco, M. and Santos, M. 2006. Oxidative stress and genotoxic effects in gill and kidney of *Anguilla anguilla* exposed to chromium with or without pre-exposure to beta-naphthoflavone. *Mut. Res.* 608, 16-28.
- Akpakpan, E. and Akpanyung, E. 2014. Stress and histopathological studies in fish from Ibaka and Ifiayong Rivers, Akwa Ibom state, Nigeria. *World Appl. Sci. J.* 32(7), 1209-1218.
- Atli, G. and Canli, M. 2008. Response of antioxidant system of freshwater fish *Oreochromis niloticus* to acute and chronic metal (Cd, Cu, Cr, Zn, Fe) exposures. *Ecotox. Environ. Safe.* 73, 1884-1889.
- Bifano, C. and Mogollón, J. 1995. Metallic contaminant profile in sediment cores from lake Valencia, Venezuela. *Environ. Geochem. Health* 17(3), 113-118.
- Cao, L., Huang, W., Liu, J., Yin, X. and Dou, S. 2010. Accumulation and oxidative stress biomarkers in Japanese flounder larvae and juveniles under chronic cadmium exposure. *Comp. Biochem. Phy. Part C* 151, 386-392.
- Cao, L., Huang, W., Shan, X., Ye, Z. and Dou, S. 2012. Tissue-specific accumulation of cadmium and its effects on antioxidative responses in Japanese

- flounder juveniles. *Environ. Toxicol. Pharmacol.* 33, 16-25.
- Chaiyarit, S. and Thongboonkerd, V. 2009. Comparative analyses of cell disruption methods for mitochondrial isolation in high-throughput proteomics study. *Anal. Biochem.* 394, 249-258.
- Corona, J. 2013. Contaminación antropogénica en el Lago de Maracaibo, Venezuela. *Biocenosis* 27(1-2), 85-93.
- Cuyppers, A., Plusquin, M., Remans, T., Jozefczak, M., Keunen, E., Gielen, H., Opdenakker, K., Nair, A., Munters, E., Artois, T., Nawrot, T., Vangronsveld, J. and Smeets, K. 2010. Cadmium stress: an oxidative challenge. *Biometals* 23, 927-940.
- Dorts, J., Bauwin, A., Kestemont, P., Jolly, S., Sanchez, W. and Silvestre, F. 2012. Proteasome and antioxidant responses in *Cottus gobio* during a combined exposure to heat stress and cadmium. *Comp. Biochem. Phys. Part C* 155, 318-324.
- Dugmonits, K., Ferencz, A., Jancsó, Z., Juhász, R. and Hermes, E. 2013. Major distinctions in the antioxidant responses in liver and kidney of Cd²⁺-treated common carp (*Cyprinus carpio*). *Comp. Biochem. Phys. Part C* 158, 225-230.
- Fondo Nacional de Ciencia, Tecnología e Innovación (FONACIT). 2011. Código de ética para la vida. Ministerio del poder popular para la Ciencia, Tecnología e Industrias Intermedias.
- Giari, L., Manera, M., Simoni, E. and Dezfuli, B. 2007. Cellular alterations in different organs of European sea bass *Dicentrarchus labrax* (L.) exposed to cadmium. *Chemosphere* 67, 1171-1181.
- Gobe, G. and Crane, D. 2010. Mitochondria, reactive oxygen species and cadmium toxicity in the kidney. *Toxicol. Lett.* 198, 49-55.
- Hatem, E., Berthonaud, V., Dardalhon, M., Lagniel, G., Baudouin-Cornu, P., Huang, M., Labarre, J. and Chédin, S. 2014. Glutathione is essential to preserve nuclear function and cell survival under oxidative stress. *Free Radical Bio. Med.* 67, 103-114.
- Hermoso, D. and Márquez, M. 2005. Evaluación de las concentraciones de metales pesados en tejidos de peces del Río Catatumbo y sus afluentes. Trabajo de pregrado. Escuela de Ingeniería Química, Universidad Rafael Urdaneta, Venezuela.
- Hernández, M. 2005. Proteínas en diferentes tejidos (hígado, riñón y branquias) del pez dulceacuicícola *Colossoma macropomum* (Cuvier, 1818) expuesto a dosis subletales de cloruro de cobre y cadmio, Grade thesis, Departamento de Bioanálisis, Universidad de Oriente, Venezuela.
- Khalid, M., Qureshi, N., Mubarak, M. and Bukhari, S. 2015. Heavy metals (chromium, copper and cadmium) induced oxidative stress in *Labeo rohita* (Hamilton, 1822) during acute and chronic toxicity experiment. *Int. J. Biosci.* 6(11), 64-72.
- Lárez, C. 2011. Genotoxicidad en células sanguíneas de la guaraguara *Ancistrus brevifilis* (Eigenmann, 1920), bajo condiciones controladas y en condiciones naturales en dos localidades del Río Manzanares, estado Sucre, Venezuela, M.S thesis, Postgrado de Biología Aplicada, Universidad de Oriente, Cumaná, Venezuela.
- Marcano, V. and Troconis, A. 2001. Evaluación del contenido de mercurio en el pescado expandido en la ciudad de Mérida, Venezuela. *Rev. Bio. Lat. Am.* 8(2), 15-24.
- Márquez, A., Senior, W., Fermín, I., Martínez, G., Castañeda, J. and González, A. 2008. Cuantificación de las concentraciones de metales pesados en tejidos de peces y crustáceos de la Laguna de Unare, estado Anzoátegui, Venezuela. *Rev. Cient. (Venez.)* 18(1), 73-86.
- Miller, R., Bittell, J. and Koeppe, D. 1973. The effect of cadmium on electron and energy transfer reactions in corn mitochondria. *Phy. Plant.* 28, 166-171.
- Mohammadi, A. and Ghazi, M. 2007. Comparative measurement of cyanide and paraquat mitochondrial toxicity using two different mitochondrial toxicity assays. *Toxicol. Mech. Method.* 17, 87-91.
- Mora, A., Alfonso, J., Baquero, J., Handt, H. and Vásquez, Y. 2013. Elementos mayoritarios, minoritarios y traza en muestras de sedimentos del medio y bajo Río Orinoco, Venezuela. *Rev. Int. Cont. Amb.* 29(3), 165-178.
- Nguyen, K., Rippstein, P., Tayabali, A. and Willmore, W. 2015. Mitochondrial toxicity of cadmium telluride quantum dot nanoparticles in mammalian hepatocytes. *Toxicol. Sci.* 146(1), 31-42.
- Osman, A. and Kloas, W. 2010. Water quality and heavy metal monitoring in water, sediments, and tissues of the African Catfish *Clarias gariepinus* (Burchell, 1822) from the River Nile, Egypt. *J. Environ. Prot.* 1, 389-400.
- Padmini, E. and Usha, M. 2011. Mitochondrial membrane potential is a suitable candidate for assessing pollution toxicity in fish. *Sci. Total Environ.* 409, 3687-3700.
- Paritha, A. and Deepak, M. 2015. The effect of Cadmium on antioxidant enzymes in the liver of fresh water fish *Cyprinus carpio* (Linn). *Biolife* 3(1), 50-53.
- Peltier, W. and Weber, C. 1985. Methods for measuring the acute toxicity OE Effluent to fresh water and marine organism. *Environ. Mon. Trans, Lab. USA* 60, 484-638.
- Perera, P., Kodithuwakku, S., Sundarabharathy, T. and Edirisinghe, U. 2015. Bioaccumulation of Cadmium in Freshwater Fish: An Environmental Perspective. *Insight Ecol.* 4, 1-12.
- Ruiz, L., Salazar, S., Pérez, J. and Alfonsi, C. 2005. Diversidad íctica del sistema hidrográfico Río

- Manzanares, estado Sucre, Venezuela. Bol. Cent. Inv. Biol. 39(2), 91-107.
- Sabullah, M., Ahmad, S., Shukor, M., Gansau, A., Syed, M., Sulaiman, M. and Shamaan, N. 2015. Heavy metal biomarker: Fish behavior, cellular alteration, enzymatic reaction and proteomics approaches. Int. Food Res. J. 22(2), 435-454.
- Salazar, R. 2009. Estado de conocimiento de las concentraciones de cadmio, mercurio y plomo en organismos acuáticos de Venezuela. Rev. Elec. Vet. 10(11), 1-15.
- Salazar, R., Pérez, R., León, A., Lemus, M. and Rojas, L. 2009. Determinación de tioles totales y tioles solubles en ácido en el pez *Colossoma macropomum* (Cuvier, 1818) expuesto a cadmio. Rev. Cient. FCV-LUZ 19(4), 414-420.
- Salazar-Lugo, R., Vargas, A., Moreno, C., Centeno, L., Astudillo, H., Lemus, M. and Rojas de Astudillo, L. 2014. Parámetros sanguíneos y metales pesados en tejido del pez del río Orinoco, Venezuela. Rev. Cient. FCV-LUZ 24(3), 261-266.
- Saz, H. and Lescure, O. 1969. The functions of phosphoenol pyruvate carboxikinase and malic enzyme in the anaerobic formation of succinate by *Ascaris lumbricoides*. Comp. Biochem. Phy. 30, 49-60.
- Sedlak, J. and Lindsay, R. 1968. Estimation of total protein bound non-protein sulfidryl groups in tissue with Ellman's reagent. Anal. Biochem. 25, 192-205.
- Sevcikova, M., Modra, H., Slaninova, A. and Svobodova, S. 2011. Metals as a cause of oxidative stress in fish: a review. Vet. Med. 56(11), 537-546.
- Soares, S., Martins, H., Gutiérrez-Merino, C. and Aureliano, M. 2008. Vanadium and cadmium in vivo effects in teleost cardiac muscle: Metal accumulation and oxidative stress markers. Comp. Biochem. Phy. Part C 147, 168-178.
- Sokolova, I. 2004. Cadmium effects on mitochondrial function are enhanced by elevated temperatures in a marine poikilotherm, *Crassostrea virginica* Gmelin (Bivalvia: Ostreidae). J. Exp. Biol. 207, 2639-2648.
- Soud, G., Souayed, N., Yaktiti, F. and Maaroufi, K. 2013. Effect of acute cadmium exposure on metal accumulation and oxidative stress biomarkers of *Sparus aurata*. Ecotox. Environ. Safe. 89, 1-7.
- Vanegas, V. 2003. Determinación de los metales pesados: Plomo, Vanadio, Zinc y Cadmio en algunas especies de peces presentes en el Río Guasare, estado Zulia, Grade thesis, Facultad Experimental de Ciencias, Universidad del Zulia, Venezuela.
- Wang, Y., Fang, J., Leonard, S. and Krishna, M. 2004. Cadmium inhibits the electron transfer chain and induces reactive oxygen species. Free Radical Biol. Med. 36(11), 1434-1443.
- Weber, C. 1993. Methods for measuring the acute toxicity of effluents and receiving waters to *Pterygoplycthis multiradiatus* freshwater and marine organisms. Fourth ed. U.S. Environmental Protection Agency.
- Zirong, X. and Shijun, B. 2007. Effects of waterborne Cd exposure on glutathione metabolism in Nile tilapia (*Oreochromis niloticus*) liver. Ecotox. Environ. Safe. 67, 89-94.