

Research

Knowledge and awareness of medical doctors, medical students and nurses about dentistry in Nigeria

Elijah Olufemi Oyetola^{1,&}, Taiwo Oyewole¹, Micheal Adedigba¹, Stephen Tunde Aregbesola², Kehinde Umezudike³, Adedotun Adewale¹

¹Department of Preventive and Community Dentistry, Obafemi Awolowo University Ile-Ife, Nigeria, ²Department of Oral and Maxillofacial surgery and Oral Pathology, Obafemi Awolowo University, Ile-Ife, Nigeria, ³Department of Preventive Dentistry, Lagos University Teaching Hospital, Lagos, Nigeria

[&]Corresponding author: Elijah Olufemi Oyetola, Department of Preventive and Community Dentistry, Obafemi Awolowo University, Ile Ife, Nigeria

Key words: Dental awareness, medical students, medical doctors, nurses

Received: 07/08/2015 - Accepted: 31/03/2016 - Published: 07/04/2016

Abstract

Introduction: Various studies have reported poor awareness and knowledge of dentistry in the Nigerian population. There is, however, paucity of information assessing the knowledge and awareness of medical doctors/students and nurses about dentistry. The present study is aimed at determining the knowledge and awareness of medical doctors/students and nurses about dentistry. **Methods:** Self-administered questionnaires were randomly distributed among medical doctors/students, and nurses of Obafemi Awolowo Teaching Hospitals' Complex, Ile-Ife, Nigeria. Information collected using the questionnaire included participants' biodata, questions evaluating dental awareness, knowledge of systemic and oral health connections as well as referral practices. The data analysis was done with STATA version 11 software. **Results:** A total of 300 questionnaires were randomly distributed among doctors/students and nurses, 206 were returned (response rate of 69%). Of the returned questionnaires, 129(63%) were males and 77(37%) were females. There were 42 medical doctors, 49 nurses and 115 medical students. The mean age of the participants was 26.7 years (SD 5.2). Majority (99.5%) was aware of dental profession, but 92% had never referred patients for dental consultation. One third (31%) of medical doctors believed that Ludwig angina was a cardiac disease. A large proportion of the respondents (61%) see no need for routine dental visit while 27% would want to visit the dentist only when they had a dental complaint. **Conclusion:** Although a large percentage of the participants claimed to be aware of dentistry, our findings revealed low level of knowledge and attitude to Dentistry. Efforts should be made towards closing this knowledge gap to achieve efficient oral health.

Pan African Medical Journal. 2016; 23:172 doi:10.11604/pamj.2016.23.172.7696

This article is available online at: <http://www.panafrican-med-journal.com/content/article/23/172/full/>

© Elijah Olufemi Oyetola et al. The Pan African Medical Journal - ISSN 1937-8688. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Despite its role in systemic health, oral health care is an aspect that is often neglected [1, 2]. The awareness and knowledge of dentistry is still grossly inadequate among many patients and health care workers [3]. The implication of this is that the vast majority of patients who should benefit from dental services or be referred for their various dental ailments would be unable to assess the services due to the poor awareness and knowledge of dentistry by the attending health care providers [4, 5]. Studies have shown that medical doctors and nurses also have vital roles in oral health care of the population and when awareness is lacking, patients suffer [6, 7]. This becomes necessary due to the well reported relationship between oral health and systemic/general health [8-11]. The need for medical practitioners to participate in oral health promotion is highly essential especially in Nigeria where there are limited number of dentist who are unequally distributed [7]. Medical doctors are the primary care givers for the vast majority of health related complaints and they are also expected to play active roles in oral health promotion [12]. As noted earlier, several studies have shown the roles of medical doctors in oral health care delivery. Such roles include the screening for oral diseases, emergency cares, pain management, referral for the alleviation of pain symptoms in general medical practice, pediatrics, and accident and emergency (A and E) department [13]. As such, family physicians and pediatricians have much more important roles to play and when their knowledge of dentistry and oral health care practices are low or poor, inappropriate dental managements and advice or referrals of patients will be the likely outcome. Similar trends were found among medical students in the study by Adeghe et al in which their knowledge of oral health was described as suboptimal as only 2.8% of the 279 medical students had good knowledge of dental specialties [13]. Nurses also play vital roles in health care of patients along with medical doctors [14]. They represent one of the first group of health care providers that give primary care to patients before referral to specialist doctors [15]. Their knowledge of oral health should also be vast so as give appropriate advice when necessary. However, studies have shown that their knowledge about dentistry is quite poor and has been described as been grossly inadequate [15-17]. Some researchers had advocated the need to lay emphasis on dentistry topics during the course of their training [15, 17]. However, Muttineni et al [14] in their study of 147 final year nursing students in India concluded that nursing students had adequate basic knowledge of oral health but were still deficient

in correct brushing techniques. Similar findings were also reported by Asif et al [18] amongst nursing students. With the poor awareness of oral health among the medical doctors, nurses and their students in different part of the world, it became necessary to consider this aspect of health care especially in this part of the world where there are limited resources. The aim of the present study was to determine the level of awareness and knowledge of dentistry amongst medical doctors, nurses and medical students in Obafemi Awolowo University Complex Ile Ife Nigeria.

Methods

Study design

The study was a descriptive, cross sectional study

Study location

The study was conducted among the Staff of Obafemi Awolowo University Teaching Hospitals' Complex (OAUTHC) Ile-Ife. The hospital is located in South Western Nigeria and provides tertiary health care for a teeming population in Osun state as well as the neighboring states of Oyo, Ogun, Ondo, and Kwara. It is a teaching hospital involve in the training of both undergraduate and post graduate medical doctors and dentists. Information was obtained from the participants with self-administered questionnaires. Three hundred questionnaires were distributed and 206 questionnaires were returned correctly filled (response rate of 68.7%), the questionnaires were distributed among Medical doctors, nurses and medical students. The participants were recruited using simple random sampling of the medical doctors, nurses and medical students in the medical and surgical wards.

Questionnaire design

The questionnaire was designed to have four sections: section A collected information on biodata such as age, sex, marital status and category of the participant such as medical students, nurse or medical doctor; section B contained information on dental awareness and knowledge of dentistry. Some of the questions asked in this section include "Have you ever been to a dentist before?" "What is the specialty of dentistry that treats abnormally arranged teeth?" "What are the causes of dental caries?" Section C assessed

participants' knowledge of the relationship between oral health and systemic health. The questions in this aspect assessed participants' knowledge of Ludwig's angina, periodontal disease and systemic health, nutrition and oral health. Section D evaluated the referral practices of the participants. They were asked when to refer patients with Halitosis, toothache and dental abscess to the dentist.

Statistical analysis

The information was collected and analyzed using SPSS (Statistical Package for Social Sciences) version 11. Descriptive statistics was used to characterize socio-demographic variables such as age, sex, and marital status. For descriptive continuous variables, the mean, median, minimum value, maximum value and appropriate measures of variability were determined. For descriptive variables that are categorical, simple frequency and percentages was determined.

Results

The sociodemographic characteristics of the respondents

A total of 206 subjects participated in the study; 129(63%) males and 77(37%) females. The mean age and standard deviation were 26.7 years and 5.2 respectively with the majority (88.8%) belonging to 21-30 year age group category and 86% had single marital status. More than half (55.8%) of the participants were medical students (Table 1).

Dental awareness and attitude of respondents to dentistry

Majority (99.5%) of the respondents was aware of dentistry; the only exception is one medical student who claimed to have never heard about dentistry. Less than half (38%) had been to a dentist for treatment or check up most of whom were medical doctors. Busy work schedules in 91(44%) participants and absence of dental complaints in 97 (47%) participants were some of the reasons why they would not want to seek dental consultation (Table 2).

Respondents' knowledge about dentistry and dental specialties

Majority (71.4%) of the respondents agreed that Oral and maxillofacial surgery is a dental specialty that treat facial fractures,

one medical student representing 0.97% of the respondents' population attributed management of facial fractures to oral medicine specialty. Less than half of the participants (48.5) were aware that orthodontics treats abnormally arranged teeth. About one-third (32.5%) of the respondents did not know the specialty of dentistry that treats a hole on the tooth and only 14.6% was able to attribute a hole in the tooth to destorative dentistry. Only 23(11%) of the participants associated the management of unusual facial pain to Oral Medicine while the majority of the respondents did not know which of the specialties of dentistry could best manage unusual facial pain (Table 3). Similarly, about two third (61.1%) of the respondents named complex sugar as aetiological factor for dental caries while 9.8% did not know the cause of dental caries.

Respondents' knowledge about systemic and oral health

Only 44(21%) respondents knew Ludwig angina to be a fascial space infection, majority of who were medical doctors. Others either believed it to be a cardiac disease (39.8%) or did not know what it was (35.4%). The Knowledge about periodontal disease exacerbating systemic health; 133(64.5%) of the participant believed periodontitis could exacerbate infective endocarditis. Majority of the respondents believed that oral health could affect the nutrition of patients (Table 4).

Referral practices among respondents

Only about one-third of the repondents strongly agreed that delayed dental treatment could lead to life threatening conditions. A large number (74.4%) also belived that the management of orofacial cancer was best done by oral and maxillofacial Surgeon. Most (92.2%) of the participants had never refered patients for dental consultation. However, majority of them believed that patients with toothache, offensive mouth odour and dental abscess should be referred to dentists for treatment (Table 5).

Discussion

Efficient health care delivery is a product of cooperation between every member of the health team [19, 20]. Members of the health team include several groups of health workers, both medical and paramedicals. Smooth interaction is needed between medical and dental team to achieve good health delivery [21]. The present

study, designed to determine dental awareness, knowledge and attitude among medical practitioners and medical students becomes relevant as findings from this study is expected to throw more light into the importance of maintaining close inter-relationship between medical and dental team. Our study showed that 205 out of 206 participants were aware of dentistry as a profession, representing 95.5% of the study population. This findings is similar to the reports of Chandra et al [22] where 100% of participants of high socio economic status group claimed they are awareness of dental diseases. Similarly, Asif et al [18] also found that 89.9% of their study population knew that there were 32 teeth in the adult cavity. The increased awareness in this study may be due to the fact that the study was done among the educated/professionals. However, results differs in studies done among medical students. Sujatha et al [12] showed that only 25% of the medical students has good oral health awareness. This is not unexpected since the students are in training. Despite the increased oral health campaign in the community at large, it appears that there has not been much of a positive change in attitude and knowledge of people to Dentistry [4, 23]. This was also reflected in our findings that showed only 78(38%) participants had ever visited the dentist for dental checkup/treatment, similar to the report of Bashiru et al [24] who noted that 71.6% of the 360 studied undergraduate students in southern Nigeria never visited a dentist before. However, other studies by Chandral et al [22] reported that 100% of the study population had at least one dental visit in the last one year. Also Doshi et al [25] studied medical students and reported that 79.4% of the participants had visited a dentist for check up at one point or the other in their life. The fear of pain is one of the reasons why people would not want to visit a dentist [2, 22, 26]. However, the present study only reported 4% of participants would not want to visit the dentist for fear of pain. Some of the reasons they gave included busy work schedule (44%) and absence of dental complaints (47%). This is probably because this study was conducted among medical practitioners who are familiar with the hospital environment and probably would have used self medication to control pain [27]. The present study showed that 73.5% of participants brushed their teeth once a day while only 26% of them brushed twice daily. Dhanasekaran et al [28] similarly studied 538 people and noted that only 8.6% of those above the age of 30 years brushed their teeth twice a day while 40% brushed once daily. Some other studies [18, 21, 29] in the scientific literature had shown that most people brush once daily. This may be due to the poor attitude of people to their oral health, busy schedule and poor dental awareness. Unlike Medical/Nursing training in Nigeria where

medical/Nursing students had little or no exposure to dental training, training as a dentist involves exposure to basic medical education in the course of their training as Dentists. This must have contributed to why some dentists show basic medical knowledge since medical training was incorporated into their undergraduate curriculum [30]. In the present study we observed gross deficiency in the knowledge of dentistry by medical doctors, nurses and medical students. Only 61.1% of the studied population could associate bacterial to dental caries, 21.4% could attribute Ludwig angina to be a complication of space infection while only 2.4% believed that periodontitis could exacerbate Diabetes mellitus. We also noted that medical doctors had a better knowledge compared to others. However, the knowledge is still unacceptable and this showed a gross lack of basic dental knowledge among medical health workers as earlier reported by many researchers [13, 31-33]. This is largely due to the little or no exposure to dental education during the course of their training. Therefore, inclusion of dental education into the medical curriculum will go a long way in improving dental knowledge and skills among medical practitioners as suggested by many studies [18, 21, 31, 33]. Oral health is an integral part of systemic health. Adequate level of understanding and cooperation is therefore needed between medical and dental health teams to ensure qualitative health care delivery. In addition, acquisition of basic knowledge of dentistry by medical health workers will go a long way in improving health care especially in the developing world with limited human resources.

Conclusion

This study has shown that majority of health workers within the hospital are aware of the dental profession but their knowledge and attitude to dentistry is still poor. Their oral hygiene practices in which majority brushed one daily and large number of them never visiting the dental clinic for check up and never refer patients for dental consultation are points to elaborate the need for understanding between medical and dental profession. Our study also demonstrated that medical doctors were better informed about Dentistry when compared to Nurses and Medical students. Inclusion of dental education at both undergraduate and postgraduate medical curriculum appear to be a promising solution to this unnecessary knowledge gap between the medical and dental profession.

What is known about this topic

- Generally there is poor awareness and knowledge of dentistry among the populace especially in the developing part of the world;
- Health workers are also included but little is known about the actual level of awareness of the various categories of health workers such as medical doctors/students and nurses.

What this study adds

- The present study determined the level of awareness and knowledge of dentistry amongst medical doctors, nurses and medical students in Obafemi Awolowo University Complex Ile Ife Nigeria.

Competing interests

The authors declare no competing interest.

Authors' contributions

Oyetola EO was involved in the conceptualisation of the idea, drafted the article and contributed to study design. Oyewole T helped in the modification of the idea and in the acquisition of data. Adedigba MA contributed extensively in analysis, interpretation of data and revision of manuscript for critical intellectual content. Aregbesola SB also worked extensively in the aspects of revising the article for critical intellectual content and in the methodology design. Umezudike K did a thorough revision of the manuscript and also helped with data interpretations and presentations. Adedotun A contributed in data collection, analysis and manuscript revision. All authors have read and agreed to the final version of this manuscript and have equally contributed to its content and to the management of the case.

Acknowledgments

The authors would like to thank Mrs Eunice Oyetola for her useful comments on the manuscript.

Tables

Table 1: The sociodemographic characteristics of the respondents

Table 2: Dental awareness and attitude of respondents to dentistry

Table 3: Respondents' knowledge about dentistry and dental specialties

Table 4: Respondents knowledge about systemic and oral health

Table 5: Referral practices among respondents

References

1. Bradbury-Jones C, Innes N, Evans D, Ballantyne F, Taylor J. Dental neglect as a marker of broader neglect: a qualitative investigation of public health nurses' assessments of oral health in preschool children. *BMC Public Health*. 2013 Apr 19; 13: 370. [PubMed](#) | [Google Scholar](#)
2. McGrath C, Sham AS, Ho DK, Wong JH. The impact of dental neglect on oral health: a population based study in Hong Kong. *International dental journal*. 2007 Feb; 57(1): 3-8. [PubMed](#) | [Google Scholar](#)
3. Umezudike Ka OA, Ayanbadejo PO. Periodontal health knowledge of nonmedical professionals and their oral hygiene behavior in a teaching hospital in Nigeria. *Eur J Gen Dent*. 2015; 4: 48-54. [Google Scholar](#)
4. Sofola OO. Implications of low oral health awareness in Nigeria. *Niger Med J*. 2010; 51(3): 131-3. [PubMed](#) | [Google Scholar](#)
5. Jain N, Mitra D, Ashok K, Dundappa J, Soni S, Ahmed S. Oral hygiene-awareness and practice among patients attending OPD at Vyas Dental College and Hospital, Jodhpur. *Journal of Indian Society of Periodontology*. 2012; 16(4): 524-528. [PubMed](#) | [Google Scholar](#)
6. Naidu R, Boodoo D, Percival T, Newton J. Dental emergencies presenting to a university-based paediatric dentistry clinic in the West Indies. *Int J Paediatr Dent*. 2005 May; 15(3): 177-84. [PubMed](#) | [Google Scholar](#)

7. Sa'adu Z, Abdulraheem I. Oral health care practice and socio-demographic findings among the physicians in Ilorin. *Niger J Med*. 2003 Oct-Dec; 12(4): 211-6. **PubMed | Google Scholar**
8. Gulsen B, Idil K, Rumezsa K, Isil B, Serdar C, Canan B et al. Oral health and inflammation in patients with end stage renal failure. *Peritoneal Dialysis International*. 2009; 29(4): 472-479. **PubMed | Google Scholar**
9. Kumar J, Samelson R. Oral health care during pregnancy: recommendations for oral health professionals. *New York State Dental Journal*. 2009; 75(6): 29-33. **PubMed | Google Scholar**
10. Lankarani KB, Sivandzadeh GR, Hassanpour S. Oral manifestation in inflammatory bowel disease: a review. *World J Gastroenterol*. 2013 Dec 14; 19(46): 8571-9. **PubMed | Google Scholar**
11. Somma F, Castagnola R, Bollino D, Marigo L. Oral inflammatory process and general health. Part 1: the focal infection and the oral inflammatory lesion. *Eur Rev Med Pharmacol Sci*. 2010; 14(12): 1085-1095. **PubMed | Google Scholar**
12. Sujatha B, Yavagal PC, Gomez MSS. Assessment of oral health awareness among undergraduate medical Students in Davangere city: a cross-sectional survey. *Journal of Indian Association Of Public Health Dentistry*. 2015; 12(1): 43-47. **PubMed | Google Scholar**
13. Adeghe H, Ehigiator O, Azodo C, Ehizele A. Nigerian clinical level medical students' knowledge of dental specialty. *Ann Med Health Sci Res*. 2012 Jul; 2(2): 157-60. **PubMed | Google Scholar**
14. Muttineni N, Bolla SC, Naheeda S, Shaik RB, Gantha NS, Reddy SS. Oral health awareness among the final year undergraduate nursing students in Khammam district, Telangana. *Journal of Health Research and Reviews*. 2014; 1(3): 70-74. **PubMed | Google Scholar**
15. White R. Nurse assessment of oral health: a review of practice and education. *British journal of nursing (Mark Allen Publishing)*. 2000; 9(5): 260-266. **PubMed | Google Scholar**
16. Costello T, Coyne I. Nurses' knowledge of mouth care practices. *British journal of nursing (Mark Allen Publishing)*. 2008 Feb 28-Mar 12; 17(4): 264-8. **PubMed | Google Scholar**
17. Sangappa SB, Bhojraj N, Godhi B, Manjunathappa T. Oral health care awareness among nursing students in an Indian school - an experimental study. *J Contemp Med Edu*. 2013; 1(4): 266-271. **PubMed | Google Scholar**
18. Asif N, Asif SM, Babu DBG, Whagrey S. Assessing the Oral Health Awareness Among Final Year the Undergraduate Nursing Students Abha, Saudi Arabia. *World Journal of Dentistry*. 2014; 5(4): 213-217. **PubMed | Google Scholar**
19. Zhang S, Edward CL, Chu CH. Attitude and awareness of medical and dental students towards collaboration between medical and dental practice in Hong Kong. *BMC Oral Health*. 2015 May 2; 15:53. **PubMed | Google Scholar**
20. Rogers JP, Stewart PR, Stapleton JV, Hribar DL, Adams P, Gale AE. An interdisciplinary approach to the management of complex medical and dental conditions. *Australian dental journal*. 2000 Dec; 45(4): 270-6. **PubMed | Google Scholar**
21. Radha G, Shaik Hyder Ali K, Pushpanjali K. Knowledge and attitude and practice of oral health among nursing staff and nursing students of Bangalore city. *Journal of Indian Association of Public Health Dentistry*. 2008; 11: 17-21. **Google Scholar**
22. Chandra Shekar B, Reddy C, Manjunath B, Suma S. Dental health awareness, attitude, oral health-related habits, and behaviors in relation to socio-economic factors among the municipal employees of Mysore city. *Ann Trop Med Public Health*. 2011; 4(2): 99-106. **PubMed | Google Scholar**
23. Patil A, Chavan S, Banghele O, Patel K, Patel K. Awareness of oral health among medical practitioners in Sangamner City- a cross-sectional survey. *Int Journal of Clinical Dental Science*. 2010; 1(1): 26-29. **PubMed | Google Scholar**
24. Bashiru B, Anthony I. Oral self-care practices among university students in Port Harcourt, Rivers State. *Niger Med J*. 2014; 55(6): 486-489. **PubMed | Google Scholar**

25. Doshi D, Baldava P. A comparative evaluation of self-reported oral hygiene practices among medical and engineering university students with access to health promotive dental care. *Journal of Contemporary Dental Practice*. 2007; 8(1): 068-75. **PubMed | Google Scholar**
26. Nagina P, Bilal A, Amjad B, Aamir MB. Oro dental health: awareness and practices. *JUMDC*. 2011; 2(2). **Google Scholar**
27. Montgomery AJ, Bradley C, Rochfort A, Panagopoulou E. A review of self-medication in physicians and medical students. *Occupational medicine (Oxford, England)*. 2011 Oct; 61(7): 490-7. **PubMed | Google Scholar**
28. Dhanasekaran R, Nayar S. Self-perceived need for dental care. *J Pharm Bioallied Sci*. 2015; 7(Suppl 1): S98-S100. **PubMed | Google Scholar**
29. Azodo CC, Unamatokpa B. Gender difference in oral Health perception and practices among medical house officers. *Russian Open Medical Journal*. 2012; 1(2): 0208. **PubMed | Google Scholar**
30. Isiekwe G, Sofola O, Sanu O, Oredugba F. Undergraduate dental curriculum of a Nigerian dental school and the behavioural science and management skills required for private dental practice. *Journal of Dentistry and Oral Hygiene*. 2013; 5(5): 40-44. **PubMed | Google Scholar**
31. Nwhator S, Olojede C, Ijarogbe O, Agbaje M. Self-assessed dental health knowledge of Nigerian Doctors. *East African Medical Journal*. 2003; 90(5): 147-155. **PubMed | Google Scholar**
32. Kumari N, Sheela S, Sarada P. Knowledge and attitude on infant oral health among graduating medical students in Kerala. *J Indian Soc Pedod Prev Dent*. 2006 Dec; 24(4): 173-6. **PubMed | Google Scholar**
33. Usman S, Bhat SS, Sargod SS. Oral health knowledge and behavior of clinical medical, dental and paramedical students in Mangalore. *J Oral Health Comm Dent*. 2007; 1(3): 46-48. **PubMed | Google Scholar**

Table 1: The socio-demographic characteristics of the respondents				
	Medical Doctors n=42 n (%)	Nurses n=49 n (%)	Medical Student n=115 n (%)	Total n=206 n (%)
Sex				
Male (%)	33 (78.6)	4 (8.2)	92 (80)	129 (63)
Female (%)	9 (21.4)	45 (91.8)	23 (20)	77 (37)
Marital Status				
Married (%)	18 (42.9)	7 (14.3)	3 (2.6)	178 (86)
Single (%)	24 (57.1)	42 (85.7)	112 (97.4)	28 (14)
Age (mean=26.7,SD=5.2)				
<20 years	0 (0)	0 (0)	1 (0.9)	1 (0.5)
21-30 years	22 (52.4)	47 (96.0)	114 (99.1)	183(88.8)
31-40 years	15 (35.7)	0 (0)	0 (0)	15 (7.3)
41-50 years	5 (11.9)	1 (2.0)	0 (0)	6 (2.9)
51-60 years	0 (0)	1 (2.0)	0 (0)	1 (0.5)

Table 2: Dental awareness and attitude of respondents to dentistry				
	Medical Doctors n=42 (%)	Nurses n=49	Medical Students n=115	Total n=206
Are you aware of a profession called Dentistry?				
Yes (%)	42 (100)	49(100)	114 (99.1)	205(99.5)
No (%)	0 (0)	0 (0)	1 (0.9)	1 (0.5)
Have you ever been to a dentist for treatment or check-up?				
Yes (%)	29 (69)	12 (24.5)	38 (33)	78 (38)
No (%)	13 (31)	35 (75.5)	77 (67)	127 (62)
Why would you not want to visit a dentist?				
Fear of pain (%)	1 (2.4)	3 (6.1)	4 (3.5)	8 (4)
Fear of needle (%)	0 (0)	0 (0)	1 (0.9)	1 (0.5)
High Cost (%)	1 (2.4)	3 (6.1)	3 (2.6)	7 (3.4)
Lack of time (%)	28 (66.7)	12 (24.5)	51 (44.4)	91 (44)
No Dental Complaints (%)	12 (28.6)	31 (63.3)	54 (46.7)	97 (47)
How often do you brush your teeth?				
Once Daily	34 (81.0)	31 (63.3)	87 (75.7)	152 (73.5)
Twice Daily	7 (16.8)	18 (36.7)	28 (24.4)	53 (26)
≥Thrice	1 (2.4)	0 (0.0)	0 (0.0)	1 (0.5)
Appropriate time for routine dental consultations				
Once every 6 months	7 (6)	0 (0)	7 (17)	14 (7)
Once every year	8 (7)	1 (2)	2 (5)	11 (5)
When there is dental pain/other dental problem	25 (22)	11 (23)	20 (48)	56 (27)
There is no need	75(65)	37 (76)	13 (31)	126 (61)
Patient with dental abscess:				
Refer the patient to dental clinic	36 (85.7)	37(75.5)	102 (88.7)	169 (82)
Give antibiotics and analgesics	6 (14.3)	11 (22.5)	12 (10.4)	29 (14)
Ignore/Leave alone	0 (0.0)	1 (2.0)	1 (0.9)	2 (4)
Patient with toothache:				
Refer the patient to dental clinic	33 (78.6)	31 (63.3)	96 (83.5)	160 (78)
Give antibiotics and analgesics	9 (21.4)	17 (34.7)	18 (15.6)	44 (21)
Ignore/Leave alone	0 (0.0)	1 (2.0)	1 (0.9)	2 (1)
Do you think dentists also admit patients?				
Yes	37 (88.1)	43 (89.6)	85 (73.9)	165 (80)
No	1 (2.4)	3 (6.3)	8 (7.0)	12 (6)
Not Sure	4 (9.5)	3 (6.3)	22 (19.1)	29 (14)

Table 3: Respondents' knowledge about dentistry and dental specialties				
	Medical Doctors n=42	Nurses n=49	Medical Students n=115	Total N=206
Specialties of dentistry that treat facial fractures				
Orthodontics	2 (4.8)	14(28.6)	11 (9.5)	27 (13)
Prosthodontics	0 (0)	4 (8.2)	0 (0)	4 (1.9)
Oral and Maxillofacial Surgery	36 (85.7)	25(51.1)	86 (74.8)	147 (71.4)
Periodontology	2 (4.8)	2 (4.1)	0 (0)	4 (1.9)
Oral Medicine	0 (0)	2 (4.1)	0 (0.0)	2 (0.9)
I don't know	2 (4.8)	2 (4.1)	14 (12.2)	18 (8.7)
Orthodontics & Maxillofacial Surgery	0 (0.0)	0 (0.0)	4 (3.5)	4 (1.9)
Total	42 (100)	49 (100)	115 (100)	206 (100)
Specialties of dentistry that treat abnormally arranged teeth				
Orthodontics	26 (61.9)	23(46.9)	51 (44.4)	100 (48.5)
Endodontics	12 (28.6)	13(26.5)	50 (43.5)	75 (36.4)
Maxillofacial Surgery	1 (2.4)	2 (4.1)	4 (3.5)	7 (3.4)
Periodontology	3 (7.1)	10(20.4)	8 (6.7)	21 (10.2)
Oral Medicine	0 (0)	1 (2.0)	1 (0.9)	2 (0.97)
Total	42 (100)	49 (100)	115 (100)	206 (100)
Specialties of dentistry that treat hole on the tooth				
Orthodontics	6 (14.3)	5 (10.2)	13 (11.3)	24 (11.7)
Odontodontics	11 (26.2)	14(28.6)	29 (25.2)	54 (26.2)
Maxillofacial Surgery	0 (0)	3 (6.1)	2 (1.8)	5 (2.4)
Periodontologist	3 (7.1)	8 (16.3)	8 (6.7)	19 (9.2)
Oral Medicine	2 (4.8)	3 (6.1)	2 (1.7)	7 (3.4)
Restorative Dentistry	10 (23.8)	8 (16.3)	12 (10.4)	30 (14.6)
I don't know	10 (23.8)	8 (16.3)	49 (42.6)	67 (32.5)
Total	42 (100)	49 (100)	115 (100)	206 (100)
Specialties of dentistry that treat unusual facial pain				
Orthodontics	2 (4.8)	4 (8.2)	2 (1.7)	8 (4)
Odontodontics	7 (16.8)	5 (10.2)	10 (8.7)	22 (10)
Maxillofacial Surgery	7 (16.8)	22(44.9)	12 (10.4)	41 (20)
Plastodontics	4 (9.5)	4 (8.2)	12 (10.4)	20 (10)
Oral Medicine	6 (14.3)	5 (10.2)	12 (10.4)	23 (11)
I don't know	16 (38.1)	9 (18.4)	67 (58.3)	92 (45)
Total	42 (100)	49 (100)	115 (100)	206 (100)
Causes Dental caries				
Smoking	1 (2.4)	1 (2)	12 (10.4)	14 (6.7)
Complex Sugars	6 (14.3)	13(26.5)	20 (17.4)	39 (18.9)
Bacteria	24 (57.1)	29(59.2)	74 (64.3)	127(61.7)
Tooth picking	10 (23.8)	3 (6.1)	7 (6.1)	20 (9.7)
I don't know	1 (2.4)	3 (6.1)	2 (1.7)	6 (2.9)
Total	42 (100)	49 (100)	115 (100)	206 (100)

Table 4: Respondents knowledge about systemic and oral health				
	Medical Doctors n=42	Nurses n=49	Medical Students n=115	Total N=206
Ludwig's Angina is a:				
Cardiac Disease	13(31.0)	21 (42.9)	48 (41.7)	82 (39.8)
Renal Disease	0 (0)	1 (2.0)	1 (0.9)	2 (0.97)
Space Infection	21(50.0)	7 (14.3)	16 (13.9)	44 (21.4)
Lung Infection	0 (0)	4 (8.1)	1 (0.9)	5 (2.4)
I don't know	8 (19.1)	16 (32.7)	49 (42.6)	73 (35.4)
Periodontal disease could exacerbate:				
Diabetes	0 (0)	2 (4.1)	3 (2.6)	5 (2.4)
Heart Attack	1 (2.4)	0 (0)	1 (0.9)	2 (0.97)
Peptic Ulcer	1 (2.4)	7 (14.3)	0 (0.0)	8 (3.8)
Infective Endocarditis	30(71.4)	30 (61.2)	73 (63.5)	133 (64.5)
I don't know	5 (11.9)	10 (20.4)	35 (30.4)	50 (24.2)
Heart Attack & Infective Endocarditis	0 (0.0)	0 (0.0)	3 (2.6)	3 (1.4)
Diabetes & Infective Endocarditis	5 (11.9)	0 (0.0)	0 (0.0)	5 (2.4)
A life threatening dental condition:				
Caries	0 (0.0)	0 (0.0)	1 (0.9)	1 (0.5)
Ludwig's Angina	33(78.6)	24 (49.0)	64 (55.7)	121(58.7)
Periodontal Disease	3 (7.1)	14 (28.6)	6 (5.2)	23 (11.2)
I don't know	5 (11.9)	11 (22.4)	44 (38.3)	60 (29.1)
Ludwig's Angina & Periodontal Disease	1 (2.4)	0 (0.0)	0 (0.0)	1 (0.5)
Life threatening condition due to untreated dental disease:				
Brain Tumour	0 (0.0)	1 (2.0)	4 (3.5)	5 (2.4)
Hemangioma	1 (2.4)	8 (16.4)	0 (0.0)	9 (4.4)
Carvenous Thrombosis	34(80.9)	27 (55.1)	57 (49.6)	118 (57.3)
I don't know	7 (16.7)	13 (26.5)	52 (45.2)	72 (35)
Oral health can affect the nutrition of patients:				
Yes	42(100.0)	49(100.0)	102(88.7)	193 (93.7)
No	0 (0.0)	0 (0.0)	2 (1.7)	2 (0.97)
Not Sure	0 (0.0)	0 (0.0)	11 (9.6)	11 (5.3)

Table 5: Referral practices among respondents				
	Medical Doctors	Nurses	Medical Students	Total
Delayed referral of dental treatments can result in life-threatening condition				
Strongly agree	15(35.7)	15(30.6)	45 (39.1)	75 (36.4)
Agree	26(61.2)	30(61.2)	57 (49.6)	113 (54.8)
Strongly Disagree	1 (2.4)	2 (4.1)	9 (7.8)	12 (5.8)
Disagree	0 (0)	2 (4.1)	4 (3.5)	6 (2.9)
Total	42 (100)	49 (100)	115 (100)	206 (100)
Cancers in the orofacial region are best managed by				
General surgeons	3 (7.1)	10(20.4)	5 (4.4)	18 (8.7)
Maxillofacial surgeons	35(83.3)	34(69.4)	85 (73.9)	154 (74.8)
Plastic surgeons	40(95.2)	1 (2)	4 (3.5)	45 (21.8)
I don't know	0 (0)	0 (0)	16 (13.9)	16 (7.7)
Have you ever referred any patient to the dentist before				
Yes	5 (11.9)	1 (2)	11 (9.6)	17 (8.3)
No	37(88.1)	49 (98)	104(90.4)	190 (92.2)
Patients with toothache				
Refer to a dentist	33(78.6)	31(62.3)	96 (83.5)	160 (77.7)
Give antibiotic/analgesic	9 (21.4)	17(34.7)	18 (15.6)	44 (21.4)
Ignore	0 (0)	1 (2)	1 (0.9)	2 (0.9)
Patients with dental abscess				
Refer to dentist	36(85.7)	37(75.5)	102 (49.5)	175 (84.9)
Give antibiotic/analgesic	9 (21.4)	11(22.4)	12 (10.4)	32 (27.8)
Ignore	0 (0)	1 (2.0)	1 (0.9)	2 (0.9)
Patients with offensive mouth odour				
Refer to a dentist	25(59.5)	30(61.2)	99 (86.1)	154 (74.7)
Prescribe mouthwash	10(23.8)	13(26.5)	15 (13.4)	28 (24.3)
Give antibiotic/analgesic	7 (16.7)	6 (12.2)	1 (0.9)	14 (6.7)