

Images in medicine

Lingual botryomycoma in the aftermath of Stevens-Johnson syndrome

Houyam Moundib^{1,&}, Fouzia Hali¹

¹Department of Dermatology and Venereology, University Hospital Ibn Rochd, Casablanca, Morocco

[&]Corresponding author: Houyam Moundib, Department of Dermatology and Venereology, University Hospital Ibn Rochd, Casablanca, Morocco

Key words: Granulation tissue, botryomycoma, tongue

Received: 10/05/2015 - Accepted: 17/05/2015 - Published: 18/05/2015

Pan African Medical Journal. 2015; 21:35 doi:10.11604/pamj.2015.21.35.7036

This article is available online at: <http://www.panafrican-med-journal.com/content/article/21/35/full/>

© Houyam Moundib et al. The Pan African Medical Journal - ISSN 1937-8688. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Image in medicine

A 55 years old woman, was hospitalized in November 2011 for a Stevens-Johnson syndrome with severe mucosal impairment appeared three weeks after taking allopurinol (Zyloric®) for an hyperuricemia associated with arthralgia. She was put under symptomatic treatment after discontinuation of the offending molecule with a good mucocutaneous improvement. After healing of lesions of the oral mucosa in December 2011, the examination of the oral cavity has found a nodular median lesion of the dorsal surface of the tongue, in favor of a granulation tissue at the histology with spontaneous regression of the rest of lesion. The current decline is two years and two months without local recurrence.



Figure 1: a nodular median lesion of the dorsal surface of the tongue