

Research

Non alcoholic fatty liver disease in a Nigerian population with type II diabetes mellitus

Titilola Osawaye Olusanya¹, Olufunmilayo Adenike Lesi^{1,&}, Adekunle Ayokunle Adeyomoye², Olufemi Adetola Fasanmade³

¹Gastroenterology and Hepatology Unit, Lagos Nigeria, ²Endocrine and Metabolism Unit, Lagos Nigeria, Department of Medicine, Lagos Nigeria, ³Department of Radiology, Lagos, Nigeria, Faculty of Clinical Sciences, College of Medicine, University of Lagos Teaching Hospital, PMB 12003, Idi-Araba, Lagos Nigeria

[&]Corresponding author: Olufunmilayo Adenike Lesi, Department of Medicine, College of Medicine, University of Lagos Teaching Hospital, PMB 12003, Idi-Araba, Lagos, Nigeria

Key words: Fatty liver disease, ultrasound scan, dyslipidaemia, Africa

Received: 12/10/2015 - Accepted: 01/12/2015 - Published: 06/05/2016

Abstract

Introduction: Worldwide, Non-alcoholic fatty liver disease (NAFLD) has become an important cause of chronic liver disease and cardiovascular morbidity, even more so in subjects with Type II Diabetes Mellitus (T2DM). The aim of this study was to determine the prevalence and risk factors of NAFLD in an African population with Type II Diabetes Mellitus. **Methods:** We performed a case control study and evaluated anthropometric and biochemical risk factors for NAFLD in 336 subjects (T2DM and non-diabetic controls). Parameters assessed included estimation of BMI (Body Mass Index), measurement of waist circumference (WC), serum cholesterol including HDL-C, LDL-C and triglyceride and serum transaminases (ALT and AST). Hepatitis B and C viral antibody screening was also performed. The diagnosis of NAFLD was confirmed by identification of hepatic steatosis on abdominal ultrasound scan evaluation and exclusion of significant alcohol consumption. **Results:** NAFLD was identified in 16.7% (28 of 168) patients with T2DM compared with 1.2% (2 of 168) non-diabetic controls (Odds Ratio 16.6; $p < 0.001$). Central obesity ($WC > 102\text{cm}$) and dyslipidaemia ($HDL-c < 40\text{mg/dl}$) were independently associated with NAFLD in male subjects with T2DM ($p=0.03$ and $p=0.04$ respectively). **Conclusion:** NAFLD occurred more frequently in patients with T2DM than controls and was associated with central obesity and dyslipidaemia. The diabetic subjects with NAFLD will require more intensive therapy to decrease the risk of hepatic, cardiovascular and other adverse events.

Pan African Medical Journal. 2016; 24:20 doi:10.11604/pamj.2016.24.20.8181

This article is available online at: <http://www.panafrican-med-journal.com/content/article/24/20/full/>

© Titilola Osawaye Olusanya et al. The Pan African Medical Journal - ISSN 1937-8688. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Non-alcoholic fatty liver disease (NAFLD) is a spectrum of clinical and pathological conditions characterized by excessive lipid deposition in the liver parenchyma in the absence of significant alcohol consumption [1]. In the past decade, NAFLD has gained worldwide prominence as an important public health condition because of the enhanced risk of progressive end-stage liver disease, liver failure and hepatocellular carcinoma [2,3]. Current evidence suggests that NAFLD is the hepatic component of the metabolic syndrome, a syndrome that also includes central obesity, dyslipidemia, hypertension and insulin resistance/impaired glucose tolerance [3-5]. Recently published cross-sectional and prospective cohort studies suggest that 19% to 30% of asymptomatic adults in the USA and Europe have NAFLD [6,7]. The prevalence of NAFLD among people with Type II Diabetes Mellitus (T2DM) is significantly higher than reported in the general population and ranges from 21%-78% [4,8-10]. In T2DM, NAFLD pursues a more aggressive clinical course with necro-inflammation and fibrosis and progression to end-stage chronic liver disease. Beyond these hepatic consequences, NAFLD is being increasingly associated with an excess risk of cardiovascular disease, coronary artery disease, as well as peripheral vascular disease [11-13]. In African populations, there is a dearth of evidence on the prevalence and characteristics of NAFLD. Kruger et al [14], histologically confirmed NAFLD in 45% of an obese population in Western Cape, South Africa. They convincingly demonstrated a strong association between obesity, NAFLD and insulin resistance. This study was the first to describe the clinical characteristics of NAFLD in South Africa, using a study population comprised mostly of Caucasian subjects with only a minority of Africans. There however remains a dearth of epidemiological and clinical evidence on NAFLD in black Africans. The aim of this study was to determine the prevalence and risk factors for NAFLD in black Africans with T2DM. The findings from this study will provide an opportunity to enhance awareness of this condition and contribute to the evaluation and clinical management of the estimated 5 million people with T2DM in Nigeria [15].

Methods

The study was conducted at a University Teaching hospital in Lagos, South-West, Nigeria. The hospital is a 760-bed federal tertiary referral centre providing both medical and surgical care to the

general population of Lagos. Consecutively presenting adults >18 years old with diabetes were evaluated at the endocrine clinic between November 2011 and January 2012. Included were patients with established T2DM diagnosed according to the American Diabetes Association (ADA) diagnostic criteria [16] (ie symptoms of **diabetes plus random plasma glucose concentration \geq 200 mg/dl or fasting Plasma Glucose (FPG) \geq 126 mg/dl**). Controls were age and sex matched, non-diabetic patients with no history suggestive of diabetes eg polyuria, polydipsia, unintentional weight loss and fasting blood sugar level less than 126 mg/dl. Exclusion criteria included alcohol intake in excess of 20 grams per day, history of jejunoileal bypass surgery or extensive small bowel resection, usage of drugs known to cause secondary steatosis like corticosteroids, methotrexate, and amiodarone. Pregnant women and subjects with severe co-morbidities such as malignancies, congestive heart failure and chronic kidney disease were excluded. A structured questionnaire was administered to each subject to obtain information such as bio-data, alcohol intake, current medications as well as family history of diabetes and liver disease. All patients were clinically evaluated and weight (kg) and height (m) were measured. The Body Mass Index (BMI=weight (kg) /height (m²) was calculated and classified based on World Health Organisation (WHO) criteria [17]. Waist circumference was measured from the right side in the mid-axillary line, midway between the lower margin of the least palpable rib and the top of the iliac crest (highest point of the hip bone on the right side) as the point of reference. Hip circumference measurement (HC) in centimetres was done at the point of widest circumference of the buttocks with the tape parallel to the floor. Both waist and hip circumference was taken twice and the average was calculated [18]. Central obesity was defined as waist circumference (WC) more than 88 cm in females and more than 102 cm in males. The average of three BP readings was calculated. Hypertension was considered at a blood pressure reading of more than or equal to 140/90 mmHg or current intake of anti hypertensive or both [19].

Blood samples were taken for evaluation of serum alanine transaminase (ALT), aspartate transaminase (AST), fasting lipid profile including total cholesterol (TCHOL), high-density lipoprotein cholesterol (HDL-C), triglyceride (TG) (Randox Laboratory Limited, UK). Low-density lipoprotein cholesterol (LDL-C) was calculated using Friedwald formulae $LDL = (TCHOL - HDL - C) - TG/5$ [20]. Dyslipidemia was defined as having one or more of the criteria: **LDL-C \geq 100 mg/dL, total cholesterol \geq 200 mg/dL, triglycerides \geq 150 mg/dL, or HDL-C $<$ 40 mg/dL in males and $<$ 50 mg/dL in**

females [19]. Samples were also analysed for detection of anti-hepatitis C virus and hepatitis B surface antigen (Diaspot, USA). Abdominal Ultrasound scan (USS) evaluation was performed (Aloka Pro-sound 3500, Japan) using a 3.75 MHz probe. Findings suggestive of hepatic steatosis included "bright liver" with increased echogenicity, hepato-renal contrast and attenuation of the diaphragm [20,21]. The diagnosis of NAFLD was based on the absence of alcohol consumption (or alcohol intake less than 20 grams per day) and the presence of hepatic steatosis on USS evaluation. Ethical approval was obtained from the Ethics and Research committee of Lagos University Teaching Hospital Idi-Araba, Lagos, Nigeria prior to commencement of the study. Pre study counselling was done and written informed consent was obtained from all patients prior to enrolment. Minimal sample size was determined using the Cochrane statistical formula [22]. Total study population was estimated to be 320 subjects. Data analysis was performed using SPSS Version 20 (SPSS Inc, Chicago, IL USA). Cases and controls were compared using Chi-square or Fisher's exact test to assess significance between proportions and Students t-test to assess significance with quantitative data. Variables associated with NAFLD, with a significant P-value < 0.05 in univariate analysis, were included in a multivariable logistic regression. Relationships were assessed by Odds ratio (OR) and 95% confidence interval (CI).

Results

Characteristics of study subjects

A total of 340 subjects were recruited of which 336 subjects (168 subjects with T2DM and 168 control subjects without T2DM) were included in data analysis. 4 subjects were excluded due to incomplete laboratory evaluation. The mean age (standard deviation) of the patients with T2DM was 53.2 (8.6) years. There were 109 (64.9%) female and 59 (35.1%) males, with a Male to Female ratio of 1: 1.8. Most of the T2DM subjects (60%, n=101) had only basic education and 88% (n=148) earned an income of less than three hundred US dollars (\$300)/month). There was no significant difference between the cases and control subjects in terms of age, gender, educational background and monthly income. The T2DM subjects were however significantly more likely to have concurrent hypertension (n=80, 47.6%) and a family history of diabetes (n=74, 44%) compared to the non-diabetic controls (Odds

Ratio 3.2, p<0.001 and Odds ratio 6.5, p<0.001 respectively). The socio demographic variables of the study population are shown in Table 1.

Anthropometric, biochemical and ultrasound characteristics of study participants

As shown in Table 2, subjects with T2DM were significantly more likely to be overweight and have higher waist/hip ratio than the controls (p=0.02 and p<0.001 respectively). The mean systolic blood pressure was significantly higher among the cases compared with the controls (p<0.001). T2DM subjects had a higher prevalence of dyslipidaemia with mean total serum cholesterol and LDL-cholesterol significantly higher compared to the control subjects (p<0.001 and p<0.001 respectively). Although the mean AST and ALT values appeared higher in diabetics than controls, this association failed to reach statistical significance (p=0.05 and p=0.17 respectively). The distribution of clinical and laboratory characteristics of the study participants is shown in Table 2. Antibodies to Hepatitis C virus (anti HCV) were negative in all the study participants. Seven (7) subjects however were positive for Hepatitis B virus. Hepatic steatosis was detected on ultrasound scan evaluation in 28 of 168 (16.7%) T2DM subjects compared to 2 of 168 (1.2%) control subjects. (Odds Ratio 16.6, p<0.001). These subjects did not have a history of alcohol use (or consumed less than 20 grams of alcohol per day) and thus fulfilled the criteria for NAFLD.

Characteristics of NAFLD among subjects with T2DM

The characteristics of NAFLD in the diabetic subjects are shown in Table 3. NAFLD occurred more frequently in men (n=12/59, 20.3%) than in women (n=16/109, 14.7%). Subjects with NAFLD also appeared to have higher levels of ALT, AST and fasting serum triglyceride than subjects without NAFLD. These differences however did not attain statistical significance (p=0.53, p=0.4 and p=0.06 respectively). A high frequency of obesity (BMI ≥ 30 kg/m²) and central obesity (WC >88 cm) was noted in women with T2DM regardless of NAFLD status (p=0.19 and p=0.35 respectively). In contrast, diabetic men with NAFLD had a significantly higher prevalence of generalized obesity (BMI ≥ 30 kg/m²) than those without NAFLD (58.3% vs. 25.5%; Odds Ratio 4.0; 95% CI; 1.1-15.3). These men also had an 8-fold risk of central obesity with WC >102 cm (9 of 12 subjects, 75%) and a higher prevalence of dyslipidaemia (HDL-c <40 mg/dl in 4 of 12 subjects, 33.3%) than

T2DM men without NAFLD ($p=0.002$ and $p=0.013$ respectively) as shown in Table 3.

Discussion

In this study, we have shown that up to 16.7% (one in six) of the subjects with T2DM had NAFLD, a 16-fold risk of NAFLD compared to non-diabetic controls. Our finding of NAFLD in both the subjects with T2DM and non-diabetic controls corroborates a prior survey from West Africa that reports a prevalence of NAFLD in 9.5% of diabetics and 4.5% of non-diabetic controls [23]. There is however a paucity of evidence in this region. A significantly higher prevalence of NAFLD ranging from 40% to 60% of subjects with T2DM has been reported in Caucasian [11], Indian [24,25] and Asian populations [26]. In the USA, the prevalence of NAFLD in the general population varies from 15% in Caucasians to 28% in Hispanics [6]. The comparatively lower prevalence of NAFLD in our study may be a reflection of the different socio-demographic and economic characteristic of this African population. The association of obesity and increasing BMI is an established risk factor for NAFLD worldwide. In this current study, the mean body mass index (BMI) and income characteristics suggest that this population is not significantly obese and appear less affluent than western populations. This is not surprising as subjects were recruited from a large urban public hospital in a resource-limited setting. In addition, diagnostic tools used for evaluation of hepatic steatosis have differing accuracy in estimating hepatic fat. The computerized tomography (CT) scan and magnetic resonance imaging (MRI), although more sensitive in detecting minimal levels of hepatic steatosis, are expensive and impractical as screening tools [20,21]. The ultrasound scan (USS) has greater than 90% sensitivity and specificity in diagnosing hepatic steatosis and was used in this study. It remains the screening modality of choice because of low cost, safety, and accessibility [21]. Another important reason for the differing distribution of NAFLD appears to be related to the racial and ethnic characteristics of the population. In the US Multi-ethnic Study of Atherosclerosis (MESA), the prevalence of NAFLD in African Americans was found to be significantly lower than in Caucasian and Asian populations [6]. Various other studies corroborate this ethnic disparity [27-29]. Recent evidence from prospective multi-ethnic cohort studies have identified a strong association between NAFLD and genetic mutation in several genes including lipolytic enzymes, patatin like phospholipase (PNPLA-3) domain, neurocan gene NCAN

and glucokinase regulatory protein gene GCKR in non-Hispanic whites and subjects of European descents respectively [28,29]. These findings suggest a racial susceptibility to NAFLD and may have implications for African and other populations and require further research.

Risk factors for NAFLD

The association of obesity, concurrent hypertension and dyslipidaemia is not surprising and has been well characterized in diabetic populations worldwide. The key pathogenesis of this association is related to insulin resistance and the metabolic syndrome [3,4]. Our study shows that central obesity and dyslipidaemia although significantly associated with T2DM, are also independent risk factors for NAFLD, more so in men than women. Diabetes and NAFLD have mutual pathogenetic mechanisms, which may interact with metabolic derangements to accelerate NAFLD progression in diabetic patients [10,11,25]. The male susceptibility to NAFLD as shown in this study has also been described in Caucasian and Hispanic populations and has been attributed to lifestyle, sex hormones, differing distribution of adiposity, and other behavioral and genetic factors that remain to be fully defined. Although our study identifies important characteristics and associations of NAFLD in this African population, there are a few limitations. The reliance on patient's estimate of their alcohol consumption is subject to recall and may be inaccurate. Additionally, the gold standard for diagnosis of fatty liver is the liver biopsy. This procedure is invasive and was neither feasible nor ethical in this diabetic cohort.

Conclusion

This study has shown that NAFLD is common in subjects with T2DM and is associated with a 16-fold risk compared to non-diabetic controls. Although obesity, hypertension and dyslipidaemia were also significantly associated with T2DM, our study suggests that central obesity and dyslipidaemia are also independent risk factors for NAFLD, more so in men than women. Our result supports a role for routine ultrasound scan evaluation in T2DM to enhance clinical decision making and enable early recognition and management of NAFLD to limit its associated adverse hepatic and cardiovascular complications.

What is known about this topic

- The association of obesity, concurrent hypertension and dyslipidaemia has been well characterized in diabetic populations worldwide;
- In Western countries, patients with diabetes are a high risk group for NAFLD;
- They have a more aggressive clinical course with necro-inflammation and progression to liver disease and enhanced cardiovascular risk.

What this study adds

- In Nigerians, T2DM is associated with a 16- fold risk of NAFLD compared to non-diabetic controls;
- Nigerian men with T2DM appear more susceptible to NAFLD than women and have a significant association of NAFLD with truncal obesity and dyslipidaemia;
- Prevalence of NAFLD in this population is lower than in Caucasians, American blacks, and Hispanics.

Authors' contributions

TOO and OAL contributed to study design, implementation, data analysis, first and final manuscript. AAA and OAF contributed to study design and review of first and final manuscript. TOO also conducted patient evaluation.

Acknowledgments

Study subjects with T2DM and non-diabetic controls who participated in the study. Staff of radiology that assisted with ultrasound scans evaluation of study subjects.

Tables

Table 1: Socio demographic variables of T2DM subjects and controls

Table 2: Comparison of anthropometric, laboratory and ultrasound scan parameters between T2DM patients and control subjects

Table 3: Univariate analysis of the clinical and biochemical factors associated with NAFLD among subjects with T2DM

References

1. Ali R, Cusi K. New diagnostic and treatment approaches in non-alcoholic fatty liver disease. *Ann Med.* 2009; 41(4): 265-78. **PubMed | Google Scholar**
2. Erickson SK. Non Alcoholic Liver Disease. *J Lipid Res.* 2009; 50 Suppl: S412-S416. **PubMed | Google Scholar**
3. Adams LA, Angulo P, Lindor KD. Nonalcoholic fatty liver disease. *CMAJ.* 2005; 172(7):899-905. **PubMed | Google Scholar**
4. Cusi K. Nonalcoholic fatty liver disease in type 2 Diabetes Mellitus. *Curr Opin Endocrinol Diabetes Obes.* 2009; 16(2): 141-149. **PubMed | Google Scholar**
5. McCullough AJ. Epidemiology of the metabolic syndrome in the USA. *J Dig Dis.* 2011;12 (5):333-40. doi: 10.1111/j.1751-2980.2010.00469.x. **PubMed | Google Scholar**
6. Foster T, Anania FA, Li D, Katz R, Budoff M. The prevalence and clinical correlates of non-alcoholic fatty liver disease in African Americans: The multiethnic study of atherosclerosis (MESA). *Dig Dis Sci.* 2013; 58(8):2392-8. **PubMed | Google Scholar**
7. Lazo M, Hernaez R, Eberhardt MS, Bonekamp S, Kamel I, Guallar E, Koteish A, Brancati FI, Clark JM. Prevalence of Nonalcoholic Fatty Liver Disease in the United States: The Third National Health and Nutrition Examination Survey, 1988-1994. *Am J Epidemiol.* 2013; 178(1): 38-45. **PubMed | Google Scholar**
8. Bellentani S, Scaglioni F, Marino M, Bedogni G. Epidemiology of non-alcoholic fatty liver disease. *Dig Dis.* 2010; 28(1): 155-61. **PubMed | Google Scholar**
9. Angulo P. Nonalcoholic fatty liver disease. *N Engl J Med.* 2002; 346:1221-1231. **PubMed | Google Scholar**

10. Clark J, Diehl A. Hepatic steatosis and type 2 diabetes mellitus. *Curr Diab Rep.* 2002; 3:210-215. **PubMed | Google Scholar**
11. Targher G, Bertolini L, Padovani R, Rodella S, Zoppini G, Pichiri I, Sorgato C, Zenari L, Bonora E. Prevalence of non-alcoholic fatty liver disease and its association with cardiovascular disease in patients with type 1 diabetes. *J Hepatol.* 2010;53(4):713-8. doi: 10.1016/j.jhep.2010.04.030. **PubMed | Google Scholar**
12. Targher G, Bertolini L, Chonchol M, Rodella S, Zoppini G, Lippi G, Zenari L, Bonora E. Non-alcoholic fatty liver disease is independently associated with an increased prevalence of chronic kidney disease and retinopathy in type 1 diabetic patients. *Diabetologia.* 2010;53(7):1341-8. doi: 10.1007/s00125-010-1720-1. **PubMed | Google Scholar**
13. Hamaguchi M, Kojima T, Takeda N, Nagata C, Takeda J, Sarui H, Kawahito Y, Yoshida N, Suetsugu A, Kato T, Okuda J, Ida K, Yoshikawa T. Nonalcoholic fatty liver disease is a novel predictor of cardiovascular disease. *World J Gastroenterol.* 2007;13(10): 1579-1584. **PubMed | Google Scholar**
14. Kruger FC, Daniels C, Kidd M, Swart G, Brundyn K, Van Rensburg C, Kotze MJ. Non-alcoholic fatty liver disease in the Western Cape: a descriptive analysis. *S Afr Med J.* 2010;100(3):168-171. **PubMed | Google Scholar**
15. International Diabetes Federation. *IDF DIABETES ATLAS- 6th edition.* 2013. Brussels, Belgium. Accessed on March 16, 2015. **Google Scholar**
16. American Diabetes Association. *Diagnosis and Classification of Diabetes Mellitus.* *Diabetes care.* 2004; 27: Suppl 1S5-S10. **PubMed | Google Scholar**
17. World Health Organization. *Waist Circumference and Waist-Hip Ratio: Report of a WHO Expert Consultation.* Geneva WHO Press. 2011; 6th edition. **Google Scholar**
18. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, Jones DW, Materson BJ, Oparil S, Wright JT Jr, Roccella EJ. The seventh report of the joint national committee on prevention, detection, evaluation and treatment of high blood pressure. *JAMA.* 2003; 289:2560. **PubMed | Google Scholar**
19. Grundy SM, Cleeman JI, Merz CN, Brewer HB Jr, Clark LT, Hunninghake DB, Pasternak RC, Smith SC Jr, Stone NJ. National Heart, Lung, and Blood Institute, American College of Cardiology Foundation, American Heart Association: implications of recent clinical trials for the National Cholesterol Education Program Adult Treatment Panel III guidelines. *Circulation.* 2004;110(2):227-39. **PubMed | Google Scholar**
20. Hernaez R, Lazo M, Bonekamp S, Kamel I, Brancati FL, Guallar E, Clark JM. Diagnosis accuracy and reliability of ultrasonography for the detection of fatty liver: a meta-analysis. *Hepatology.* 2011;54(3):1082-1090. **PubMed | Google Scholar**
21. Schwenzer NF, Springer F, Schraml C, Stefan N, Machann J, Schick F. Non-invasive assessment and quantification of liver steatosis by ultrasound, computed tomography and magnetic resonance. *J Hepatol.* 2009;51(3):433-445. **PubMed | Google Scholar**
22. Fleiss JL. *Statistical Methods for Rates and Proportions.* New York: Wiley. 1981; 2nd ed. **Google Scholar**
23. Onyekwere CA, Ogbera AO, Balogun BO. Non-alcoholic fatty liver disease and the metabolic syndrome in an urban hospital serving an African community. *Ann Hepatol.* 2011;10(2):119-24. **PubMed | Google Scholar**
24. Prashanth M, Ganesh HK, Vima MV, John M, Bandgar T, Joshi SR, Shah SR, Rathi PM, Joshi AS, Thakkar H, Menon PS, Shah NS. Prevalence of Nonalcoholic fatty liver disease in patients with type 2 Diabetes Mellitus. *J Assoc Physicians India(JAPI).* 2009; 57:205-210. **PubMed | Google Scholar**

25. Leite, NC, Salles GF, Araujo AL, Villela-Nogueira CA, Cardoso CR. Prevalence and associated factors of non-alcoholic fatty liver disease in patients with type-2 diabetes mellitus. *Liver International*. 2009; 29(1):113-119. **PubMed | Google Scholar**
26. Chan WK, Tan AT, Vethakkan SR, Tah PC, Vijayanathan A, Goh KL. Non-alcoholic fatty liver disease in diabetics--prevalence and predictive factors in a multiracial hospital clinic population in Malaysia. *J Gastroenterol Hepatol*. 2013;28(8):1375-83. **PubMed | Google Scholar**
27. Pan J, Fallon MB. Gender and racial differences in non-alcoholic fatty liver disease. *World J Hepatol*. 2014; 6(5):274-283. **PubMed | Google Scholar**
28. Romeo S, Kozlitina J, Xing C, Pertsemlidis A, Cox D, Pennacchio LA, Boerwinkle E, Cohen JC, Hobbs HH. Genetic variation in PNPLA3 confers susceptibility to nonalcoholic fatty liver disease. *Nat Genet*. 2008;40(12):1461-5. **PubMed | Google Scholar**
29. Speliotes EK, Yerges-Armstrong LM, Wu J, Hernandez R, Kim LJ, Palmer CD, Gudnason V, Eiriksdottir G, Garcia ME, Launer LJ, Nalls MA, Clark JM, Mitchell BD, Shuldiner AR, Butler JL, Tomas M, Hoffmann U, Hwang SJ, Massaro JM, O'Donnell CJ, Sahani DV, Salomaa V, Schadt EE, Schwartz SM, Siscovick DS. GOLD Consortium: Genome-wide association analysis identifies variants associated with nonalcoholic fatty liver disease that have distinct effects on metabolic traits. *PLoS Genet*. 2011;7(3):e1001324. **PubMed | Google Scholar**

Table 1: Socio demographic variables of T2DM subjects and controls			
VARIABLES	T2DM Subjects n=168 (%)	Controls n=168(%)	P-Value
AGE mean, years (SD)	53.17±8.6	52.01±8.9	0.1
Gender			0.11
Females	109 (64.9)	97 (57.7)	
Males	59 (35.1)	71 (42.3)	
Level of Education			0.08
None	4 (2.4)	3 (1.8)	
Primary	43 (25.6)	28 (16.7)	
Secondary	58 (34.5)	73 (43.5)	
Tertiary*	63 (37.5)	64 (38.1)	
Monthly income (US \$)			0.10
<300	148(88.1)	149 (88.7)	
>300	20 (11.9)	19 (11.3)	
Concurrent			<0.001 ^a
Hypertension	80 (47.6)	37(22.0)	
Family history of DM	74(44.0)	18(10.7)	
*Inclusive of University, Polytechnic, Teacher training college			
^a - Odds Ratio 3.2, 95% Confidence Interval 2.00-5.17,			
^b Odds Ratio 6.5, 95% Confidence Interval 3.6-11.7			

Table 2: Comparison of anthropometric, laboratory and ultrasound scan parameters between T2DM patients and control subjects

VARIABLE	T2DM subjects Mean±SD, (n=168)	Control subjects Mean±SD, (n=168)	p-value
BMI (kg/ m²)	28.47±4.39	27.42±5.09	0.02
Waist Hip Ratio;			
Females	0.93±0.07	0.89±0.07	0.00
Males	0.94±0.06	0.89±0.07	0.00
Systolic BP (mmHg)	130.41±18.99	123.05±17.74	0.00
Diastolic BP (mmHg)	80.59±12.87	79.92±11.54	0.30
AST (IU/L)	6.49±3.41	5.86± 3.70	0.05
ALT (IU/L)	9.07± 4.34	8.43±7.59	0.17
TCHOL (mg/dl)	179.49±40.89	170 ±31.69.	0.00
LDL-C (mg/dl)	94.46±40.57	81.26±32.94	0.00
HDL-C (mg/dl)	71.07 ±25.25	73.89±23.32	0.14
Triglyceride (mg/dl)	75.22±37.96	80.07±31.47	0.10
Hepatic steatosis Identified on USS (n,%)	28 (16.7%)	2 (1.2%)	0.00*

SD standard deviation, AST, aspartate aminotransferase, ALT alanine aminotransferase, TCHOL total cholesterol, LDL-C Low density lipoprotein, HDL-C high density cholesterol, USS ultrasound scan. *Odds Ratio, 16.6; 95% CI (3.9-70.9)

Table 3: Univariate analysis of the clinical and biochemical factors associated with NAFLD among subjects with T2DM				
Variables	ALL N=168	NAFLD Positive N=28 (%)	NAFLD Negative N=140 (%)	P VALUE
Mean Age, yrs ±SD	53.17 ± 8.6	53.1±8.3	53.2±8.7	0.98
Gender				0.4
Male:	59 (35.1)	12 (20.3)	47 (79.6)	
Female	109 (64.9)	16 (14.7)	93 (85)	
BMI female (≥30 kg/m²)	42 (38.5)	9 (56.3)	33 (35.5)	0.19
BMI Male (≥30 kg/m²)	19 (32.2)	7 (58.3)	12 (25.5)	0.04*
WC Males > 102cm	21 (35.6)	9(75)	12 (25.5)	0.002**
WC Females >88cm	84 (77.1)	14 (87.5)	70 (75.3)	0.35
AST >12 IU/L	8 (4.8)	2(7.1)	6(4.3)	0.40
ALT > 12 IU/L	16 (9.5)	3(10.7)	13(9.3)	0.53
T-CHOL ≥ 200 mg/dl	46 (27.4)	5(17.9)	41(29.3)	0.16
HDL-c				
Female <50 mg/dl	30 (28.3)	5 (31.3)	25 (26.9)	0.76
Male <40 mg/dl	6 (10.2)	4 (33.3)	2 (4.3)	0.013
LDL-c ≥ 100 mg/dl	76 (45.2)	15(53.6)	61(43.6)	0.22
Triglyceride >150mg/dl	10 (5.9)	4 (14.3)	6 (4.3)	0.06

WC waist circumference, BMI Body mass index, AST aspartate aminotransferase, ALT alanine aminotransferase, T-CHOL total cholesterol, LDL-c Low density lipoprotein, HDL-c high density lipoprotein-cholesterol. NAFLD non-alcoholic fatty liver disease.
* Odds ratio 4; 95% CI; 1.1-15.3, ** Odds ratio 8.7; 95% CI:2.1-37