

Images in medicine



Giant anal condyloma

George Kirkilessis^{1,&}, Ioannis Karavokyros²

¹Surgical Department, General Hospital of Pirgos, Pirgos, Greece, ²First Department of Surgery, Medical School, National and Kapodistrian University of Athens, Laikon General Hospital, Athens, Greece

[&]Corresponding author: George Kirkilessis, Surgical Department, General Hospital of Pirgos, Pirgos, Greece

Key words: Anal condyloma, ulcerated tumor mass, condyloma acuminatum

Received: 24 Aug 2019 - Accepted: 02 Sep 2019 - Published: 24 Mar 2020

Pan African Medical Journal. 2020;35:86. doi:10.11604/pamj.2020.35.86.20147

This article is available online at: <http://www.panafrican-med-journal.com/content/article/35/86/full/>

© George Kirkilessis et al. The Pan African Medical Journal - ISSN 1937-8688. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Image in medicine

A 61 years old male patient presented to the Accident and Emergency (AE) Department reporting bleeding during defecation, difficulty and low pain while walking and sitting. On physical examination we saw an exophytic cauliflower like, ulcerated tumor mass measuring 20 x 14 cm that covered the whole perineum and a similar satellite lesion 3 x 4 cm, with malodorousness. The mass was there for at least three years, progressively growing.



Figure 1: a giant anal condyloma of a 61 years old man with the satellite lesion before the operation