

Research



Awareness about stroke among high and low risk individuals in Khartoum, Sudan: a cross-sectional study

Ola Ahmed Abdulmjeed Mohammed, Fatima Abd Alraheem Osman Ahmed, Abubaker Emadeldin Adlan Koko, Sufian Elshafee Osman Khalifa, Hind Abdelaziz Mohamed Abdelaziz, Mohamed Elmojtaba Adil Mohamed, Francis Harrington, Sulaf Ibrahim Abdelaziz, Ihab Babiker Abdalrahman

Corresponding author: Ola Ahmed Abdulmjeed Mohammed, Soba University Hospital, Khartoum, Sudan. olamageed95@gmail.com

Received: 26 Apr 2020 - **Accepted:** 04 May 2020 - **Published:** 21 Aug 2020

Keywords: Stroke, awareness, risk factors, warning symptoms

Copyright: Ola Ahmed Abdulmjeed Mohammed et al. Pan African Medical Journal (ISSN: 1937-8688). This is an Open Access article distributed under the terms of the Creative Commons Attribution International 4.0 License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cite this article: Ola Ahmed Abdulmjeed Mohammed et al. Awareness about stroke among high and low risk individuals in Khartoum, Sudan: a cross-sectional study. Pan African Medical Journal. 2020;36:318. 10.11604/pamj.2020.36.318.23107

Available online at: <https://www.panafrican-med-journal.com/content/article/36/318/full>

Awareness about stroke among high and low risk individuals in Khartoum, Sudan: a cross-sectional study

Ola Ahmed Abdulmjeed Mohammed^{1,&}, Fatima Abd Alraheem Osman Ahmed², Abubaker Emadeldin Adlan Koko², Sufian Elshafee Osman Khalifa³, Hind Abdelaziz Mohamed Abdelaziz⁴, Mohamed Elmojtaba Adil Mohamed³, Francis Harrington⁵, Sulaf Ibrahim Abdelaziz², Ihab Babiker Abdalrahman²

¹Soba University Hospital, Khartoum, Sudan, ²Faculty of Medicine, University of Khartoum, Khartoum, Sudan, ³Faculty of Medicine, Omdurman Islamic University, Khartoum, Sudan, ⁴Academy Charity Teaching Hospital, Khartoum, Sudan, ⁵Royal Cornwall Hospital, Truro, United Kingdom

&Corresponding author

Ola Ahmed Abdulmjeed Mohammed, Soba University Hospital, Khartoum, Sudan

Abstract

Introduction: stroke causes 10.17% of all deaths in Sudan. Levels of stroke awareness amongst patients in Sudan are unknown. The aim of this study is to assess the level of awareness of stroke risk factors, symptoms and immediate management amongst high and low risk patients.

Methods: using descriptive cross-sectional study, participants of high and low risk groups were recruited from the referral clinics of three tertiary hospitals in Khartoum. Data was collected through interviews using structured questionnaire. Knowledge score was devised to assess the awareness about stroke symptoms, risk factors, and management. **Results:** of the 286 participants, 150 were females. The mean age was 44.66 years. About 79.4% reported that stroke is preventable. Hypertension was the most identified risk factor (71.6%) while genetics (0.2%) and alcohol (0.2%) were the least identified risk factors. Twenty-seven percent (27.6%) did not know any stroke risk factors, while 32.9% did not know any warning symptoms. Paralysis of one side of the body was the most identified warning symptom (30.7%). The mean awareness score was 21.9 ± 3.4 among the high risk group vs. 22.1 ± 3.6 among the low risk group with no statistically significant difference between the two groups ($p = .717$). The mean awareness score was statistically associated with the level of education ($p < 0.001$). **Conclusion:** the awareness level was relatively low and not statistically different between high and low risk groups. We recommend the development of an effective educational program for the whole community.

Introduction

Stroke is the second most common cause of death, and it is a major cause of disability. Approximately, two thirds of deaths occur in low and middle income countries. Globally, 70% of strokes and 87% of both stroke-related deaths and disability-adjusted life years occur in low and middle income

countries. Over the last four decades, stroke incidence in low and middle income countries has increased more than double [1]. Non communicable diseases such as stroke are becoming more common in Africa [2]. Stroke services are less developed in most African countries [3] and are in their infancy in Sudan. According to the latest World Health Organization (WHO) data published in 2017, stroke deaths in Sudan reached 27,222 or 10.17% of total deaths. The age adjusted death rate is 136.47 per 100,000 of the population which ranks Sudan in the 27th position in the world regarding deaths from stroke [4]. A study done among patients with acute stroke in Khartoum teaching hospital showed that mortality from stroke in Sudan is higher than in other countries [5]. There are multiple causes of stroke including: embolization, arterial thrombosis, and hemorrhage [6]. Less common causes are venous infarction, carotid or vertebral artery dissection, polycythemia and hyper-viscosity syndromes, fat and air embolism. Common risk factors of stroke are: hypertension, smoking, lifestyle, increased hematocrit, raised cholesterol, atrial fibrillation, obesity, diabetes, and severe carotid stenosis. Rapid identification, quick transfer to medical care and immediate and appropriate medical care are key factors in improving outcome of stroke [6]. Awareness about stroke amongst patients, caretakers and medical staff has been studied in some African countries [3,7-13], most of which revealed a generally poor level of awareness. Locally, levels of stroke awareness amongst medical staff, patients and caretakers in Sudan are unknown. Studies assessing awareness regarding other chronic diseases like hypertension showed low awareness and lack of adherence to medications [14]. Risk factors for stroke are becoming more prevalent among the Sudanese community according to the STEPwise approach to surveillance survey of 2005 (STEPS) [15]. In addition, a low level of knowledge in patients has resulted in sub-optimal adherence to risk modifications [16].

Rationale: stroke is a preventable disease, with prevention strategies mainly based on knowledge about stroke risk factors and their modification, recognition of stroke warning symptoms and early hospital arrival which will all improve the outcome [17]. It is expected for those who are at high risk of developing stroke to have more knowledge about stroke risk factors, warning symptoms and treatment than the low risk population [18], so this study will assess the potential difference in knowledge between the two groups, which will also give an indication of whether or not the role of medical practitioners in educating patients is well-activated. In addition, there are no published studies from Sudan comparing awareness about stroke symptoms; risk factors and management among high and low risk groups.

Study aim: the aim of this study was to assess the level of awareness of stroke risk factors, symptoms and immediate management amongst high and low risk patients attending the clinics at 3 hospitals in Khartoum. In addition, we aimed to investigate the potential difference between the two groups' levels of awareness. The findings of this study aim to inform the design and delivery of public stroke educational programs.

Methods

Study design and setting: this is a descriptive cross-sectional hospital-based study. Participants were recruited from outpatient clinics within three of the largest tertiary hospitals in Khartoum State (Omdurman teaching hospital, Bahri teaching hospital and Soba university hospital). The study targeted patients with high (hypertension, type 2 diabetes, cardiac diseases), and low or no risk for development of stroke. Inclusion criteria were adult Sudanese males and females diagnosed with hypertension, or diabetes attending the follow up clinics at Soba, Bahri or Omdurman teaching hospitals; and adult Sudanese males and females with no identified risk of stroke attending the surgery, paediatrics and obstetrics referral clinics as

co-patients. Patients attending the clinics during the period from April 2018 to July 2018 were approached to be included in the study if they met the inclusion criteria. Participants were recruited and interviewed by the authors. The sample size was calculated to be 290 using Epi Info 7 software, taking into consideration a prevalence of 22% [10], acceptable margin of error of 5%, design effect of 1, and a 10% non-response rate. Only 4 subjects refused to participate in the study resulting in a non-response rate of 1.4%.

Data collection and analysis: data was collected through structured face to face interviews using standardized questionnaire that was used in a previous study [13]. The questionnaire was pretested in a pilot study of 30 patients. A knowledge score was devised assessing knowledge about organ affection, risk factors, and warning signs of stroke. The score was computed from 45 questions. Questions assessing organ affection and general information were 11 questions; and those assessing risk factors were 14, while those assessing symptoms and warning signs were 20. Each correct option was given 1 score, while each wrong option was given 0, so that the total score would be 45. Internal consistency of the scale was assessed using Cronbach's alpha which was found to be 0.7. Data was entered and analysed using SPSS [statistical package social science] version 21.0. Descriptive statistics were conducted and presented as frequency tables, means, medians [Md], and standard deviations [SD]. Since the data was not normally distributed, non-parametric tests [Mann-Whitney U and Kruskal-Wallis tests] were performed to assess the difference between groups regarding knowledge about stroke, a p-value < 0.05 was considered significant for all purposes.

Ethical consideration: ethical approval was obtained from Soba Centre for Audit and Research, and Khartoum State ministry of health research department and from each hospital's administration. Research purpose and objectives were explained to the participants verbally. Written informed consent was obtained from participants.

Participants had the right to withdraw at any time without any detriment to their care.

Results

Demographic data: a total of 286 participants were included in this study, the mean age was 44.66 ranging from 18 to 90 years, with the most frequently reported age range being 41-50 years (23.8%). Fifty-four percent of the participants (150) were females. The majority (52.4%) had attained secondary school or university education. Demographic characteristics of the participants are shown in Table 1. Sixty-six percent of the participants (190) resided in Khartoum State; 35% (100) were from Soba university hospital, 33.6% (93) were from Omdurman teaching hospital and 31.5% were from Bahri teaching hospital. Participants were divided into low risk group 48.6% (139) and high risk group 51.4% (147) of the total sample. As shown in Figure 1 the most prevalent chronic medical condition among participants was diabetes. Reported sources of information regarding stroke were family and friends 58.7% (206), TV 14.8% (52), health care providers 10% (35), electronic social media (Facebook, Twitter, WhatsApp) 1.1% (4), and others 5.7% (20) (Figure 2). Only 10.8% (31) of the respondents had a personal experience of stroke while 22.4% (64) had an experience of stroke in a first degree relative.

Knowledge about stroke: about 44% of the participants correctly identified the brain as the body organ affected by stroke, 22.8% of them identified the limbs, and 16% did not know which body organ is affected by stroke. Two hundred and twenty-seven of the participants (79.4%) reported that stroke is a preventable disease, 73.1% reported that a person can have a stroke more than once, and the majority of the participants (92%) believe that having a stroke affects daily activities.

Stroke risk factors: as shown in Table 2, the most commonly identified risk factor by 71.6% (149/207) is hypertension followed by diabetes (17.1%). Only

0.2% identified genetics as a risk factor for stroke, similar with alcohol (0.2%), while only 0.4% identified lack of exercise and 0.8% identified cancer as risk factors, furthermore none identified oral contraceptives as a risk factor. Twenty-seven percent (27.6%) (79/286) did not know any risk factors of stroke.

Stroke warning symptoms: ninety-four (32.9%) of the study participants did not know any stroke warning symptoms. Of the 192 (67.1%) who knew about stroke warning symptoms, paralysis of one side of the body was the most commonly identified warning symptom by 30.7% (59/192), followed by sudden difficulty in speaking or understanding 27.1% (52/192) and dizziness 20.3% (39/192).

Planned response to stroke: two hundred and one (70.0%) of the participants reported that they would go to hospital if stroke symptoms were experienced; while 46 of them (16.3%) would react by giving self-medications, specifically aspirin (Table 3).

Difference in knowledge about stroke between high risk and low risk groups: the mean awareness score was found to be 22 ± 3.48 of a total of 45 for the whole sample, ranging from 15 to 30 ($n = 286$). Mean awareness score for high risk group was 21.9, and for the low risk group 22.1 with no statistically significant difference between the two groups according to Mann-Whitney U test ($p = .717$). In addition, no significant difference in awareness scores was detected between males ($Md = 23$, $n = 127$) and females ($Md = 22$, $n = 150$), $p = .894$. Also, a Kruskal-Wallis test indicated a statistically significant difference in the awareness scores between different levels of education as greater level of education correlated with higher scores (None: $n = 66$, $Md = 19.5$; Primary: $n = 70$, $Md = 21$; Secondary: $n = 93$, $Md = 23$; University: $n = 57$, $Md = 24$), $\chi^2 (3, n = 286) = 46.53$, $p < 0.001$.

Discussion

The study identified a low level of stroke knowledge among the participants. No statistically significant difference was identified between high and low risk groups. The mean score of knowledge was 22 out of 45, reflecting a suboptimal or below average level of awareness among both high and low risk individuals. This is similar to other studies done in African countries where predictors of good knowledge included male sex and educational status [7,12], while areas of poor knowledge included risk factors [13] and symptoms [12]. However, some studies reported contradicting results of good knowledge among high risk patients [8]. The most reported source of knowledge was family and friends, which is similar to a study from Pakistan [19], reflecting that most of the knowledge is gained from non-professional medical sources. Though this may affect the reliability of information delivered, it represents a great opportunity to educate families and communities on a much wider base.

In our study, only 44% identified the brain as the organ affected by stroke. It is significantly lower than studies in Sokoto, Nigeria [8] and Taif, Saudi Arabia [20] (87.1% and 81.1%, respectively), but approximately similar to other studies in Kampala, Uganda and Benin city, Nigeria in which 51.8% and 56.3% respectively correctly mentioned the brain as the organ affected in stroke [10,12]. However, it was significantly higher compared to rural Uganda [13] and Oman [21] (26.1% and 35%, respectively). Hypertension is the single most important stroke risk factor and was the most commonly identified risk factor in our study as well as other studies [20,22,23]. Diabetes is a chronic systemic disease with wide range of complications that directly correlate with poor management and control. The incidence and prevalence of diabetes is rising [24], and stroke is a known complication of diabetes. A study from Uganda [13] showed that 8% were able to link diabetes to stroke, while a similar study from Nigeria [7] showed that only 2.9% of the participants associated stroke with diabetes. In our

study 29.4% of the population was able to make the association. The common stroke symptoms of weakness, paralysis and speech difficulties were identified most commonly in our study as well as in other studies [8,13,22].

Assessing the planned response to a stroke event by a patient or a witness is very important in any stroke awareness assessment study. We found that 70% of the participants would seek medical care and transfer the patient to the nearest health care facility. Nevertheless, a worrying percentage had some misbeliefs about dealing with an acute event as 16.3% stated that they would give the patient aspirin pills which may be very harmful in cases of hemorrhagic stroke. Many studies have addressed the importance of education in primary and secondary stroke prevention [25-27]. The role of education in primary and secondary prevention is well established. This role is far greater with non-communicable diseases [17]. Becoming educated about the risks and ways to reduce risk factors can decrease the number of strokes which occur each year. In some instances, people may not be ignoring their health- they may simply be unaware that they are at risk [28]. Community education regarding recognition of stroke can be an excellent way to increase public awareness and decrease mortality due to stroke.

Conclusion

Awareness level was relatively low and not statistically different between high and low risk groups. The most identified risk factors were hypertension and diabetes; the most identified symptom was paralysis of one side of the body. Knowledge score was statistically associated with the level of education. Although being a small study, it covers a broad range of patients, both with and without risk factors, with diverse educational levels and age (ranging from 18 to 90 years) so it does give a good grasp of the state of knowledge about stroke. It is also the first study assessing knowledge about stroke in Sudan. We recommend that efforts should be concentrated on developing

an educational program. We plan to use the low levels of stroke knowledge and awareness found in this study to inform the design of an effective stroke educational program targeting both high and low risk groups. We also plan to study the medical care of patients with acute stroke, when they present to hospital, how they are managed and followed as in and outpatients. Of the limitations we faced was the fact that although the questionnaire is validated the score has never been used before, so we scored, pretested and validated it for this study. This led to the fact that scores could not be compared to other studies in which this questionnaire was used. Also, the study covered patients mostly living in the central Sudan area; Khartoum, not representing those who live in other areas of Sudan, which are mostly rural areas. Furthermore, assessors were not completely blinded and separated to analysers.

What is known about this topic

- *Stroke is a major cause of morbidity and mortality, and incidence in Africa is increasing;*
- *Causes 10.17% of deaths in Sudan;*
- *No previous studies assessing its awareness among patients in Sudan.*

What this study adds

- *Assess the awareness level of stroke among patients;*
- *Compare the awareness level among high and low risk patients.*

Competing interests

The authors declare no competing interests.

Authors' contributions

Francis Harrington, Ihab Abdelrahman, Sulaf Abdelaziz formulated the research idea; Francis Harrington, Sulaf Abdelaziz, Ihab Abdelrahman, Ola Mohammed, Fatima Ahmed, Abubaker Koko, and Mohamed Mohamed designed the study; Sufian Alshafee, Mohamed Mohamed, Hind Abdelaziz, Ola

Mohammed, Fatima Ahmed, Abubaker Koko collected data; Abubaker Koko, Fatima Ahmed, Ola Mohammed, Sufian Alshafee, Mohamed Mohamed, Hind Abdelaziz undertook the data analysis and interpretation: Ola Mohammed, Fatima Ahmed, Abubaker Koko, Francis Harrington, Ihab Abdelrahman drafted the manuscript. All authors read and approved the final version of the manuscript.

Acknowledgments

The authors would like to acknowledge Liz Tremayne-Ward, Stroke Occupational Therapist (Royal Cornwall Hospital, Truro, UK, Liz.tremayne-ward@nhs.net) for her invaluable comments on the making of this manuscript, we also like to convey thanks to Mohammed K.A. Saad (university of Khartoum, faculty of medicine) for English language revision and proofreading.

Tables and figures

Table 1: demographic characteristics of the study participants

Table 2: differences in knowledge between high and low risk groups

Table 3: planned response to an event of stroke among the respondents

Figure 1: chronic medical conditions among the high risk group

Figure 2: source of information regarding stroke among respondents

References

1. Johnson W, Onuma O, Owolabi M, Sachdev S. Stroke: a global response is needed. Bull World Health Organ. 2016 Sep 1;94(9):634-634A. **PubMed** | **Google Scholar**
2. Moffat Nyirenda J. Non-communicable diseases in sub-Saharan Africa: understanding the drivers of the epidemic to inform intervention strategies. International Health. 2016 May;8(3):157-8. **PubMed** | **Google Scholar**

3. Bertha Chioma Ekeh. Challenges of the management of stroke in sub-Saharan Africa: evaluating awareness. *Journal of Pediatric Neurology and Medicine*. January 2017.
4. World Health Ranking. South Sudan: stroke. World Life Expectancy. 2018.
5. El Zein A, Bukhari E, Homeida S, Adam I. Stroke in CT-scan Department of Khartoum Hospital, Sudan. *Tropical Doctor*. 2007;37(4):244-245. **PubMed | Google Scholar**
6. Parveen Kumar Michael Clark. Kumar & Clark's Clinical Medicine. Imprint: Elsevier. 28th July 2016; 9th ed:1456.
7. Vincent-Onabajo G, Moses T. Knowledge of stroke risk factors among stroke survivors in Nigeria. *Stroke Res Treat*. 2016;2016:1902151. **PubMed | Google Scholar**
8. Sarafadeen Adeniyi Ariseji, Kehinde Joseph Awosan, Mansur Oche Oche, Anas Ahmad Sabir, Mohammed Taofeek Ibrahim. Knowledge and practices related to stroke prevention among hypertensive and diabetic patients attending specialist hospital, Sokoto, Nigeria. *Pan African Medical Journal*. 2018;29:63. **PubMed | Google Scholar**
9. Leopold Ndemnge Aminde, Noah Takah, Calypse Ngwasiri, Jean Jacques Noubiap, Maxime Tindong, Anastase Dzudie *et al*. Population awareness of cardiovascular disease and its risk factors in Buea, Cameroon. *BMC Public Health*. 2017 Jun 5;17(1):545. **PubMed | Google Scholar**
10. Mark Kaddumukasa, James Kayima, Jane Nakibuuka, Leviticus Mugenyi, Edward Ddumba, Carol Blixen *et al*. A cross-sectional population survey on stroke knowledge and attitudes in Greater Kampala, Uganda. *Cogent Med*. 2017;4(1):1327129. **PubMed | Google Scholar**
11. Akinyemi R, Ogah O, Ogundipe R, Oyesola O, Oyadoke A, Ogunlana M *et al*. Knowledge and perception of stroke amongst hospital workers in an African community. *European Journal of Neurology*. 2009;16(9):998-1003. **PubMed | Google Scholar**
12. Odiase Francis Ehidiamen, Ogbemudia Judith Ehinwenma. Awareness of stroke risk factors and warning symptoms amongst hypertensive patients in Benin city. *Ann Med Health Sci Res*. 2018; 8: 40-44. **Google Scholar**
13. Kaddumukasa M, Kayima J, Kaddumukasa MN, Ddumba E, Mugenyi L, Pundik S *et al*. Knowledge, attitudes and perceptions of stroke: a cross-sectional survey in rural and urban Uganda. *BMC Res Notes*. 2015;8:819. **Google Scholar**
14. Suliman A. The state of heart disease in Sudan. *Cardiovasc J Afr*. 2011;22(4):191-196. **PubMed | Google Scholar**
15. World Health Organization. Sudan-Khartoum STEPS Noncommunicable Disease Risk Factors Survey 2005-2006. World Health Organization. Accessed May 01 2020
16. Slark J, Sharma P. Risk awareness in secondary stroke prevention: a review of the literature. *JRSM Cardiovasc Dis*. 2014;3: 2048004013514737. **PubMed | Google Scholar**
17. Green T, Haley E, Eliasziw M, Hoyte K. Education in stroke prevention: efficacy of an educational counselling intervention to increase knowledge in stroke survivors. *Canadian Journal of Neuroscience Nursing*. 2007;29(2):13-20. **PubMed | Google Scholar**
18. Saengsuwan J, Suangpho P, Tiamkao S. Knowledge of stroke risk factors and warning signs in patients with recurrent stroke or recurrent transient ischaemic attack in Thailand. *Neurol Res Int*. 2017;2017:8215726. **PubMed | Google Scholar**

19. Aly Z, Abbas K, Kazim S, Taj F, Aziz F, Irfan A *et al.* Awareness of stroke risk factors, signs and treatment in a Pakistani population. *Journal Of Pakistan Medical Association.* 2009;59(7):495-499. **PubMed | Google Scholar**
20. Alzahrani Y, Alomairi N, Althomali M, Althagafi A, Kutbi A, Alzaidi S *et al.* Awareness and knowledge of outpatients in Taif, Saudi Arabia regarding stroke. *World journal of pharmaceutical and medical research.* 2018;4(2):5-10. **Google Scholar**
21. Mohammed Al Shafae A, Shyam Ganguly S, Abdullah Al Asmi. Perception of stroke and knowledge of potential risk factors among Omani patients at increased risk for stroke. *BMC Neurol.* 2006 Oct;6:38. **PubMed | Google Scholar**
22. Kamran S, Bener AB, Deleu D, Khoja W, Jumma M, Al Shubali A *et al.* The level of awareness of stroke risk factors and symptoms in the Gulf Cooperation Council Countries: Gulf cooperation council stroke awareness study. *Neuroepidemiology.* 2007;29(3-4):235-242. **PubMed | Google Scholar**
23. Jarou Z, Harris N, Gill L, Azizi M, Gabasha S, LaBril R *et al.* Public stroke knowledge: those most at risk, least able to identify symptoms. *Medical Student Research Journal.* 2013;3(Fall):3-8. **Google Scholar**
24. Sudan Federal Ministry of Health. Annual Health Statistics Reports 2014. Repub Sudan. 2015;1:313.
25. Siti Noorkhairina S, Sakinah H, Che Rabiaah M. Secondary stroke prevention through patient education intervention on lifestyle risk Factors?: a review. *Heal Environ J.* 2013;4(2):127-151. **Google Scholar**
26. Boden-Albala B, Quarles L. Education strategies for stroke prevention. *Stroke.* 2013;44(6 Supplement 1):S48-S51. **PubMed | Google Scholar**
27. Yan LL, Li C, Chen J, Miranda JJ, Luo R, Bettger J *et al.* Prevention, management, and rehabilitation of stroke in low- and middle-income countries. *ENeurologicalSci.* 2016;2:21-30. **PubMed | Google Scholar**
28. Community Education in Recognition of Stroke. Certification Institute.

Table 1: demographic characteristics of the study participants

| | Character | N | Percent (%) |
|--|-----------------------|----------|--------------------|
| Gender | Male | 127 | 45.8 |
| | Female | 150 | 54.2 |
| Age range | 18-30 | 63 | 22.3 |
| | 31-40 | 54 | 19.1 |
| | 41-50 | 67 | 23.8 |
| | 51-60 | 60 | 21.3 |
| | More than 60 | 38 | 13.5 |
| Residence | Khartoum State | 190 | 66.4 |
| | Gazira State | 63 | 22.0 |
| | Others | 33 | 11.5 |
| Origin | Central Sudan | 137 | 48.1 |
| | Northern Sudan | 32 | 11.2 |
| | Western Sudan | 64 | 22.5 |
| | Eastern Sudan | 13 | 4.6 |
| | Southern Sudan | 39 | 13.7 |
| Marital status | Married | 227 | 79.4 |
| | Single, never married | 50 | 17.5 |
| | Divorced | 3 | 1.0 |
| | Separated | 0 | 0.0 |
| | Widowed | 6 | 2.1 |
| Highest level of education attained | None | 66 | 23.1 |
| | Primary | 70 | 24.5 |
| | Secondary | 93 | 32.5% |
| | University | 57 | 19.9% |

| Table 2: differences in knowledge between high and low risk groups | | | |
|---|------------------------|-----------------------|--------------|
| Knowledge Question | High risk group | Low risk group | p |
| Stroke affects the brain | 83 [56.5%] | 87 [62.5%] | 0.175 |
| Stroke is preventable | 116 [78.9%] | 111 [79.9%] | 0.816 |
| Person can have stroke more than once | 113 [76.9%] | 96 [69.1%] | 0.254 |
| Stroke have an effect on daily activities | 137 [93.2%] | 126 [90.6%] | 0.133 |
| What are the risk factors that you know? | | | |
| Old age | 5 [3.4%] | 8 [5.7%] | 0.251 |
| Hypertension | 75 [53.9%] | 74 [53.2%] | 0.399 |
| Diabetes | 43 [29.2%] | 41 [29.5%] | 0.533 |
| Cigarette smoking | 4 [2.7%] | 5 [3.6%] | 0.465 |
| Heart disease | 6 [4.0%] | 23 [16.5%] | 0.001 |
| Alcohol | 1 [0.7%] | 0 [0%] | 0.33 |
| Atherosclerosis | 13 [8.8%] | 12 [8.6%] | 0.95 |
| High cholesterol | 15 [10.8%] | 9 [6.4%] | 0.256 |
| Obesity | 6 [4.3%] | 11 [7.9%] | 0.214 |
| Stress | 33 [22.4%] | 34 [24.4%] | 0.688 |
| Genetics [hereditary] | 0 [0%] | 1 [0.7%] | 0.486 |
| Lack of exercise | 0 [0%] | 2 [1.4%] | 0.235 |
| Poor hygiene | 0 [0%] | 0 [0%] | |
| Headache [migraine] | 1 [0.7] | 3 [2.2%] | 0.359 |
| Cancer | 0 [0%] | 4 [2.9%] | 0.055 |
| Bad diet | 13 [8.8%] | 2 [1.4%] | 0.005 |
| Oral contraceptives | 0 [0%] | 0 [0%] | |
| Tremors | 1 [0.7] | 0 [0%] | 0.33 |
| Don't know | 42 [28.6%] | 37 [26.6%] | |
| What are the warning signs that you know? | | | |
| Dizziness | 15 [10.2%] | 24 [17.2%] | 0.082 |
| Blurred or double vision or loss of vision | 6 [4.0%] | 2 [1.4%] | 0.284 |
| Headache | 23 [15.6%] | 15 [10.8%] | 0.227 |
| Sudden difficulty in speaking or understanding or reading | 24 [16.3%] | 28 [20.1%] | 0.403 |
| Tiredness | 5 [3.4%] | 10 [7.2%] | 0.15 |
| Fever/sweating | 2 [1.4%] | 4 [2.9%] | 0.437 |
| Shortness of breath | 5 [3.4%] | 11 [7.9%] | 0.097 |
| Chest pain/chest tightness | 10 [6.8%] | 13 [9.4%] | 0.428 |
| Nausea and vomiting | 3 [2.0%] | 0 [0%] | 0.248 |
| Weakness of any part of the body | 10 [6.9%] | 22 [15.8%] | 0.016 |
| Weakness of one side of the body | 16 [10.9%] | 14 [10%] | 0.823 |
| Paralysis of any part of the body | 16 [10.9%] | 21 [15.1%] | 0.287 |
| Paralysis of one side of the body | 32 [21.8%] | 27 [19.4%] | 0.624 |
| Fainting, black out, collapse | 14 [9.5%] | 8 [5.7%] | 0.232 |
| Numbness tingling of any body part | 19 [12.9%] | 7 [5.0%] | 0.02 |
| Numbness tingling of one side of the body | 7 [4.8%] | 3 [2.1%] | 0.337 |
| Don't know | 49 [33.3%] | 45 [32.4%] | |
| Bold p-values indicate statistically significant difference between high and low risk groups [p < 0.05] | | | |

Table 3: planned response to an event of stroke among the respondents

| | | N | Percent |
|---|---|----------|----------------|
| What would be your planned response to a stroke ? | Ask family members or relatives to help | 3 | 1.1% |
| | Call general practitioner or family doctor | 19 | 6.7% |
| | Go to chemist for advice or medication | 3 | 1.1% |
| | Self-medication with aspirin | 46 | 16.3% |
| | Ask friend or neighbors for help | 8 | 2.8% |
| | Go to hospital | 201 | 70.2% |
| | Visit community health center | 9 | 3.2% |
| | Visit alternative health care providers [herbal med, traditional healers] | 2 | 0.7% |
| | Seek spiritual healing [prayer] | 2 | 0.7% |
| | Combination of hospital and tradition | 1 | 0.4% |
| | Call a Physiotherapist | 2 | 0.7% |
| | Don't know | 26 | 9.2% |
| | Others | 17 | 6.0% |




Figure 1: chronic medical conditions among the high risk group




Figure 2: source of information regarding stroke among respondents