

Supplementary Information, *S. Afr. J. Chem.*

***In vitro* investigation of the antimicrobial activity of a series of lipophilic phenols and naphthols**

Thavendran Govender,^a Usha Govinden,^a Chunderika Mocktar,^a Hendrik G. Kruger,^a Jelena Veljković,^b Nikola Cindro,^b Damir Bobinac,^b Ivana Žabčić,^b Kata Mlinarić-Majerski,^b Nikola Basarić^{b*}

^a Catalysis and Peptide Research Unit, School of Health Sciences, University of KwaZulu-Natal, Durban, South Africa. Fax: +27 31 2603091;

^b Department of Organic Chemistry and Biochemistry, Ruđer Bošković Institute, Bijenička cesta 54, 10 000 Zagreb, Croatia. Fax: +385 1 4680 195; E-mail: nbasari@irb.hr

TABLE OF CONTENTS

1. Zones of inhibition of compounds (Table S1)	S2
--	----

Table S1: Zones of inhibition (in mm) of the compounds (n=3)

Compound	<i>Staphylococcus aureus</i> ATCC 25923	SD	<i>Bacillus subtilis</i> ATCC 6633	SD	<i>Candida albicans</i> ATCC 10231	SD	<i>Escherichia coli</i> ATCC 25922	SD
3	-		-		8.83	0.29	-	
14	14.33	0.6	24.33	0.6	-		-	
16	13.33	0.5	17.67	0.6	-		-	
20	14.67	0.6	15.33	0.5	-		-	
21	10.33	0.6	13.33	0.6	-		-	
23	17.33	0.6	16.33	0.6	-		-	
25	9	0.1	-		-		-	
31	13.67	0.6	13.67	0.5	-		-	
37	13.67	0.5	18.33	0.6	-		10.33	0.6
41	20.33	0.6	19.67	0.6	11.67	0.6	-	

Compounds were dissolved in DMSO and DMSO was used as a control.

(-) indicates no compound activity