

Saw-toothed fish bone ingestion: A method for propulsion

EMILY BROADIS, M.B. CH.B., M.R.C.S. (GLASG.)

WAKISA MULWAFU, F.C.O.R.L. (S.A.)

Department of Surgery, College of Medicine, Chichiri, Private Bag 414, Blantyre 3, Malawi

Summary

We describe the case of an 11-year-old girl who presented with a lump in her neck. On exploration a 4 cm saw-toothed fish bone was extracted from the soft tissue of the anterior triangle of the neck.

Ingested fish bones are a common presentation in accident and emergency departments, but there are only a small number of reports of migration through the pharynx into soft tissues. Rare complications can include oesophageal perforation, haematoma formation, retropharyngeal abscesses and recurrent infections. To the best of our knowledge this is the first reported case of migration of an ingested fish bone in Malawi.

Case report

An 11-year-old girl presented at our department with a swelling in the anterior triangle of the neck. She had first noticed it 9 days previously, and she denied any pain, discharge or changes in size of the swelling. She was otherwise well with no significant past medical history. Interestingly, on further questioning she reported having swallowed a fish bone 2½ weeks earlier. This had caused severe pain, but after the pain subsided she was able to continue eating with no further problems.

On examination she was a well-looking, well-nourished child with normal haemodynamic parameters. She had a small (1×1 cm), firm, non-tender lump at the anterior border of the right sternocleidomastoid muscle, at the junction of its upper third and lower two-thirds. It was relatively immobile and gave the impression of extending deeper. The skin moved over it easily and there were no skin changes or warmth.

Our differential diagnoses included branchial arch remnant, cervical rib, lymph node or ingested migrated fish bone. However, the swelling was considered to be too anterior for a cervical rib, and she was at the upper end of the age group in which we would have expected a branchial arch remnant to present.

A lateral cervical radiograph was performed and was unremarkable.

The patient was taken to theatre for exploration, where a 4 cm saw-toothed fish bone (Fig. 1) was discovered just beneath



Fig. 1. Posteriorly angled spinous processes.

the skin, protruding from under the anterior border of the right sternocleidomastoid muscle. The wound was washed out and sutured. She made an uneventful postoperative recovery and was discharged home the following day.

Discussion

Although ingested fish bones are a relatively common presentation to accident and emergency departments, migration of the bone through the pharynx is a rare complication. Most reports are from East Asia and South Korea, where fish are commonly consumed.^{1,2}

This report emphasises the importance of proper history taking in the diagnosis of ingestion of foreign bodies. If a patient reports having ingested a foreign body, the clinician must maintain a high index of suspicion and should endeavour to obtain a full history.

Complications of migration include oesophageal perforation, retropharyngeal abscess formation, chronic infections and haematoma formation. Occasionally the bone may migrate through the tissues, causing no damage.³

Lake Malawi provides a source of fish for many of the people who live along its shores. Our patient described the fish as a *nkholokolo*, which is thought to be a local name for *Synodontis njassae*.⁴ We believe that the saw-toothed spine enabled propulsion of the bone through the tissues in one direction only. With the spike at the end the tissues could be pierced easily, and as the soft tissues moved around the bone it could be pushed forwards but was unable to move backwards because of the posteriorly angled spines.

A computed tomography scan is often recommended for investigation of possible fish bone ingestion if the examination and plain radiograph are unhelpful.^{5,6} However, because of

the superficial location of the lump and the clear history from the patient and her guardian in our case, we did not feel it was necessary to investigate further prior to exploration.

Conclusion

This report emphasises the importance of proper history taking in the diagnosis of ingestion of foreign bodies. To the best of our knowledge this is the first reported case of migration of an ingested fish bone in Malawi.

REFERENCES

1. Chung SM, Kim HS, Park EH. Migrating pharyngeal foreign bodies: a series of four cases of saw-toothed fish bones. *Eur Arch Otorhinolaryngol* 2008; 265: 1125-1129.
2. Ng SK, Tong MC. A swallowed fish bone. *Hong Kong Med J* 2009; 15: 73-74.
3. Maseda E, Ablanedo A, Baldó C, Fernández MJ. Migration and extrusion from the upper digestive tract to the skin of the neck of a foreign body (fish bone). *Acta Otorrinolaringol Esp* 2006; 57: 474-476.
4. Ambali A, Kabwazi H, Malekano L, et al. Relationship between local and scientific names of fishes in Lake Malawi. *African Study Monographs* 2001; 22: 123-154.
5. Akazawa Y, Watanabe S, Nobukiyo S, et al. The management of possible fish-bone ingestion. *Auris Nasus Larynx* 2004; 31: 413-416.
6. Braverman I, Gomori JM, Poly O, Saah D. The role of CT imaging in the evaluation of cervical esophageal foreign bodies. *J Otolaryngol* 1993; 22: 311-314.



SAGES 2010 | 29 July - 1 August

Dear Colleagues,

We invite you to attend the combined SAGES SASES meeting taking place in Johannesburg, Sandton Convention Centre. We have many prestigious international speakers, including a group from Johns Hopkins University and the academic programme promises to be exciting which will include a live workshop. We offer a platform for free papers and we eagerly anticipate hearing about new research being conducted in South Africa.

Reid Ally
Reid Ally
Chairman SAGES 2010

CONFIRMED INTERNATIONAL SPEAKERS

- Willem Bemelman, Academic Medical Center, Amsterdam
- Steven Brant, Meyeroff Inflammatory Bowel Disease Genetics Laboratory, USA
- Marcia (Mimi) Canto, Johns Hopkins University, USA
- Jacques Deviere, Erasme University Hospital, Brussels, Belgium
- Paul Fockens, Academic Medical Center, Amsterdam
- Charles Blanke, University of British Columbia, Vancouver, Canada
- Martin Goetz, University of Mainz, Germany
- Anthony Kalloo, Johns Hopkins University, USA
- Zhiping Li, Johns Hopkins University, USA
- Patrick Marcellin, Beaujon Hospital and University of Paris, France
- Martin (Marty) Makary, Johns Hopkins University, USA
- Chris Mulder, University Medical Center, Amsterdam
- Patrick Okolo, Johns Hopkins University, USA

PARTICIPATING SOCIETIES

- SA Gastroenterology Society
- SA Society of Endoscopic Surgery
- American Society of Gastroenterology
- SA Gastrointestinal Nurses Society

WORKSHOP – Friday 30 July 2010

Procedures will be televised live from the operating theatres at Donald Gordon Medical Centre to delegates at the Sandton Convention Centre. Watch the website for updated details.

LOCATION

Johannesburg, Joburg, Jozi or Egoli. Our city has as many names as it does faces. Vibrant, energetic and dynamic, its diversity is both intriguing and exciting; this is a city of contrasts where the soul of Africa is fused with eastern and western influences.

One of the 40 largest metropolitan areas in the world, our city is the financial and commercial hub of Africa; cultures from every nation unite in this city that attracts people from all over the globe drawn by economic opportunity and a fast-paced urban lifestyle.

Whether for business or pleasure, whatever the length of your stay, Joburg has something for everyone; the very best conference venues, spas of splendour, colourful casinos, amazing golf courses, majestic malls and trendy hotels, all fused with a unique urban African essence. Fall in love with Joburg – the sound, the rhythm, the soul and the passion of Africa.

VENUE

Sandton Convention Centre is Southern Africa's international leading and most prestigious, multi-purposes exhibition and convention centre.

Situated in northern Johannesburg, in the heart of Sandton's business, hotel and entertainment district, Sandton Convention Centre offers easy access to over 5 000 hotel rooms, is adjacent to the country's top shopping and entertainment complexes and provides nearly 11 000 secure parking bays.

EXHIBITION & SPONSORSHIP

Please contact Tanya Schmidt for details of sponsorship and exhibition packages on +27 (0) 41 374 5654 or at Tanya@easternsun.co.za. Details will also be on the congress website www.sages.co.za

ACCOMMODATION

Special rates will be negotiated and block bookings made for congress delegates in the vicinity of the convention centre.

CONFERENCE ORGANISERS

Eastern Sun Events: P O Box 12612, Centrahil, 6006, South Africa.
Tel: +27 (0) 41 374 5654. Fax: +27 (0) 41 373 2042.
Email: info@easternsun.co.za

SAGES SECRETARIAT

Karin Fenton: Tel: +27 (0) 21 404 3062 (9am-2pm).
Fax: +27 (0) 21 447 0582. Email: karin.fenton@uct.ac.za

for more information visit WWW.sages.co.za

AstraZeneca World Leaders in Gastroenterology
AstraZeneca Pharmaceuticals (Pty) Ltd. Reg. No. 1992/005854/07. 5 Leuwakop Road, Sunninghill, 2157. Tel: (011) 797-6000. Fax: (011) 797-6001. www.astrazeneca.co.za
Nexiam® 20 mg, Reg. No. 35/11.4.3/0263, Nexiam® 40 mg, Reg. No. 35/11.4.3/0264, Nexiam® IV 40 mg, Reg. No. A39/11.4.3/0394. For full prescribing information refer to Nexiam® package inserts for MUPS and IV respectively. EPI date MUPS: 14/01/2006. EPI date IV: 06/03/2009. Date compiled: May 2010.
A HASA member company

Nexiam
esomeprazole

20mg 40mg iv