

Oesophageal atresia without tracheo-oesophageal fistula and an anorectal malformation: Advantages of a primary laparoscopically assisted anorectal pull-through

M. ARNOLD, M.B. CH.B., D.C.H.

D. SIDLER, M.D. (BERNE), M.PHIL. (APPLIED ETHICS), F.C.S. (S.A.)

S. W. MOORE, M.B. CH.B., F.R.C.S. (ED.), M.D.

Division of Paediatric Surgery, Stellenbosch University and Tygerberg Children's Hospital, Tygerberg, W Cape

Summary

We report on a primary laparoscopically assisted anorectal pull-through (LAARP) performed in a neonate with pure oesophageal atresia and imperforate anus with recto-bulbo-urethral fistula, representing a unique case for the LAARP approach owing to the undistended nature of the bowel and sterile meconium. Further evaluation of the applicability of LAARP in the management of infants with anorectal malformations is needed, but in this case it held major advantages for the patient. A laparoscopically assisted gastrostomy was facilitated during the same procedure, while avoidance of a colostomy and its associated complications also facilitated preservation of the left colon for subsequent use in oesophageal replacement. Long-term outcome remains to be assessed.

Success with posterior sagittal anorectoplasty and Pfannenstiel minilaparotomy-assisted approaches in primary repair of high anorectal malformations has been reported in neonates,¹⁻⁴ but the risk of sepsis without a covering colostomy remains a concern. Only 14 laparoscopically assisted attempts at primary repair have been reported.⁵⁻⁹ However, single-stage surgery remains attractive, particularly in resource-limited settings where multi-stage surgery represents a much higher financial burden,¹⁰ also considering the mixed social acceptability of colostomies¹¹ and their high associated morbidity (8 - 80%)¹²⁻¹⁷ as well as mortality.¹⁷ We present the case of a 2-day-old neonate who presented with oesophageal atresia without a distal tracheo-oesophageal fistula, and an anorectal malformation (ARM). The case provided the unique opportunity for a one-stage repair of the ARM without fear of faecal contamination of the peritoneal cavity, owing to the undistended bowel and sterile meconium, while the laparoscopic approach allowed excellent visualisation of the urethral fistula. Avoidance of a colostomy in this case also preserved the left colon for possible use in oesophageal replacement.

Case presentation

A male infant was born with a high imperforate anus and pure oesophageal atresia. A LAARP and a gastrostomy were performed on the second post-natal day. Primary oesophageal repair at 6 weeks failed, and the infant was discharged home to continue anal dilations. Stooling was regular with no constipation. Oesophageal substitution is planned after 10 months of age.

Surgical technique of neonatal LAARP

The rectum was identified and dissected down to the level of the proximal bulbo-urethral fistula. The fistula was transected as close as possible to the urethra. Perineal electrostimulation identified the centre of the anal sphincter, and an oval disc of skin was excised within this area. Visualising the midline of the puborectalis sling from above, we then dissected bluntly into the peritoneal cavity, after which a 5 mm port was inserted and used to pull the rectum through. A standard anoplasty was then fashioned and a urethral catheter left for 5 days.

Discussion

Concomitant ARMs are found in about 10 - 15% of patients with oesophageal atresia (OA),¹⁸⁻²¹ although the incidence may be as high as 23% in patients with OA without tracheo-oesophageal fistulae.¹⁹ OA without fistula represents about 7% of OA,²² and is therefore likely to be present in about 0.8% of ARM patients, based on a 12% incidence of OA occurring with ARM.²³

Primary anorectoplasty is routinely performed in low ARMs.^{24,25} Definitive repair of ARMs with recto-urethral fistulae in the newborn period is not recommended, owing to risk of injury to genito-urinary structures from 'blind' exploration without the anatomical information gained from a colostogram, and the increased risk of infection and wound dehiscence without faecal diversion.²⁴ Pena's posterior sagittal anorectoplasty (PSARP) is the gold standard in the management of ARMs with recto-urethral fistulae, replacing the Kiesewetter-Rehbein pull-through. In 2000, Georgeson introduced a

laparoscopically assisted approach.^{7,26} While the advantage of laparoscopy in bladder-neck and prostatic-urethral fistulae is clear,^{27,28} as laparotomy would otherwise be required, only a few small series report on laparoscopy in the management of other 'high/intermediate' malformations with recto-urethral fistulae.²⁹ One possible limitation of the laparoscopic approach is the imprecise angulation of the ano-rectal canal, which is easily positioned under direct vision in the open Pena dissection, although possibly at the expense of increased damage to nerves and muscle fibres. Various adjunctive techniques such as ultrasound guidance³⁰ and magnetic resonance imaging (MRI)³¹ have been used to improve this approach. Other concerns are inadequate excision of the recto-urethral fistula,³² with creation of a urethral diverticulum, although evidence for this actually causing problems is limited.³³ Comparative studies have shown equivalent outcomes between laparoscopically assisted pull-through and posterior sagittal anorectoplasty, although the number of patients and long-term follow-up reported is very limited at this stage,^{8,34-37} with only one reported randomised study.³⁸

One-stage neonatal pull-through

Georgeson⁷ and Vick *et al.*⁶ reported on 4 and 6 LAARPs in newborns respectively. However, experience with this approach has been very limited. The difficulties of stool contamination without a covering stoma make it a risky procedure, particularly in the intrinsically immune-compromised neonate.

The LAARP technique

Since the original Georgeson technique, various modifications have been applied, including methods to improve identification of the correct placement of the rectum within the levator and striated muscle complexes.^{39,40} We find levator response to nerve stimulation from the perineum can be readily identified from the peritoneal cavity, avoiding the need for a laparoscopic stimulator. It is safe and simpler to divide the fistula without suture ligation, which has been shown to be unnecessary,^{41,42} and follow-up micturating cysto-urethrograms in our series confirm this. Various techniques to determine the correct tract to the perineum through the muscle complex have been described. These include placement of a Veress needle followed by successively larger ports over the needle; balloon dilation of the tract, and use of Hegar dilators rail-roaded over the laparoscopic suction catheter.⁴² The importance of avoiding electrocautery close to the course of the pelvic nerve complexes has been emphasised, with a preference for sharp dissection.⁴²

Conclusion

The co-morbid oesophageal atresia without a distal tracheo-oesophageal fistula in this case of high ARM provided the opportunity to apply a laparoscopic approach, because of the absence of gut distension and the sterile gut contents. Laparoscopic surgery is still under-developed in resource constrained settings, but this case demonstrates its feasibility. While the problems associated with the management of the oesophageal atresia caused morbidity in this patient, these do not detract from the initial success of the LAARP, which held

major advantages for this patient because of the avoidance of colostomy and the preservation of the left colon for use in future oesophageal replacement. Long-term functional outcome will need to be assessed when the child is older.

REFERENCES

- Liu G, Yuan J, Geng J, Wang C, Li T. The treatment of high and intermediate anorectal malformations: one stage or three procedures? *J Pediatr Surg* 2004;39(10):1466-1471.
- Moore TC. Advantages of performing the sagittal anoplasty operation for imperforate anus at birth. *J Pediatr Surg* 1990;25(2):276-277.
- Mirshemirani A, Kouranloo J, Rouzrokh M, Sadeghiyan MN, Khaleghnejad A. Primary posterior sagittal anorectoplasty without colostomy in neonates with high imperforate anus. *Acta Med Iran* 2007;45(2):121-125.
- Khalil BA, Morabito A, Bianchi A. Transanoproctoplasty: a 21-year review. *J Pediatr Surg* 2010;45(9):1915-1919.
- Albanese CT, Jennings RW, Lopoo JB, Bratton BJ, Harrison MR. One-stage correction of high imperforate anus in the male neonate. *J Pediatr Surg* 1999;34(5):834-836.
- Vick LR, Gosche JR, Boulanger SC, Islam S. Primary laparoscopic repair of high imperforate anus in neonatal males. *J Pediatr Surg* 2007;42(11):1877-1881.
- Georgeson KE, Inge TH, Albanese CT. Laparoscopically assisted anorectal pull-through for high imperforate anus – a new technique. *J Pediatr Surg* 2000;35(6):927-930, discussion 30-1.
- Al-Hozaim O, Al-Maary J, AlQahtani A, Zamakhshary M. Laparoscopic-assisted anorectal pull-through for anorectal malformations: a systematic review and the need for standardization of outcome reporting. *J Pediatr Surg* 2010;45(7):1500-1504.
- Sydorak RM, Albanese CT. Laparoscopic repair of high imperforate anus. *Semin Pediatr Surg* 2002;11(4):217-225.
- Sidler D, Millar AJ, Rode H, Brown RA, Franck M, Cywes S. Neonatal Soave endorectal pull-through for Hirschsprung's disease. *S Afr J Surg* 1999;37(2):47-49, discussion 49-50.
- Ameh EA, Mshelbwala PM, Sabiu L, Chirdan LB. Colostomy in children--an evaluation of acceptance among mothers and caregivers in a developing country. *S Afr J Surg* 2006;44(4):138-139.
- Pena A, Migotto-Krieger M, Levitt MA. Colostomy in anorectal malformations: a procedure with serious but preventable complications. *J Pediatr Surg* 2006;41(4):748-756, discussion 756.
- Bischoff A, Levitt MA, Lawal TA, Pena A. Colostomy closure: how to avoid complications. *Pediatr Surg Int* 2010; Aug 17.
- Chirdan LB, Uba FA, Ameh EA, Mshelbwala PM. Colostomy for high anorectal malformation: an evaluation of morbidity and mortality in a developing country. *Pediatr Surg Int* 2008;24(4):407-410.
- Ekenze SO, Agugua-Obianyo NE, Amah CC. Colostomy for large bowel anomalies in children: a case controlled study. *Int J Surg* 2007;5(4):273-277.
- Sheikh MA, Akhtar J, Ahmed S. Complications / problems of colostomy in infants and children. *J Coll Physicians Surg Pak* 2006;16(8):509-513.
- Cigdem MK, Onen A, Duran H, Ozturk H, Otcu S. The mechanical complications of colostomy in infants and children: analysis of 473 cases of a single center. *Pediatr Surg Int* 2006;22(8):671-676.
- Moore SW, Alexander A, Sidler D, et al. The spectrum of anorectal malformations in Africa. *Pediatr Surg Int* 2008;24(6):677-683.
- Chittmitrapap S, Spitz L, Kiely EM, Brereton RJ. Oesophageal atresia and associated anomalies. *Arch Dis Child* 1989;64(3):364-368.
- Cho S, Moore SP, Fangman T. One hundred three consecutive patients with anorectal malformations and their associated anomalies. *Arch Pediatr Adolesc Med* 2001;155(5):587-591.
- Engum SA, Grosfeld JL, West KW, Rescorla FJ, Scherer LR 3rd. Analysis of morbidity and mortality in 227 cases of esophageal atresia and/or tracheoesophageal fistula over two decades. *Arch Surg* 1995;130(5):502-508; discussion 508-509.
- Spitz L. Oesophageal atresia. *Orphanet J Rare Dis* 2007;2:24.
- Piekarski DH, Stephens FD. The association and embryogenesis of tracheo-oesophageal and anorectal anomalies. *Prog Pediatr Surg* 1976;9:63-76.
- Levitt MA, Pena A. Anorectal malformations. *Orphanet J Rare Dis* 2007;2:33.
- Menon P, Rao KL. Primary anorectoplasty in females with common anorectal malformations without colostomy. *J Pediatr Surg* 2007;42(6):1103-1106.
- Georgeson K. Laparoscopic-assisted anorectal pull-through. *Semin Pediatr Surg* 2007;16(4):266-299.
- Lopez M, Kalfa N, Allal H, et al. Anorectal malformation (ARM) with bladder fistula: Advantages of a laparoscopic approach. *Eur J Pediatr Surg* 2009 Jun 10.
- Hay SA. Transperineal rectovesical fistula ligation in laparoscopic-assisted abdominoperineal pull-through for high anorectal malformations. *J Laparoendosc Adv Surg Tech A* 2009;19 Suppl 1:S77-79.
- Lima M, Antonellini C, Ruggeri G, Libri M, Gargano T, Mondardini MC. Laparoscopic surgical treatment of anorectal malformations. *Pediatr Med Chir* 2006;28(4-6):79-82.

30. Kubota A, Kawahara H, Okuyama H, et al. Laparoscopically assisted anorectoplasty using perineal ultrasonographic guide: a preliminary report. *J Pediatr Surg* 2005;40(10):1535-1538.
31. Raschbaum GR, Bleacher JC, Grattan-Smith JD, Jones RA. Magnetic resonance imaging-guided laparoscopic-assisted anorectoplasty for imperforate anus. *J Pediatr Surg* 2010;45(1):220-223.
32. Koga H, Kato Y, Shimotakahara A, et al. Intraoperative measurement of rectourethral fistula: prevention of incomplete excision in male patients with high-/intermediate-type imperforate anus. *J Pediatr Surg* 2010;45(2):397-400.
33. Lopez PJ, Guelfand M, Angel L, et al. Urethral diverticulum after laparoscopically-assisted anorectal pull-through (LAARP) for anorectal malformation: is resection of the diverticulum always necessary? *Arch Esp Urol* 2010;63(4):297-301.
34. Ichijo C, Kaneyama K, Hayashi Y, et al. Midterm postoperative clinicoradiologic analysis of surgery for high/intermediate-type imperforate anus: prospective comparative study between laparoscopy-assisted and posterior sagittal anorectoplasty. *J Pediatr Surg* 2008;43(1):158-162; discussion 162-163.
35. Kudou S, Iwanaka T, Kawashima H, et al. Midterm follow-up study of high-type imperforate anus after laparoscopically assisted anorectoplasty. *J Pediatr Surg* 2005;40(12):1923-1926.
36. El-Debeiky MS, Safan HA, Shafei IA, Kader HA, Hay SA. Long-term functional evaluation of fecal continence after laparoscopic-assisted pull-through for high anorectal malformations. *J Laparoendosc Adv Surg Tech A* 2009;19 Suppl 1:S51-54.
37. Lin CL, Wong KK, Lan LC, Chen CC, Tam PK. Earlier appearance and higher incidence of the rectoanal relaxation reflex in patients with imperforate anus repaired with laparoscopically assisted anorectoplasty. *Surg Endosc* 2003;17(10):1646-1649.
38. Yang J, Zhang W, Feng J, et al. Comparison of clinical outcomes and anorectal manometry in patients with congenital anorectal malformations treated with posterior sagittal anorectoplasty and laparoscopically assisted anorectal pull through. *J Pediatr Surg* 2009;44(12):2380-2383.
39. Iwanaka T, Arai M, Kawashima H, et al. Findings of pelvic musculature and efficacy of laparoscopic muscle stimulator in laparoscopy-assisted anorectal pull-through for high imperforate anus. *Surg Endosc* 2003;17(2):278-281.
40. Yamataka A, Segawa O, Yoshida R, Kobayashi H, Kameoka S, Miyano T. Laparoscopic muscle electrostimulation during laparoscopy-assisted anorectal pull-through for high imperforate anus. *J Pediatr Surg* 2001;36(11):1659-1661.
41. Rollins MD, Downey EC, Meyers RL, Scaife ER. Division of the fistula in laparoscopic-assisted repair of anorectal malformations – are clips or ties necessary? *J Pediatr Surg* 2009;44(1):298-301.
42. Bhandary KS, Kumaran V, Rajamani G, et al. Laparoscopic assisted anorectal pull through: Reformed techniques. *J Indian Assoc Pediatr Surg* 2009;14(4):210-214.

SAMF

South African Medicines Formulary



Ninth Edition

The ESSENTIAL REFERENCE for every healthcare professional!

The carefully and thoroughly updated 9th edition of the South African Medicines Formulary (SAMF) can now be ordered. It is your essential reference to rational, safe and cost-efficient use of medicines. That is why you should not prescribe without it.

The newly published SAMF provides easy access to the latest, most scientifically accurate information – including full drug profiles, clinical notes and special prescriber's points. The convenient pocket-size design enables you to fit it comfortably into your bag or hospital coat pocket – always at hand for ready reference.

WHY YOU SHOULDN'T BE WITHOUT THE SAMF 9TH EDITION

The new 9th edition of SAMF provides expanded information on key issues facing South African healthcare professionals today, including antiretrovirals, TB treatment guidelines, management guidelines for asthma and chronic heart failure, other common chronic conditions and prescribing in sport.

- It presents practical, new approaches to the management of venomous bites and stings.
- It outlines extensively the acute adverse reactions to drugs of abuse, and their management.
- It features new as well as existing drugs, indexed by both trade and generic names.
- It offers fresh insights into informed prescribing and carries cautionary guidelines on drug interactions and a range of special risk patients and conditions.

And, as always, you can rely on...

- the professional compilation and editing by a team from the Division of Clinical Pharmacology, UCT
- an independent and unbiased guide on prescribing in South Africa today
- the indication of agents included in the SA and WHO essential drug lists
- support of the SA national drug policy
- guidance for prescribing during pregnancy and lactation, and in patients with porphyria, liver disease and renal impairment (including tables with drug dosage adjustments); and
- indexed and page tabs for quick and easy access to each section.

YOUR SATISFACTION IS GUARANTEED

3 easy order options:

1. PHONE EDWARD OR BYRON – 021 6817000
2. FAX the completed SAMF order form to 0866006218
3. EMAIL: edwardm@hmpg.co.za OR byronm@hmpg.co.za

