

# AN ASSESSMENT OF GROWTH IN HIGH AND LOW SOCIO- ECONOMIC STATUS SCHOOLCHILDREN IN SOUTH AFRICA

G J Louw, S M Naidoo

'We are guilty of many errors and many faults, but our worst crime is abandoning the children, neglecting the foundation of life. Many of the things we need can wait. The child cannot. Right now is the time his bones are being formed, his blood is being made and his senses are being developed. To him we cannot answer "Tomorrow". His name is "Today".'

(Gabriela Mistral, 1948, World Health Organisation, 1997)

The patterns of physical growth (height, weight, length of trunk and limbs, circumference of trunk and limbs, and limb breadths) and function (grip strength of both hands and neuromuscular reaction time) of Cape Coloured (specifically mixed origin) schoolchildren from urban and rural areas and contrasting socio-economic status (SES) levels, were measured. The mixed longitudinal study accumulated data over more than a decade, and included 929 male and 1 160 female pupils of high SES (HSES) in the Cape Town urban area, aged 5 - 20 years, and 954 male and 1 030 female pupils of low SES (LSES) in rural areas of the Little Karoo, aged 5 - 19 years. Means for every anthropometric character were calculated and matched against age for each of the four groups, for comparative purposes. Standard deviations were recorded for each character. Figs 1 - 16 show the means for each of the anthropometric characters, matched with age groups, for HSES and LSES boys and girls.

The results reflect the importance of positive intervention in the growth and development of the LSES children at different periods for the girls and boys, namely pre-pubertally in girls, particularly during the period of possible pre-pubertal growth spurt (8 - 10 years of age), and continuously and consistently for the boys, from 5 to 19 years and onwards. Intervention should take the form of improved diet, increased exercise, and plenty of time spent outdoors in the sun.

*Department of Human Biology, University of Cape Town*

G J Louw, DVSc

S M Naidoo, BSc




Fig. 1. Means of total body height (mm) for age (yrs).




Fig. 4. Means of upper limb length (mm) for age (yrs).




Fig. 2. Means of body weight (kg) for age (yrs).




Fig. 5. Means of lower limb length (mm) for age (yrs).




Fig. 3. Means of trunk length (mm) for age (yrs).




Fig. 6. Means of arm circumference (mm) for age (yrs).




Fig. 7. Means of bi-acromial diameter (mm) for age (yrs).




Fig. 10. Means of bi-epicondylar breadth of femur (mm) for age (yrs).




Fig. 8. Means of bi-iliacristal diameter for age (yrs).




Fig. 11. Means of chest circumference (mm) for age (yrs).




Fig. 9. Means of bi-epicondylar breadth of humerus (mm) for age (yrs).




Fig. 12. Means of subscapular skinfold thickness (mm) for age (yrs).




Fig. 13. Means of abdominal skinfold thickness (mm) for age (yrs).




Fig. 15. Means of left-hand grip strength (units) for age (yrs).




Fig. 14. Means of right-hand grip strength (units) for age (yrs).




Fig. 16. Means of neuromuscular reaction time (ms) for age (yrs).

Four patterns of growth and development emerge from this analysis, and the anthropometric characters falling into each pattern are grouped accordingly.

The majority of characters fell into the first pattern, namely total body height, body weight, trunk length, upper extremity length, lower extremity length, arm circumference, bi-acromial diameter, bi-iliocristal diameter, bi-epicondylar width of humerus, bi-epicondylar width of femur, and chest circumference. The rates of increase of the anthropometric characters, as reflected by the slopes of the graphs, are similar for HSES and LSES males and females from age 5 until puberty, where the values for the girls drop off rapidly as the growth plates of their bones close. The different values observed along the graph are a reflection of differing starting points at age 5 for the HSES and LSES groups, and there is no 'catch-up'

achieved at any time. The males have higher final values because they continue growing postpubertally, into their twenties. For all four groups, only a small pre-pubertal growth spurt is visible.

The second pattern of growth was similar for the various skinfold thicknesses. The skinfold thicknesses, as a reflection of 'fatness', fluctuate continuously, but a pre-pubertal peak is visible in all groups. Although both HSES and LSES groups commence at similar values at age 5, the HSES group rises rapidly, with females exceeding the males at all ages. Both HSES and LSES males show lower values after puberty since their linear characters all continue to increase, and the LSES females overtake the HSES males at the time of puberty. A girl's 'fatness' is an important trigger for menarche.

The third pattern of development was for left and right grip

strengths, which are virtually identical. The pattern for boys is markedly different from that of the girls in that there is a dramatic increase in the strength just before puberty right through to the mid-teens, after which the graphs plateau. The values for the girls increase at the same rate as those of the boys, as reflected by similar slopes, but the girls reach their maximum strength at a younger age, shortly after puberty.

The last pattern of development was shown by neuromuscular reaction time. The graphs for all four groups of schoolchildren are similar in shape from 6 to 16 years in that there is a declining reaction time from 6 to 8 years, then a slow increase from 8 to 12 years, after which the time decreases again until 16 years of age. The graphs demonstrate that HSES males have the quickest reaction times, followed by the LSES males. The girls of both groups show considerable fluctuation, but the LSES girls have the slowest reaction times of all groups at all ages, and do not achieve a 'catch-up' period.

The authors wish to thank the staff and students of this Department for their assistance in collecting the data over the years, and the staff and pupils of the schools visited for their wonderful co-operation.

---