

UNUSUAL BEHAVIOUR OF FROG AND HUMAN SERUM PROTEINS LOADED WITH ^{131}I -LABELLED THYROID HORMONES DURING PAPER ELECTROPHORESIS

A. VAN ZYL and B. WILSON

Department of Physiology, Medical School, University of Cape Town

It has been reported earlier that the serum protein-bound iodine in the South African toad (*Xenopus laevis*) is higher than that of most animals.¹ Since it is thought that the serum proteins to which thyroid hormones are bound, determine to some extent the speed with which the hormones enter the cells,² a greater affinity of the specific binding proteins for thyroid hormones or a slower turnover rate of the binding proteins in cold-blooded animals could explain in part the relatively high serum protein-bound iodine and, therefore, the decreased metabolic rate in the frog.

During studies on the binding properties of serum proteins with ^{131}I -labelled thyroid hormones, the association of ^{131}I -labelled tri-iodothyronine ($^{131}\text{T}_3$) and thyroxine ($^{131}\text{T}_4$) with the proteins of frog serum was compared with that of human serum. The specific binding proteins, in most cases, were overloaded with $^{131}\text{T}_3$ and $^{131}\text{T}_4$. The hormones were labelled with ^{131}I chemically by exchange³ and also endogenously by injecting $200\mu\text{c}$ of ^{131}I into a mouse. After 48 hours the thyroid was digested enzymically. The hydrolysate was chromatographed, radio-autographed, and the radioactive bands corresponding to tri-iodothyronine and thyroxine markers were eluted (Fig. 1*). The ^{131}I -labelled thyroid hormones were added *in vitro* to serum and electrophoresed on paper in barbiturate-acetate buffer at pH 8.6.

The electrophoresed frog serum, unlike human serum, gave only 3 distinct bands on the anode side of the origin (Fig. 2). The albumin band of frog serum shows a greater electrophoretic mobility than that of human serum. Mid-way between the albumin and the origin, a broad band appeared in frog serum which corresponded to the area between the α_1 and the α_2 globulin of human serum. This protein band is referred to as the β -globulin, although it may not necessarily be the same as β -globulin of human serum. In some cases the γ -globulin is fairly well defined, but in most cases it is just visible when $20\mu\text{l}$. of frog serum is used.

The radioactivity from endogenously prepared $^{131}\text{T}_4$ added to serum was associated with both the inter-alpha globulins and albumin when human serum was used. On the other hand, in frog serum it was mainly associated with β -globulin and to a lesser extent with albumin (Fig. 3).

In order to gain some information about the binding capacity of frog and human serum proteins, chemically prepared $^{131}\text{T}_4$ was added in known concentrations to serum, and electrophoresed. At a concentration of $0.05\mu\text{g}$. $^{131}\text{T}_4$ /ml. of serum the bulk of radioactivity was again associated with the inter α -globulins in human serum and with the β -globulins in frog serum (Fig. 4). However, when the chemically prepared $^{131}\text{T}_4$ was increased to $0.5\mu\text{g}$. $^{131}\text{T}_4$ /ml. of serum, most of the radioactivity shifted onto the human serum albumin, whereas the albumin of frog

serum had not taken up much $^{131}\text{T}_4$. In frog serum such high concentrations of $^{131}\text{T}_4$ were mainly associated with β -globulin and another radioactive peak appeared ahead of the albumin (Fig. 5).

When endogenously prepared $^{131}\text{T}_3$ was added to frog and human sera, an unusual electrophoretic migration of the radioactivity was noted. With human serum the major portion of the radioactivity moved during electrophoresis for a distance of about 5 cm. from the origin towards the cathode, while smaller portions were associated mainly with albumin and pre-albumin. With frog serum the major portion of the radioactivity was similarly located as an intense band about 5 cm. on the cathode side of the origin whereas the rest of the activity coincided with the positions of albumin and β -globulin (Fig. 6).

Staining of the electrophoretograms showed no protein bands corresponding to the ^{131}I -labelled substance which migrated towards the cathode. Electrophoresis of the same sera loaded with endogenously prepared $^{131}\text{T}_3$ was repeated and the same results were obtained.

In an attempt to identify the radioactive substance which migrated towards the cathode, the human and frog sera containing the endogenously prepared $^{131}\text{T}_3$ and $^{131}\text{T}_4$ were chromatographed one-dimensionally in butanol:dioxan:2-N NH₄OH (4:1:5) with carriers T_3 and T_4 (Fig. 7). The darkened areas on the radio-autograph of the chromatogram indicated that the $^{131}\text{T}_4$ corresponded exactly with carrier T_4 in frog and human sera, but that the $^{131}\text{T}_3$ was slightly ahead of the carrier T_3 spots. The concentration of carrier T_3 was greater than that of carrier T_4 , and this may have caused the $^{131}\text{T}_3$ to move slightly ahead of carrier T_3 .

The section of the electrophoretogram corresponding to the ^{131}I -labelled substance, which migrated towards the cathode, was cut out, eluted, and the extract chromatographed two-dimensionally in butanol:dioxan:2-N NH₄OH and in butanol:acetic acid:water (120:30:50). Most of the activity corresponded to non-radioactive T_3 and iodide (Fig. 8). Other carriers tested out, like 3-mono-iodothyronine, 3:3'-di-iodothyronine, and 3:5:3'-tri-iodothyroacetic acid, did not coincide in two-dimensional chromatograms with the radioactive spots.

It is concluded that (1) frog serum, unlike human serum, carries thyroxine on the β -globulins, (2) the binding capacity of the β -globulins of frog serum for thyroid hormones is greater than that of the inter- α -globulins of human serum, and (3) the bulk of the ^{131}I -labelled substance, which migrated towards the cathode during electrophoresis of frog and human sera under the conditions of the experiment, was in fact $^{131}\text{T}_3$.

REFERENCES

1. van Zyl, A. and Kerrieh, J. E. (1955): *S. Afr. J. Med. Sci.*, **20**, 9.
2. Solomon, D. H. and Dowling, J. T. (1960): *Ann. Rev. Physiol.*, **22**, 615.
3. Taurog, A. and Chaikoff, I. L. (1957): *Methods in Enzymology*, **IV**, p. 860, New York: Academic Press Inc.

* Figs. 1-8 are on p. 1047.




Fig. 1. Characterization of the ¹³¹I-labelled amino acids in a mouse thyroid chromatographed in butanol:dioxan:2-N NH₄OH. The dotted areas indicate the positions of the inactive carriers di-iodothyrosine (DIT), mono-iodothyrosine (MIT), iodide (I), thyroxine (T₄) and tri-iodothyronine (T₃).

Fig. 2. Stained electrophoretic patterns of (A) normal human serum, and (B) frog serum.

Fig. 3. The association of endogenously prepared ¹³¹I-T₄ with frog (F+T₄) and human (H+T₄) sera. The radio-autograms are shown to the right of the stained electrophoretograms in each case.

Fig. 4. The association of 0.05 μg ¹³¹I-T₄/ml. with (A) human serum, and (B) frog serum. The histograms represent the radioactive counts per cm. length of the electrophoretograms.

Fig. 5. The association of 0.5 μg T₄/ml. with (A) human serum, and (B) frog serum. The radio-autogram (above) and the graph indicate qualitatively and quantitatively the activity associated with the various protein fractions on the electrophoretogram below the graph.

Fig. 6. The association of endogenously prepared ¹³¹I-T₃ with human (H) and frog (F) serum proteins. The stained electrophoretogram is indicated on the left of each pair while the radio-autogram is on the right.

Fig. 7. Radio-autogram of chromatographic analyses of frog (A and B) and human (C and D) sera to which endogenously prepared ¹³¹I-T₄ and ¹³¹I-T₃ had been added.

Fig. 8. Characterization of the endogenously prepared radioactive substance which migrated to the cathode on electrophoresis, indicating the positions of non-radioactive carriers: iodide (I), tri-iodothyronine (T₃) and tri-iodothyroacetic acid (TRIAC). Solvents: 1. Butanol:dioxan:2-N NH₄OH (BDA) 2. Butanol:acetic acid:water (BAW).