

Short communication

NUTRIENT DYNAMICS OF IMO RIVER ESTUARY, SOUTH EASTERN NIGERIA

Osondu Christopher Akoma

Department of Botany, University of Benin, Benin-City, Nigeria. Email:osonduakoma@yahoo.com

ABSTRACT: The physical and chemical properties of the Imo River estuary, Southeastern Nigeria were investigated between April 2001 and June 2002. Variations in water temperature, total dissolved solids, conductivity, dissolved oxygen, nitrate, sulphate, phosphate and silicate are reported. With the exception of dissolved oxygen all parameters were significantly different ($P < 0.05$) in the study stations. Conductivity was positively correlated with total dissolved solids. The same holds true for phosphate and nitrate. The parameters showed both spatial and temporal variations. Temperature, TDS, conductivity, were high during the dry season while nitrate, sulphate, phosphate and silica recorded high wet season values. Flood influx into the estuary during the wet season and tidal inflow of seawater during the dry season were the two major factors that influenced the nutrient dynamics of the Imo River estuary.

Key words/phrases: Estuary, Imo River, Nigeria, nutrients, variations

INTRODUCTION

It is common knowledge that estuaries are exposed to continuous eutrophication through daily enrichment from natural systems and human wastes. Specifically, nutrient supplies to estuaries are usually coming from the sea, inflowing rivers, adjacent marshes and vegetation and from anthropogenic sources. This leads to nutrient enrichment of water and with adequate solar radiation gives rise to hydrobiological changes which may result in bloom formations (Opute, 1990). The shallow and well-mixed waters of the estuaries are often ideal sites for high rates of photosynthesis and secondary production (Pearl, 1988).

Reports on the physical and chemical characteristics of Nigerian rivers include those of Egborge (1971) on River Oshun, Adebisi (1981) on Ogun River, Opute (1990) on the Warri/Forcados estuary, Erondu and Chindah (1991) on the New Calabar River. Others include Chikere and Okpokwasili (2001) on Eleme River, Nwankwo and Jaiyeola (2001) on Lagos lagoon, Kadiri (2002) on the eastern Niger Delta and Akpan *et al.* (2002) on the Great Kwa River. There is scanty information on the limnology of Imo River despite the fact that it is one of the major rivers in south eastern Nigeria and its estuary part of the anastomosing network of creeks in the Niger Delta. The present study was undertaken to provide baseline data on the physical and chemical conditions in the

estuary, investigate spatial and temporal variations and sources of nutrient input into the estuary.

MATERIALS AND METHODS

Water samples were collected for fifteen months (April 2001 to June 2002) from locations along 45 km stretch to the mouth of the river discharge into the Atlantic Ocean (Fig. 1). Station 1 was considered as the headwaters of the estuary and characterized by dense tropical rain forest swamp fringing vegetation with species of *Alstonia boonei* De Wild; *Raffia hookeri* Mann *et al.*, Wendl., *Emilia sonchifolia* (L.) DC, *Spondianthus preusii* and *Mitragyna ciliata*. Stations 2 and 3 were freshwater stations with marked diurnal tidal amplitude and fringing vegetation mainly mangrove swamp and dense vegetation of *Nypa fruticans*. Stations 4, 5 and 6 were characterized by brackish water, semi-diurnal tides, coastal alluvium and sandy beaches. Generally, the study area falls within the tropical rainforest zone of Nigeria characterized by rainy season (March to November) and dry season (December to February) with two rainfall peaks in July and September (Iloeje, 1991). Other activities include marine transport, fishing and lumbering.

Water temperature was measured *in situ* with mercury in glass thermometer. Dissolved oxygen was fixed on site by adopting the alkali-azide modification of Winkler's method. Total dissolved solids, conductivity, colour and turbidity were

recorded with a HACH DR2000 spectrophotometer. Nutrient analyses for nitrate, sulphate, phosphate and silica were performed as outlined by APHA (1998). The samples were transported in an ice box and later stored in a deep freezer pending analyses due to the difficulty in logistics between the sampling points and laboratory facility. Another major limitation in this study is the lack of filtration

of the water samples before analyses and this had a great impact on the result considering the effect of suspended matter in the water samples.

Mean and standard error for each parameter were calculated for the various sampling stations, while one way analysis of variance was computed to know the level of significant difference among the stations.




Fig. 1. Map of Imo River estuary showing sampling stations.

RESULTS AND DISCUSSION

Table 1 shows range, mean and standard error values of physico-chemical parameters obtained from the six sampling stations in the Imo River estuary. The surface water temperatures varied from 25°C to 30°C. Temperature variations were due to changes in solar radiation, cloud cover and time of sampling. Temperature variations in tropical water bodies are of minimal effect and also vary according to the seasons (Talling, 1986). The difference in sampling time as a result of the distance between the first and last stations could be responsible for the observed trend in temperature variation.

Total dissolved solids ranged from minimum value of 4.2-168mg^l⁻¹ at station 1 to maximum range of 516-19890mg^l⁻¹ at station 5. There was progressive increase in the total dissolved solids concentration downstream and this is attributable to rainy season run off from tributary streams and the fact that the lower reaches of the estuary is interconnected with the Niger Delta creek system with its attendant seasonal flood influx and tidal effect of the ocean. Conductivity values ranged from 15.3µScm⁻¹ recorded at station 2 to 39780µScm⁻¹ at station 5. This range is comparable with 12.0 to 46,000µScm⁻¹ recorded in Bonny River – a brackish water river in the Niger Delta by Chindah and Pudo (1991). The Imo river estuary conductivity conforms widely with the general range of conductivity of the brackish waters of the Niger Delta and other Nigerian coastal waters (Nwankwo and Jaiyeola, 2001; Kadir, 2002).

Dissolved oxygen content ranged from 4.6mg^l⁻¹ recorded in station 1 to 10.2mg^l⁻¹ in station 6. However lower values have been reported for various rivers in Nigeria. Examples include 3.5 to 7.2mg^l⁻¹ for Bonny river (Chindah and Pudo, 1991),

3.0-6.0mg^l⁻¹ for the Lagos lagoon (Nwankwo, 1994), 3.0-4.3mg^l⁻¹ for the Warri/Forcados estuary (Opote, 1990) and 4.0-8.33mg^l⁻¹ for Great Kwa river, Akpan, Offem and Nya (2002). Dissolved oxygen concentrations recorded in this study could be due to photosynthetic activity of phytoplankton population especially at stations 5 and 6. Flocculation of colloidal suspension and the attendant increase in transparency could be responsible for increased primary productivity and oxygen concentration. (Reynolds and Decsy, 1996).

The nitrate, sulphate, phosphate and silica are some of the constituents of the nutrient composition in any body of water and their range, mean and standard error are shown in also shown in Table.1. In the present study these nutrients increased downstream in the estuary except silica which showed a reverse trend of decreasing values. Generally the nutrients variations were subject to seasonal changes due to flood influx from land drainage and tributary stream during the rainy season. Table 2 shows correlation coefficient of the various parameters investigated in the present study and reveals positive correlation between conductivity and total dissolved solids and also between nitrate and phosphate (the major limiting nutrients for phytoplankton growth and primary productivity). Tidal action could also lead to resuspension of nutrients from the sediments. Inundation of the fringing mangrove vegetation at high tides yield decaying plant and animal materials into the estuary and these are potential sources of nutrient enrichment (John, 1986). It is important to note the effect of suspended matter on the result owing to non-filtration of the samples. This no doubt could have contributed to the high nutrient concentrations recorded in this study.

Table 1. Summary of physico-chemical properties of Imo River estuary.

Parameters	Station 1	Station 2	Station 3	Station 4	Station 5	Station 6	P value
Temperature (°C)	26-29 27.33±0.23	25-29 27.4±0.29	26-30 27.75±0.30	26-30 28.2±0.30	25-30 28.6±0.33	28-30 29.0±0.26	P<0.05
TDS (mg ^l ⁻¹)	4.3-168 26.57±10.33	7.6-483 76.54±34.35	8.77-4136 904.55± 435.07	232-8240 4121.44±674.37	516-19890 6215.33±1239.42	5250-15529.5 7469.90±1387.02	P<0.001
Conductivity (µScm ⁻¹)	18.5-242 47.79±14.44	15.3-540 128.87±46.50	17.45-8273 1809.74± 870.23	463-16250 8233.89±1335.56	1038-39780 1452.73±2258.73	10896-31059 21523±2092.37	P<0.001
DO (mg ^l ⁻¹)	4.6-8.0 6.84±0.24	5.0-9.1 7.16±0.30	6.0-8.6 7.45±0.27	5.8-8.8 7.43±0.24	5.9-9.6 7.86±0.30	5.2-10.2 7.27±0.48	P>0.05
Nitrate (mg ^l ⁻¹)	0.01-0.84 0.29±0.06	0.04-0.88 0.38±0.07	0.01-0.31 0.19±0.02	0.04-1.27 0.50±0.10	0.95-4.32 2.28±0.42	1.09-4.3 2.39±0.36	P<0.001
Sulphate (mg ^l ⁻¹)	0.08-0.98 0.38±0.07	0.09-0.9 0.32±0.06	0.29-3.2 1.91±0.26	0.09-9.02 3.86±0.79	0.18-10.88 5.82±0.86	7.20-18.50 10.07±1.02	P<0.001
Phosphate (mg ^l ⁻¹)	0.01-0.25 0.10±0.01	0.03-0.75 0.19±0.05	0.1-1.44 0.52±0.23	0.01-0.48 0.23±0.03	0.01-2.01 0.59±0.17	0.15-1.61 0.74±0.16	P<0.05
Silicate (mg ^l ⁻¹)	3.0-27.0 15.16±2.36	3.2-20.0 12.71±1.66	2.0-14.44 6.23±1.14	1.45-6.50 3.61±0.37	0.38-1.02 0.65±0.05	0.25-1.05 0.61±0.10	P<0.001

Table 2. Correlation coefficient of physico-chemical parameters in the Imo River estuary.

	TEMP	TDS	COND	ALK	DO	NITRATE	SULPH	PHOSPH	SILICATE
TEMP									
TDS	.357								
CONDUCTIVITY	.096	.870**							
DO	-.389	-.380	-.514*	-.237					
NITRATE	.076	.081	.025	.197					
SULPHATE	-.445*	-.252	-.017	.306	.031				
PHOSPHATE	.048	.182	.476	.133	.457				
SILICATE	.035	.101	.126	.330	-.148	.423			
	.451	.361	.327	.115	.299	.058			
	-.353	-.047	.106	.384	-.232	.811**	.404		
	.098	.435	.353	.079	.203	.000	.068		
	-.588*	-.456*	-.398	-.207	.428	.116	-.651**	.152	
	.011	.044	.071	.229	.056	.340	.004	.295	

* Correlation is significant at the 0.05 level (1-tailed), ** Correlation is significant at the 0.01 level (1-tailed).

CONCLUSION

Unlike other estuarine systems in the Niger Delta, the Imo river estuary is devoid of serious oil exploratory activity and its attendant pollution. As a result of reduced anthropogenic impact allochthonous input of nutrients from wet season run off, decaying organic matter from the fringing swampy vegetation and flocculation at the freshwater and brackish water interface coupled with biological activities are responsible for high nutrient concentrations. The present study was a preliminary study to provide baseline information on the physico-chemical conditions in the estuary and proffer suggestions about possible sources of nutrient enrichment. A more detailed and articulated program could be carried out in the future to correct the lapses in the present study.

REFERENCES

- Adebisi, A.A. (1981). The physico-chemical hydrology of a tropical seasonal river, upper Ogun River. *Hydrobiologia* 79:157-165.
- Akpan, E.R., Offem, J.O. and Nya, A.E. (2002). Baseline ecological studies of the Great Kwa River, Nigeria I: Physico-chemical studies. *Afr. J. Env. Poll. & Health* 1(1):83-90.
- APHA (1998). *Standard Methods for the Examination of Water and Wastewater*, 20th edition. American Water Works Association, Water Environment Federation, Washington D.C. 1287 pp.
- Chikere, B.O. and Okpokwasili, G.C. (2001). Organic pollution in a Niger Delta River receiving petrochemical effluents. *Trop. Freshw. Biol.* 10:19-33.
- Chindah, A.C and Pudo, J.K. (1991). A preliminary check of algae found in plankton of Bonny River in Niger Delta, Nigeria. *Fragm. Flor. Geobot* 36:177-126.
- Egborge, A.B.M. (1971). The chemical hydrology of the River Oshun, Western State, Nigeria. *Freshwater Biol.* 1:257-272.
- Erondu, E.S. and Chindah, A.C. (1991). Physico-chemical and phytoplankton changes in a tidal freshwater station of the New Calabar River, South Eastern, Nigeria. *Environment and Ecology* 9(3):561-570.
- Iloje, N.P. (1991). *New Geography of Nigeria*. Longman, Lagos-Ibadan, 201 pp.
- John, D.M. (1986). The inland waters of tropical West Africa. *Archivum Hydrobiologia* 23:1-244.
- Kadiri, M.O. (2002). A spectrum of phytoplankton flora along salinity gradient in the eastern Niger Delta area of Nigeria. *Acta Bot. Hungarica* 44(1-2):75-83.
- Nwankwo, D.I. (1994). Hydrochemical properties and bottom dwelling diatoms of a Lagos Lagoon sewage disposal site. *Pollskje Arch. Hydrobiol.* 41:35-47.
- Nwankwo, D.I. and Jaiyeola, M.A. (2001). Surf diatoms of Lagos lagoon beach, Nigeria. *J. Aquat. Sci.* 16:35-38.
- Opute, F.I. (1990). Phytoplankton flora of the Warri/-Forcados estuary of southern Nigeria. *Hydrobiologia* 208:101-109.
- Pearl, H.W. (1988). Nuisance algae blooms of coastal estuaries and inland waters. *Limnol. Oceanogr.* 33:823-847.
- Reynolds, C.S. and Descy, J.P. (1996). The production, biomass and structure of phytoplankton and large rivers. *Arch. Hydrobiol. Suppl.* 113:161-187.
- Talling, J.F. (1986). Seasonality of phytoplankton in African Lakes. *Hydrobiologia* 138:139-160.