

A Look into Colostomy in Pediatric Patients Presented To Al Ribat University Hospital December 2013-December 2014

Majed A A Mohammad¹ and Omer Al Amin²

Abstract

Background: Colostomy is a very common procedure in pediatric pts. It has a lot of indications and complications.

Objectives: 1) To observe the frequency of indications complications of colostomies between patients done in Al- Ribat university hospital and other patients done in other hospitals who presented there for definitive procedure. 2) To determine the final outcome of colostomy in pediatric patients.

Materials and Methods: Observational prospective study, performed in patients presented to Al Ribat university hospital including patients admitted for colostomy or for final treatment in the period between December 2013 and December 2014. The stoma related complications were tested using Chi-square. The level of significance was taken as $p < 0.05$.

Results: There were 50 patients with the peak age group of 4 years (range: 10 days to 8 years). The male percentage was 70% and female 30%. The indications of colostomy were HSD 29 (58 %), anorectal malformations 19 (38%), rectal injury 1 (2%) and anal malignancy 1 (2%). Complications occurred in 24 patients (48%) of the studied sample. The commonest complication was prolapse 7(29, 2%) followed by stenosis 5(20.8%). Complications were higher in patients operated as an elective cases in compare to emergency operations (51.8% vs. 43.5), and patients operated by registrars in comparison to other surgical operating levels, and in transverse colostomy in comparison to sigmoid colostomy(57,1% vs 46,5%), and in loop colostomy in comparison to divided colostomy (51,2% vs 33,3%), but all these results were statistically not significant. One mortality case was reported during this study (2%).

Conclusion: colostomy being a common procedure in pediatrics with high rates of serious complications deserves special attention for a better outcome.

Key words: colostomy, complications, Hirschsprungs disease, anorectal malformations.

Colostomy is a very common part in the management of wide range of congenital and acquired gastrointestinal conditions in pediatrics. HSD and anorectal malformations are amongst the major indications of colostomy in neonates and infants and children. It is done basically for diversion of fecal stream, waiting for the definitive procedure to be done. It is almost always a temporary procedure in compare to adult indications which can be permanent for malignant diseases. Colostomy can be divided into loop or divided, sigmoid or transverse. In spite of being a temporary procedure, there are serious complications that can occur from

improper technique and follow up care. When colostomy is incorrectly constructed, it can complicate or delay the management of these malformations and even can lead to death. A variety of complications with relatively high incidence have been reported. Even with careful technique, there is marked morbidity and mortality associated with the construction of colostomy. Despite this procedure being done commonly in pediatric age groups, few studies have been reported in local as well as in western literature on this issue.

In Sudan, there are little studies about the commonest indications of colostomy. Also, there are no clear rules about the constructional techniques of colostomy, regarding its type (sigmoid or transverse) or its mode (loops or divided). Also no clear

1. Senior registrar general surgery (SMSB)

2. Consultant pediatric surgeon

correlation between the incidence of complications and other factors related to level of operator or urgency of construction or prolong age of colostomy within the child's body...etc. This study is focusing on these issues, especially the complications and its outcome, hoping that we can provide away to decrease the incidence of this growing problem for a better life of children, and their families.

MATERIALS AND METHODS:

This is a prospective observational study, of all patients presenting to Al Ribat hospital for constructing colostomy or for final treatment, between December 2013 and December 2014. The hospital is a tertiary hospital with a capacity of 20 beds in the pediatric surgery department. The department has 3 consultants. In all the selected cases, a detailed and focused history and examination were done. The stoma site and type differed according to the preference of the operator. Colostomies were constructed by consultants, specialists and registrars. For sigmoid loop colostomy, the colon was brought out as a loop and prevented from retracting using nasogastric tube size 8 as a bridge and removed in most cases after 48 hours. For transverse colostomy, the same technique applied in the right hypochondrial area. Additional information was taken from the operation sheet about the type of anastomosis and suture materials used intraoperative and duration of postoperative fasting. All patients received antibiotics for different periods. All patients were checked again after 2 months either in the clinic or by telephone for further complications. Verbal consent obtained from all patients' relatives, data collected in questionnaire forms.

Analysis obtained using SPSS package version 20 and by applying Chi-square test, the level of statistical significance was $p < 0.05$. Results expressed in tables and graphs.

RESULTS:

Fifty patients were studied in the study. Residence of these children was as follows: center of Sudan 30 pts (60%), north of Sudan

7 pts (14%), west of Sudan 7 pts (14%), east of Sudan 3 pts (6%), south of Sudan 3 pts (6%). Boys were 35 (70%), girls were 15 (30%). The commonest age group was 4 years ($n=6$; 12%).

The most common indication of performing colostomy was Hirschsprung's disease (HSD) in 29 pts (58%). Anorectal malformations (ARM) are the causative factors in 19 pts (38%). Other causes include, perianal malignancy in 1 pt (2%), traumatic rectal injury in 1 pt (2%). Number of colostomies done in Al Ribat hospital was 31 (62%), while the number of colostomies done elsewhere were 19 (38%). Number of complicated colostomies was 24 pts (48%). The type of colostomy was sigmoid in 43 patients, complications occurred in 20 of them. Right transverse colostomy performed in 7 patients, complications occurred in 4 patients (57.1%). The ratio of complications between these two categories was (46.5% vs 57.1%) with a p -value = 0.602. Mode of colostomy was loop in 41 patients; complications occurred in 21. Divided colostomy done in 9 patients, complications happened in 3 of them. The ratio of complications between these two categories was (51.2% vs 33.3%) with a p -value = 0.331. The time of occurrence of complications was classified into groups. Those occurred within 1 month were 3 cases (12.5%). Those occurred between: 1 month -1 year were 10 cases (41.7%). Those occurred after 1 year were 11 cases (45.8%).

The total number of colostomies that underwent refashioning was 22 (91.7%). The other 2 (8.3%) complicated colostomies were observed in the referral clinic & there was no need for re-fashioning. Number of colostomies done as emergency was 31 (62%). Number of colostomies done in the elective list was 19 (38%).

The surgical staff that performed the colostomy was also classified into groups; consultant operated 9 cases, complications observed in 4 patients. Specialist operated 18 cases & complications observed in 8 patients. Registrars operated the rest of cases 23 & complications observed in 12 cases. The ratio of complications was high in registrars

(52,1%) in comparison to specialist and consultants (44,4% for each). P-value= 0.862.

Colostomies were complicated by stenosis occurred in 5 patients (20,8%), Prolapse observed in 7 patients, (29,2%), retraction occurred in 3 patients (12,5%) (Table 1).

Table (1): The type of Complications.

	Frequency	Percent
Stenosis	5	20.8
Hernia	1	4.2
Colostomy in aganglionic segment	1	4.2
Prolapse	7	29.2
Wound infection	3	12.5
Nec	1	4.2
Retraction	3	12.5
Faecaloma	1	4.2
I.o.	1	4.2
Death	1	4.2
Total	24	100.0

Unfortunately, colostomy was done in aganglionic segment in 1 patient and death occurred in 1 pt due to fecal peritonitis complicated by sepsis and multi organ failure.

CONCLUSION:

The study observes the frequency of indications and complications in 50 patients presented to al-ribat university hospital. The commonest indication was the Hirschsprungs disease and the prolapse was the largest complication group observed in these children. Nevertheless, no significant mortality reports were observed (2%).

Surgical training in pediatric colostomy must be improved for better outcome. The construction and the closure of colostomy should not be taken as a minor procedure and should be done by/under supervision of the most senior personnel.

A prompt postoperative care is essential, and parent education is of paramount importance to minimize colostomy morbidity.

REFERENCES:

1. A.F.Uba and L.B.Chirdan. Colostomy complications in children. *Annals of African medicine*. Vol 2, No.1; 2003:9-12.

2. Lister J, Webster PJ, Mirza S. Colostomy complications in children. *Practitioner* 1983; 227(1376): 229-37.

3. Mollitt DL, Malangoni MA, Ballantine TV, Grosfeld JL. Colostomy complications in children. An analysis of 146 cases. *Arch Surg* 1980; 115: 455-8.

4. Rickwood AM, Hemalatha V, Brooman P. Closure of colostomy in infants and children. *BrJ Surg* 1979; 66: 273-4.

5. Cilley RE, Statter MB, Hirschl RB, Coran AG. Definitive treatment of Hirschsprung's disease in the newborn with a one-stage procedure. *Surgery* 1994; 115: 551-6.

6. Nixon HH. Colostomy: a simple technique, and observations on indication. *Z Kinderchir* 1966; 3: 98-103.

7. Al-Salem AH, Grant C, Khawaja S. Colostomy complications in infants and children. *Int Surg* 1983; 77: 164-6.

8. Schier F, Antoniu D, Waldschmidt J. Complications of colostomy in childhood. *Zentralbl Chir* 1990; 15: 1435-40.

9. Ein SH. Divided loop colostomy that does not prolapse. *Am J Surg* 1984; 147: 250-2.

10. Gauderer MW, Izant RJ Jr. A technique for temporary control of colostomy prolapse in children. *J Pediatr Surg* 1985; 20: 653-5.

11. Golladay ES, Bernay F, Wagner CW. Prevention of prolapsed in pediatric enterostomas with purse string technique. *J Pediatr Surg* 1990; 25: 990-1.

12. Krasna IH. A simple purse string suture technique for treatment of colostomy prolapse. *J Pediatr Surg* 1979; 14:801-2.

13. Lau JT. Proximal end transverse colostomy in children. A method to avoid colostomy prolapse in Hirschsprung's disease. *Dis Colon Rectum* 1983; 26: 221-2.

14. Lau JR, Saing H, Ong GB. Double purse string suture technique for loop colostomy prolapse in infants. *Aust PediatrJ* 1982; 18: 58-9.

15. Narasimharao KL, Chatterjee H. A new technique of prolapse-free transverse colostomy. *Surg Gynecol Obstet* 1984; 158: 283.

16. Kelly MJ. Split loop colostomy: a modification. *Ann R Coll Surg Engl* 1995; 77: 313-14.

17. Lazar L, Kvalivker M, Erez I, Motovic A. Simple method of dilatation for strictured colostomy in children. *Dis Colon Rectum* 1993; 36: 199.

18. Pearl RK, Prasad ML, Orsay GP, Abcarian H, Tan AB, Melzl MT. Early local complications from intestinal stomas. *Arch Surg* 1985; 120: 1145-7.

19. Freund HR, Raniel J, Muggia-Sulam M. Factors affecting the morbidity of colostomy closure: a retrospective study. *Dis Colon Rectum* 1982; 25: 712-15.

20. Kirkegaard P, Luke M, Rasmussen JG, Christiansen J. Closure of terminal and loop colostomy. *Dis Colon Rectum* 1982; 25: 567-8.

21. Nmadu PT. Complications of colostomy closure in Zaria, Nigeria: a report of 70 cases. *Cent Afr J Med* 1990; 36: 287-91.
22. Samhoury F, Grodsinsky G. The morbidity and mortality of colostomy closure. *Dis Colon Rectum* 1979; 22: 312-14.
23. Yajko RD, Norton LW, Boemendal L, Eiseman B. Morbidity of colostomy closure. *AmJ Surg* 1976; 132: 304-6.
24. Brain AGL, Kiely EM. Use of single layer extramucosal suture for intestinal anastomosis in children. *Br J Surg* 1985 ;72: 483-4
25. Saleem M, Saqi Z, Shaikh AH, Malik N, Imran A, Shaikat M, et al. Complications of colostomy in infants and children. *Ann KE Med Coll* 1998; 4:20-3.
26. Wilkins S, Pena A. The role of colostomy in the management of anorectal malformations. *Pediatr Surg Int* 1998; 3: 105-9.
27. Rogers J. Hirschsprung's disease: diagnosis and management in children. *Br J Nurs* 2001; 10: 640-9.
28. Weber TR, Tracy TF Jr, Silen ML, Powell MA. Enterostomy and its closure in newborns. *Arch Surg* 1995; 130:534-7.
29. Nour S, Beck J, Stringer MD. Colostomy complications in infants and children. *Ann R Coll Surg Engl* 1996; 78:526-30.
30. Patwardhan N, Kiely EM, Drake DP, Spitz L, Pierro A. Colostomy for anorectal anomalies: high incidence of complications. *J Pediatr Surg* 2001; 36:795-8.
31. Rokhsar S, Harrison EA, Shaul DB, Phillips JD. Intestinal stoma complications in immunocompromised children. *J Pediatr Surg* 1999;34:1757-61.
32. Khan MNZ, Abbasi Z, Jan IA, Akhtar N, Shirazi H. Critical analysis of complications in 125 colostomies in children *Specialist* 1992; 8: 39-43.
33. Steinau G, Ruhl KM, Hornchen H, Schumpelick V. Enterostomy complications in infancy and childhood. *Langenbecks Arch Surg* 2001; 386:346-9.
34. Rolstad BS, Erwin-Toth PL. Peristomal skin complications: prevention and management. *Ostomy Wound Manage* 2004; 50: 68-77.
35. Millar AJ, Lakhoo K, Rode H, Ferreira MW, Brown RA, Cymes S. Bowel stomas in infants and children: a 5-year audit of 203 patients. *S Afr J Surg* 1993; 31:110-3.
36. Khan Y. Complications of colostomy and their management in infancy, Karachi (dissertation). *J Coll Physicians Surg Pak* 200
37. Chandramouli B, Srinivasan K, Jagdish S, Ananthkrishnan N.
38. Morbidity and mortality of colostomy and its closure in children. *J Pediatr Surg* 2004; 39:596-9.
39. Khan K, Khan MY, Waheed T. Management of colostomies in infancy. *J Postgrad Med Inst* 2003;17:7-10.
40. Muhammad Ali Sheikh, Jamshed Akhtar and Soofia Ahmed. Complications /problems of colostomy in infants and children. *JCPSP* 2006, Vol. 16 (8): 509-513