

Body Surface Area as a Surrogate Measure of Gallbladder Sizes and Indices: a Predictive Equation in Humans

Ugwu AC*

Abstract

Aims: To Find out the relationship between body surface area and gallbladder size; body surface area and gallbladder roundness index; and body surface area and gallbladder spherical index.

Methods: The length, width of 250 normal subjects comprising 140 males and 110 Females were sonographically measured. The gallbladder volumes, roundness index and spherical index were computed for each subject. Pearson's correlation established relationships between body surface area and gallbladder variables. A regression equation of gallbladder volume and body surface area was computed. Tests were two-tailed with $P < 0.05$ indicating statistical significance.

Results: There was a significant linear correlation between gallbladder size and body surface area ($r = 0.400$; $p = 0.007$). The correlation between body surface area and gallbladder roundness index ($r = -0.053$, $p = 0.731$) and body surface area and gallbladder spherical index ($r = -0.133$, $p = 0.389$) were not significant. An equation relating gallbladder volume and body surface area based on a simple linear regression gave $GBV (cm^3) = 32.036 BSA (m^2) - 25.943$.

Conclusion: Body surface area significantly correlates with gallbladder size and hence can be used to predict gallbladder sizes prior to cholecystosonography.

Keywords: Body surface area, gallbladder size, cholecystosonography



Introduction

Measurement of gallbladder size is an established diagnostic procedure in the assessment of gallbladder hydrops. Gallbladder spherical index which is a 3-dimensional equivalent of gallbladder roundness index has been suggested as a method for the assessment of gallbladder size. There is an increased fasting gallbladder volume in non-insulin-dependent diabetes mellitus¹. A positive correlation between fasting gallbladder volume and body weight have been reported². Exogenous motilin reduced fasting gallbladder volumes³ while long term oral L-arginine (L-Arg) increases fasting and postprandial gallbladder volume⁴.

Estimation of body surface area (BSA) is most commonly done with a formula that was derived in 1916 by Du Bois and Du Bois but remains the most popular way to estimate body surface area in nomograms or computer programs⁵.

An indirect method of predicting gallbladder sizes would be clinically invaluable. This would help in the estimation of gallbladder sizes in patients with suspected cases of hydrops or diabetes induced gallbladder dilatation.

To the best of our knowledge, no prediction formula or regression equation between gallbladder sizes and body surface area has been reported. In this study my aim was to study the relationship between body surface area and gall

bladder size, roundness and spherical indices in a descriptive environment and establish a predictive equation.

Subjects and methods

A total of 250 apparently healthy volunteers of Ibo tribe ranging from 18 years to 61 years were studied in Ebonyi state, Nigeria between January 2005 and November 2006. 140 were males and 110 Females. Studies were carried out with an abdominal 2 D ultrasound (siemens medical systems, USA Inc, ultrasound Group, Issaquah WA) with a 3.5 MHz sector transducer. Subjects with history of hepatobiliary disease were not included. Diabetic patients were also not included in the study. The study was approved by the local ethical committee of Jeomedics ultrasound centre and informed consent was obtained from each subject.

Each subject was scanned in the morning after an overnight fasting, having previously given an informed consent to participate in the study. Subjects had their weight measured on a bathroom weighing scale: model H 89 LT Blue, and the height measured on a calibrated vertical wall.

Subjects were scanned on supine position. After visualization of the maximal gallbladder longitudinal outline, the length and maximal antero-posterior diameter (height) measurements were taken on arrested respiration with calipers crossing each other at 90°. Subsequently the probe was rotated through 90° to obtain the maximal transverse dimension. All measurements were taken thrice and the maximum value recorded. The length, width and height

* Radiology Department, Federal Medical Centre, P.M.B 102, Abakaliki, Ebonyi State, Nigeria. (+2348076241297, Email: tonybullng@yahoo.co)

measurements were taken from the outer to outer hyperechoic walls of the gallbladder (Fig 1a and 1b). Gallbladder volumes were calculated using ellipsoid formula. The gallbladder roundness index (GBRI) (in percentage) was calculated⁶: antero-posterior diameter/length X 100/1. The gallbladder spherical index (GBSI) was calculated¹:

$$\frac{(\text{Antero-posterior diameter} + \text{width})}{2 \times \text{length}} \times 100/1$$

Body surface area (m²) was obtained using the formula derived by Du Bois and Du Bois⁵; BSA = (Weight^{0.425} X Height^{0.725}) X 0.007184, where weight is in kilograms and height in centimeters.

Values were obtained by one imaging scientist (AC) to avoid possible inter-observer variations. Summary statistics were generated using SPSS 11.0 software while graph was plotted and regression equation was developed in MS excel office 2000. Correlation coefficients (r) and linear regression analysis were performed to assess the relationship between body surface area and gallbladder size. Correlations between BSA and gallbladder indices (GBSI and GBRI) were also calculated. Tests were two-tailed with P < 0.05 indicating statistical significance.

Result

The result of the statistical analysis of the data on body surface area and gallbladder parameters are shown in Table 1.




Fig 1a : A line drawing showing a longitudinal section of the gallbladder.




Fig 1b: A line drawing showing a transverse section of the gallbladder.

Table 1. Descriptive statistics of subjects

| Variable | N | Minimum | Maximum | Mean | Std. Deviation | Skewness | Kurtosis |
|----------|-----|---------|---------|---------|----------------|----------|----------|
| BSA | 250 | 1.35 | 2.17 | 1.7573 | 0.17301 | -0.123 | 0.450 |
| GBV | 250 | 7.78 | 71.21 | 29.2893 | 13.75489 | 1.320 | 1.777 |
| GBRI | 250 | 22.54 | 55.38 | 42.2382 | 8.34084 | -0.190 | -0.745 |
| GBSI | 250 | 26.76 | 57.69 | 42.2509 | 8.37422 | 0.167 | -1.070 |

There was a significant linear correlation between body surface area and gallbladder sizes ($r = 0.4$, $P = 0.007$). The BSA did not correlate significantly with GBRI and GBSI. The Pearson's correlation coefficient between BSA and GBSI was $r = -0.133$, $P = 0.389$ while the correlation

between BSA and GBRI was $r = -0.053$, $P = 0.731$.

Fig 2 is a graph of gallbladder size (cm³) against BSA (m²). A regression equation of gallbladder volume on BSA gives: $GBV = 32.036$ BSA -26.943 .




Fig 2: A graph of Gall Bladder Volume (cm³) against Body Surface Area (m²)

Discussion

Oral cholecystography, which depends on adequate absorption of the oral contrast medium from the small intestine, its excretion from the liver with bile, the patency of the cystic duct and finally adequate concentration by the gallbladder, was the first popular technique for the assessment of gallbladder sizes and structural changes. With the increasing use of high-resolution sonography in the gallbladder, sonography has become the commonest imaging modality in evaluating gallbladder sizes.

Gallbladder contours differ much in humans. In the assessment of gallbladder function, a study⁷ has shown that the estimation of gallbladder volume based on one direct measurement of length exhibits satisfactory precision to be applied in all the situations where the assessment of gallbladder function should be more rapid and simple. The accuracy of this method can be even greater than that of the widely accepted and applied ellipsoid method^{8,9} since the former method is based only upon one gallbladder

parameter. When the new method is applied, it is recommended to select the proper formula depending on gallbladder shape⁷.

When the shape resembles the ellipsoid or the sphere, the ellipsoid method is recommended. When the gallbladder is thinner, the use of method of cone¹⁰ is suggested. When the gallbladder is rather thick and exhibits atypical shape, the method of cylinder¹¹ can provide more accurate results.

In this study, a significant positive correlation was established between BSA and gallbladder volume, while correlations between BSA and gallbladder indices (GBSI and GBRI) were noted to be insignificant. This result is similar to a previous study², which established a positive correlation between gallbladder volume and body weight, body mass index and body fat weight. The significant finding in this study agrees with a previous study¹² which established a moderate and significant correlation between body surface area and gallbladder volume ($r=0.57$; $p<0.01$).

To the best of our knowledge, this is the first time a prediction equation for gallbladder volume was established.

An obvious limitation in this study was the use of ellipsoid method for the evaluation of all the assessed gallbladder (shapes). Future studies adopting the supposed formula for different gallbladder shapes are advised. Three dimensional ultrasonography is fast becoming highly utilized in the ultrasound laboratory and may well prove to be the appropriate technology in future studies¹³. In this study, all the measurements were taken by one imaging scientist (AC). This was done to avoid possible interobserver errors, as it has not been previously investigated. Future studies on intra- and inter-observer variation in the 2 dimensional assessment of gallbladder dimensions and volumes are recommended. Despite these obvious limitations, this study has shown that body surface area significantly correlates with gallbladder volume and hence can be used to predict gallbladder sizes (± 2 standard deviation) prior to cholecystosonography.

Acknowledgement

The authors acknowledge with deep appreciation, Mr Chy Femi for his assistance in statistical analysis.

References

1. Guliter S, Tlmaz S, Karakan T. Evaluation of gallbladder volume and motility in non-insulin-dependent diabetes mellitus. *Journal of clinical Gastroenterology* 2003; 37 (4): 288 -- 291.
2. Sari R, Balci MK, Coban E et al. Sonographic evaluation of gallbladder volume and ejection fraction in obese women without stones. *Gastroenterology* 1990; 98 (4): 1001-1007
3. Luiking Y C, Peeters T L, Stolk MF et al. Motilin induces gallbladder emptying and antral contractions in the fasted state in humans. *Gut* 1998; 42 (6): 830-835.
4. Luiking T C, Weusten BLAM, Portincasa P. et al Effects of long – term oral L-arginine on esophageal motility and gallbladder dynamics on healthy humans. *Am J Physiol Gastrointest Liver Physiol* 1998; 274 (6): 984-991.
5. Du Bois D, Du Bois EF. A formula to estimate the approximate surface area if height and weight be known. *Arch Intern Med* 1916; 17:863-871.
6. Gourtsoyiannis N C, Damilakis J E, Cheroulakis N Z et al. Relationship of gallbladder contour, fasting volume and emptying to body size indices in normal subjects and patients with gallstones. *Digestion* 1995, 56 (5): 395-399.
7. Romanski K W. Accuracy of estimation of canine gallbladder volume from one measurement. *Bull Vet Inst Pulawy* 2004; 48:159-162.
8. Dodds WJ, Groh WJ, Darweesh RMA et at. Sonographic measurement of gallbladder volume. *AJR* 1985; 145:1009-1011.
9. Jonderko K, Ferre LP, Buerio L. Real-time ultrasonography as a non-invasive tool for the examination on canine gallbladder emptying. A validation study. *J pharmacol Toxicol Meth* 1992; 27: 107-111.
10. Cano N, Cicero F, Ranieri F et at. Ultrasonographic study of gallbladder motility during total parental nutrition. *Gastroenterology* 1986; 91:131-317.
11. Romanski K. Evaluation of the methods used for the estimation of gallbladder volume. *Ultrason Pol* 2003; 13:62-66.
12. Yoo J, Kwak H, Lee M, Suh J, Rhee C. Sonographic measurements of normal gallbladder sizes in children. *Journal of clinical ultrasound* 2003; 31(2):80-84.
13. Hashimoto S, Goto H, Hirooka Y et al. An evaluation of three-dimensional ultrasonography for the measurement of gallbladder volume. *American Journal of Gastroenterology* 1999; 94:3492