



Fatal abomasal sand impaction in a giraffe calf (*Giraffa camelopardalis*) at the University of Ilorin zoological garden

HO Jegede^{1*}, AY Adenkola², A Obalolu¹, FR Olowoleni¹ & PO Odeniran³

- ^{1.} *Veterinary Teaching Hospital, University of Ilorin, Ilorin, Nigeria*
- ^{2.} *Department of Physiology and Pharmacology, College of Veterinary Medicine, University of Agriculture, Makurdi, Nigeria*
- ^{3.} *Department of Veterinary Microbiology and Parasitology. Faculty of Veterinary Medicine. University of Ibadan, Ibadan, Nigeria*

*Correspondence: Tel.: +2348038070602, E-mail: drlanrejeg@gmail.com

Abstract

A post-mortem examination was carried out on a 4-month-old giraffe which was reported dead early hours of the morning in the zoological garden, University of Ilorin. The carcass of the animal appeared slightly emaciated and on opening of the carcass the abomasum was distended with a hard mass felt inside the organ. On opening of the organ, it was filled with sand and weighing 3.8kg. Geophagia due to various factors were queried in the cause of the condition including seasonal prevalence, nutrient deficiencies, feeding regimen and also housing inadequacies. Although poor milk intake, absence of maternal nurturing and inadequate captive conditions are the most likely causes of geophagia which eventually led to the death of the animal.

Keywords: Abomasum, Sand impaction, Geophagia, Giraffe, Nigeria

Received: 16-10- 2015

Accepted: 10-11-2015

Introduction

Sand impaction occurs when sand adheres to food, soil is ingested directly, or dirt is eaten, as has been reported in foals of domestic horses (Radostits *et al.* 2000).

Sand impaction occurs in a wide variety of animals including cattle, elephants (*Elephas maximus*) (Warren *et al.* 1996), and ostriches (*Struthio camelus*) (Mushi *et al.* 1998), as well as humans (Singh, 1983). It has been reported in a cria by Sameeh & Lyall (2006) where the animal was observed eating soil.

Clinical signs of sand impaction in other animal species include depression; either diarrhea or little or no passage of feces; and mild, moderate, or even severe colic with mucosal damage (Johnston & Freeman, 1997; Fowler, 1998).

Geophagia, the deliberate ingestion of soil, has been classified as a form of pica. Geophagia in mammals has been associated with deficiencies of elements such as phosphorus, sodium, magnesium, sulphur, copper, cobalt and manganese (Kreulen & Jager, 1984). Trace element deficiencies, in particular copper, zinc and cobalt; have been incriminated in the aetiology of alopecia and wool eating habit in sheep (Fahmy *et al.*, 1980; Meyer & Lohse, 2002).

Numerous health problems that are suspected to be of nutritional origin have been documented in captive giraffe. Pathologies that may relate to vitamin and mineral intake or metabolism include white muscle disease (Strafuss & Kennedy, 1973; Burton & Dierenfeld), urolithiosis (Wolfe *et al.*, 2000), and dental disease (Enqvist, 2003). Pancreatic pathologies (Fox, 1938; Fowler, 1978; Lechowski *et al.*, 1991; Ball *et al.*, 2002), decreased ruminal absorptive surface area (Hofmann & Matern, 1988), ruminal acidosis (Clauss, 1998; Clauss *et al.*, 2002), fermentative gastritis or rumenitis (Fox, 1938; Ball *et al.*, 2002) and gastrointestinal ulceration (Fox, 1938; Fowler, 1978) also have been documented. This study looks to provide wildlife veterinarians and biologists working in captive conditions e.g. zoos, with best husbandry and management practices to avoid future mortalities to their captive giraffes especially calves.

Case History

The giraffe arrived at the University of Ilorin zoological garden in March, 2012 with the umbilical cord still attached indicating the animal



Plate 1: Emaciated carcass

was less than 2 months old (Langman, 1977). The animal was fed with about 4 litres of whole milk (cattle) twice daily for the first month of introduction to the zoo without any mineral supplementation. It was housed in a 7 by 7 foot pen with sharp sand as bedding material. On the second month of introduction the animal, milk feeding was ceased; instead the animal was fed about 1 kg of wheat offal, guinea corn and leaves of the African copaiba balsam tree (*Igi-iyá*) *Daniellia oliveri* (Quattrocchi, 2012) as food. The animal was found dead 3 weeks later by the zoo workers. No signs of colic or abdominal discomfort were noticed before closing hours a day prior to death.

A postmortem examination was carried out in an attempt to diagnose the cause of death. The entire carcass was slightly emaciated (Plate 1) with a resolved wound on the medial aspect of the right forelimb at the level of the carpus. On opening of the carcass it was noticed that the abomasum was very large and distended with a hard mass felt inside. On dissecting through the gastrointestinal tract, there were large deposits of sand filling the abomasum which weighed 3.8kg (Plate 2). No other peritoneal lesions were observed. All other organs appeared normal.

Discussion

First-year giraffe calf mortality in captivity may be as high as 45% as reported by Lackey & LaRue, (1997) and as high as 75% in the wild due to predation (Foster, 1966). Wasting and sudden death are frequently reported in the literature and anecdotally; malnutrition and peracute mortality syndrome (Fox, 1938), serous fat atrophy and peracute mortality syndrome (Fowler, 1978), mineral imbalances and wasting (Junge & Bradley, 1993), poor milk intake (Flach *et al.*, 1997), hypoglycemia and chronic energy malnutrition (Ball *et al.*, 2002), therefore the sudden death is



Plate 2: Enlarged abomasum filled with sand

not a strange occurrence in giraffes. At this time, the true proportion of captive giraffe mortality caused by nutritional pathologies is unknown.

Geophagia and osteophagia is a common feature of the feeding routine of the southern giraffe (*Giraffa camelopardis giraffa*) during the months from April to November and geophagia was primarily exhibited by sub-adult giraffe (Langman, 1978). Although all age classes can be affected as it is a common behaviour in free-ranging giraffe (Seeber *et al.*, 2012). The ultimate cause of the geophagia is still uncertain (Neser, 2001). This incidence occurred in the month of May.

Due to the association of the condition to mineral deficiencies (Kreulen & Jager, 1984); A deficiency in cobalt has been associated with pica in cattle (Kreulen & Jager, 1984) and a zinc deficiency has been implicated as a cause of geophagia in children (Hambidge *et al.*, 1987). It is possible that lack of supplementation of essential minerals in the milk could also be the cause of the problem as hand reared giraffe calves should be given supplements like calcinol, evion and other feed concentrates in milk (Khadri & Valandikar, 2002). In calves and sheep that are temporarily deserted by their dams, hunger was suggested as a driver for geophagy (Neser, 2001). The recorded quantity to be fed to hand reared giraffe calves at 2-4 months is 1.2-1.6 litres of milk 6 times daily and supplement with browse and concentrates (Khadri & Valandikar, 2002).

The use of sharp sand as bedding gave an avenue and availability of sand to be consumed hence materials like straw, saw-dust or mulch can be used instead (Lorraine, 2003). Control of sand impaction has been achieved by preventing access to pastures that contain sand and by not feeding animals on the ground (Johnston & Freeman, 1997).

Hediger (1955) and Hediger (1964) considered restricted movement due to space limitations to be one of the primary contributors to captivity-induced stress, and some of the earliest studies of abnormal behaviour in captive animals supported this contention (Levy, 1944). More recent work has also suggested an adverse effect of confinement in small spaces. In one study of the impact of captivity on 35 different species of carnivore, infant mortality in captivity and stereotypic locomotion in the form of pacing was found to correlate positively with species home range size in the wild (Clubb & Mason, 2003; Clubb & Mason, 2007). The 7 by 7 foot pen was therefore considered grossly inadequate and a larger perimeter fencing of 200 by 300 feet was constructed.

References

- Ball RL, Kearney C, Burton M, Dumoneux G & Olsen JH (2002). Morbidity and mortality related to hypoglycemia and chronic energy malnutrition in captive giraffe. In: *Proceedings of 35th Annual American Association of Zoo Veterinarians Conference held Milwaukee, Wisconsin*. Pp 180-185.
- Burton MS & Dierenfeld ES (1990). Comparisons of vitamin E injectables in giraffes (*Giraffa Camelopardalis reticulata*). In: *Proceedings of 23rd Annual American Association of Zoo Veterinarians Conference held Padre Island Texas*. Pp 198-201.
- Clauss M (1998). Feeding giraffe (*Giraffe camelopardis*) Unpublished Masters thesis. Royal Veterinary College, University of London. London, UK: University of London. Pp 114.
- Clauss M, Lechner-Doll M, Flach EJ, Wisser J & Hatt J-M (2002). Digestive tract pathology of captive giraffe (*Giraffa camelopardalis*) an unifying hypothesis. *European Association of Zoo and Wildlife Veterinarians Fourth Scientific Meeting*, Heidelberg, Germany. Pp 99-107.
- Clubb R & Mason G (2003). Captivity effects on wide-ranging carnivores. *Nature*, **425**(7): 473-474.
- Clubb R & Mason G (2007). Natural behavioural biology as a risk factor in carnivore welfare: How analysing species differences could help zoos improve enclosures. *Applied Animal Behaviour Science*. **102**(3-4): 303-328.
- Enqvist KE (2003). Dental disease and serous atrophy of fat syndrome in captive giraffes (*Giraffa camelopardalis*). Minneapolis, MN: In: *Proceedings of the 36th Annual conference of the American Association of Zoo Veterinarians conference held at Minneapolis, Minnesota*. Pp 86-89.
- Fahmy F, Amer AA, Abd-el-Aziz H & Abd-el-Roaf M (1980). Wool as an effective tool of some deficiency diseases. *Assiut Veterinary Medical Journal*, **7**(13/14): 262-270.
- Flach EJ, Thornton SM, Kirkwood JK & Sainesbury AW (1997). Chronic loss of condition with persistent neutrophilia in a reticulated giraffe (*Giraffa camelopardalis*). In: *Proceedings of the 33rd Autumn British Veterinary Zoological Society*. London, UK. Pp 33-37.
- Foster JB (1966). The giraffe of Nairobi National Park: home range, sex ratios, the herd, and food. *East African Wildlife Journal*, **4**(1) 139-148.
- Fowler ME (1978). Peracute mortality in captive giraffe. *Journal of American Veterinary Medical Association* **173**(9): 1088-1093.
- Fowler ME (1998). *Medicine and Surgery of South American Camelids. Second edition*, Iowa: State University, Iowa. Pp 364- 370,337-350.
- Fox H (1938). The giraffe. Some notes upon the natural characters of this animals, its care and its misfortune. In: *Zoo, editor. Report of the Penrose Research Laboratory, formerly the Laboratory and Museum of Comparative Pathology*. Philadelphia, PA: Zoological Society of Philadelphia. Pp 35-67.
- Graham TW, Thurmond MC, Mohr FC, Holmberg ML & Keen CL (1994). Relationships between maternal and foetal liver copper, iron, manganese and zinc concentrations and fetal development in California

- Holstein dairy cows. *Journal of Veterinary Diagnostic Investigation*, **6**(1): 77–87.
- Hambidge KM, Casey CE & Krebs NF (1987). Zinc. In: *Trace Elements in Human and Animal Nutrition. Fifth edition*, (Mertz W, editor). Academic Press Inc., San Diego, USA, 2: 50–51.
- Hediger H (1955). *Studies of the Psychology and Behavior of Captive Animals in Zoos and Circuses*. Criterion Press, New York. Pp 3-15.
- Hediger H (1964). *Wild Animals in Captivity: An Outline of the Biology of Zoological Gardens*. Dover Publications, New York. Pp 6-33.
- Hofmann RR & Matern B (1988). Changes in gastrointestinal morphology related to nutrition in giraffes. *Conservation Science and Zoos*, **22**(3):168-176.
- Johnston JK & Freeman DE (1997). Diseases and surgery of the large colon. *Veterinary Clinics of North America: Equine Practice*. **13**(4): 317–340.
- Junge RE & Bradley TE (1993). Peracute mortality syndrome of giraffes. In: *Zoo and Wild Animal Medicine: Current Therapy 3*. (Fowler ME, editor) Philadelphia, PA: Saunders Company. Pp 547-549.
- Khadri SSMS & Valandikar SC (2002). Hand rearing of giraffe calves at Sri Chamarajendra Zoological gardens, Mysore. *Indian Forester*. Pp 1153-1158.
- Kreulen DA & Jager T (1984). The significance of soil ingestion in the utilization of arid rangelands by large herbivores with special reference to natural licks on the Kalahari pans. In: *Herbivore Nutrition in the Tropics and Subtropics (FMC Gilchrist & RI Mackie, editors)*. The Science Press, Craighall, South Africa. Pp 204 - 221.
- Lackey BL & LaRue FV (1997). *North American Regional giraffe Studbook*. Dallas: Dallas Zoo.
- Langman VA (1977). Cow-calf relationships in giraffe (*Giraffa camelopardalis giraffa*). *Zeitschrift fuer Tierpsychologie*, **43**:265-286.
- Langman VA (1978). Giraffe Pica Behavior and Pathology as Indicators of Nutritional Stress. *The Journal of Wildlife Management* **42**(1): 141-147
- Lechowski R, Pisarski J, Goslawski J & Lenarcik M (1991). Exocrine pancreatic insufficiency-like syndrome in giraffe. *Journal of Wildlife Diseases* **27**:728-780.
- Levy DM (1944). On the problem of movement restraints, stereotyped movements, hyperactivity. *American Journal of Orthopsychiatry*. **14**(7): 644–671.
- Lorraine J (2003). *Giraffe Husbandry Manual*. <http://www.aszk.org.au/husbandry.mam.mals.ews>, retrieved 14-07-2015.
- Meyer H & Lohse K (2002). Ca and P supply of ruminants in the 19th and beginning of 20th century in Middle Europe. *Dtsch Tierarztl Wochenschr Journal*, **109**(1): 34-37.
- Mushi EZ, Isa JFW, Chabo RG, Binta MG, Modisa L & Kamau JM (1998) Impaction of stomach ostriches (*Struthio camelus*) in Botswana. *Avian Disease*, **42**(5):597–599.
- Neser JA (2001). Enzootic geophagia of calves and lambs in the Northern Cape and North-West Provinces of the Republic of South Africa, and the possible role of chronic manganese poisoning. *Applied Animal Husbandry & Rural Development Journal*, **2**(1): 9-12.
- Quattrocchi U (2012). *CRC World Dictionary of Medicinal and Poisonous Plants: Common Names, Scientific Names, Eponyms, Synonyms, and Etymology (5 Volume Set)*. Pp 1328.
- Radostits OM, Gay CC, Blood DC & Hinchcliff KW (2000). *Veterinary Medicine, A Textbook of the Diseases of Cattle, Sheep, Pigs, Goats and Horses. Ninth edition*. London: WB Saunders. Pp 229–230.
- Sameeh MA & Lyall P (2006). Fatal sand impaction of the spiral colon in a 1-month-old alpaca. *The Canadian Veterinary Journal*, **47**(7): 683–684.
- Seeber PA, Ndlovu HT, Duncan P & Ganswindt A (2012). Grazing behaviour of the giraffe in Hwange National Park, Zimbabwe. *African Journal of Ecology*, **50**(2): 247–250.
- Singh M (1983). Sandy fecal impaction caused by severe pica. *Tropical and Geographical Medicine*. **35**(6):393–394.
- Strafuss AC & Kennedy GA (1973). Degenerative myopathy in a giraffe. *Journal of the American Veterinary Medical Association* **163**(6): 551-552.
- Warren K, Bolton J, Swan R, Gaynor W & Pond L (1996). Treatment of gastrointestinal tract impaction of a 2-year-old Asian elephant (*Elephas maximus*) *Australian Veterinary Journal*. **73**(1):37–38.
- Wolfe BA, Sladky KK & Loomis MR (2000). Obstructive urolithiasis in a reticulated giraffe (*Giraffa camelopardalis reticulata*). *The Veterinary Record* **146**(9): 260-261.