



Seroprevalence of infectious bursal disease virus antibodies in some species of poultry in Maiduguri, Nigeria

YM Shettima¹, AD El-Yuguda¹, OD Oluwayelu², MB Abubakar¹, MY Zanna¹, TM Hamisu^{1*}, MM Maina¹, SS Baba¹ & A Ali¹

1. Department of Veterinary Microbiology, Faculty of Veterinary Medicine, University of Maiduguri, Borno State, Nigeria
2. Department of Veterinary Microbiology and Parasitology, Faculty of Veterinary Medicine, University of Ibadan, Oyo State, Nigeria

*Correspondence: Tel.: + 2348068783895; E-mail: alabsola@yahoo.com

Copyright: © 2018 Shettima *et al.* This is an open-access article published under the terms of the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Publication History:
Received: 04-06-2017
Accepted: 24-11-2017

Abstract

This study was aimed at determining the antibodies of IBDV in some poultry species in Maiduguri, Nigeria. A total of 944 serum samples were collected from village chickens, broilers, layers, ducks, turkeys and geese in Maiduguri and tested for IBDV antibodies using inzyme linked Immunosorbent assay (ELISA) and a seroprevalence of 46.6% was recorded. The species distribution showed that 33.4% of local chickens (134/401), 67.3% of layers (35/52), 50.8% of broilers (159/313), 60.6% of turkeys (77/127), 65.5% of ducks (19/29) and 72.7% of geese (16/22) sampled were positive for IBDV antibodies. Males showed a seroprevalence of 70% while females recorded 30%. IBDV seropositive sera showed that samples from broilers reacted with 31.5% middle OD values and turkeys with 9.4% middle OD values. Samples from other species reacted with lower OD values. Samples from other species reacted with lower OD values. Presence of IBDV antibodies in species other than chickens suggested that different bird species might have IBDV and could serve as reservoirs for IBDV transmission. Because of this threat to poultry industry, there is need for continuous surveillance of IBDV in all poultry species so as to institute effective preventive measures against the disease.

Keywords: ELISA, Infectious bursal disease virus, IgG, Poultry species, Serum samples

Introduction

Infectious bursal disease (IBD) also called Gumboro, avian infectious bursitis or avian phrosis-nephritis, is an acute contagious disease of young birds 3-6 weeks and even 8 weeks of age (El-Yuguda & Baba, 2004) characterized by destruction of lymphoid cells in the bursa of fabricius (Abdu *et al.*, 1986; Oluwayelu, 2010). Infectious bursal disease is caused by infectious bursal disease virus, a bisegmented double stranded RNA belonging to the genus *Avibirnavirus* in the family *Birnaviridae* (Parkhurst, 1964; Lukert & Saif, 1997). Both village and

commercial chickens are equally affected (Akoma & Baba, 1995; El-Yuguda & Baba, 2002). The virus can be transmitted mainly from infected to susceptible birds through water, feeds, droppings and fomites (Akoma & Baba, 1995). Other means of transmission such as through the intermediary role of vermines like the lesser meal worm, mites and mosquitos have also been reported (Abdu *et al.*, 1986). The disease is characterized by a short incubation period of 2-3 days (Awolaja & Adene, 1995). Clinical signs include depression, mucoid or watery diarrhoea,

dehydration, rapid loss of weight, high morbidity of up to 100% and a mortality of 20-30% with a rapid recovery in 5-7 days (Lasher & Davis, 1997). The cloacal bursa and spleen are used for the isolation of the virus (Lukert & Saif, 2003). The virus can be found in other organs such as thymus, liver and bone marrow but in significantly low quantities than in bursa (Cheville, 1967; Tanimura & Sharma, 1997). Primary cell structures of chicken embryo fibroblasts (CEF), bursa, and kidney have been used to propagate the virus. McFerran *et al.* (1980) reported that embryonating eggs have also been used with good result. Serological tests generally used for the detection of IBDV antibodies are ELISA, viral neutralization and agar gel immunodiffusion test. However, ELISA is the most commonly used test for the detection of antibodies to IBDV (Lukert & Saif, 2003). No treatment has been found to have an effect on the course of IBD and in addition, there are no reports of the use of antiviral compounds and interferon inducers for the treatment of IBD (Lukert & Saif, 2003). Consequently, proactive measures must be in place to prevent and control IBD occurrence. To achieve this, a regular and sustained surveillance of the disease is needed to know the current status in the study area and this is what this research was aimed to achieve.

Materials and Methods

Study area

This study was carried out in Maiduguri, Borno State, Nigeria. Maiduguri is the capital city of Borno State and lies between 10.20° & 13.40° North and 9.80° & 14.40° East and occupies an area of 69.436km². Borno state shares international borders with Niger to the north, Chad to the north east and Cameroon to the east (Waziri *et al.*, 2009).

Study population

A total 944 birds comprising local chickens (401), broilers (313), layers (52), turkeys (127), ducks (29) and geese (22) were used in the study. The male broilers were separated from females by their noticeable nature of big and brighter combs and wattles and thicker legs and feet.

Sampling

Blood samples were collected from different species of poultry using convenient sampling technic. The samples were collected from poultry at slaughter

slabs at the point of slaughter (554); live bird markets (60); commercial farms and backyard poultry (330) using 23G sterile hypodermic needles and 2ml syringes. The blood samples were collected into plain vacutainer tubes and allowed to clot at room temperature. The sera were harvested into appropriately labeled cryotubes and stored at -20°C until used.

Serology

The sera were tested for the presence of IBDV antibodies using commercial ELISA kit (X-Ovo FlockscreenTM) following the manufacturer's protocol in Virus Research Laboratory, Department of Veterinary Microbiology, Faculty of Veterinary Medicine, University of Maiduguri. Briefly, 50µl of diluted serum was added to each well using multichannel automated pipette and incubated at 37°C for 30 minutes and then washed to remove excess unbound antibodies. An alkaline phosphate labeled rabbit anti-chicken IgG conjugate was added and then incubated at 37°C for 30 minutes. The plates were washed four times with wash buffer (300µl per well) to remove unbound conjugate and phenolphthalene monophosphate (PMP) substrate was added to the wells and then incubated at 37°C for 30 minutes. The reaction was then stopped using the stop solution (sodium hydroxide). The degree of colour development (optical density) greater than 0.306 is considered positive while less than that is considered negative. In general, optical density value less than 0.306 is considered low OD value, while OD values equals and greater than 0.306 are considered middle and high respectively.

Results

The result of this study showed that IBDV antibody was detected in 440 (46.6%) of the 944 serum samples collected from some species of poultry. The species distribution showed that the sera of 33.4% of local chickens, 67.3% of layers, 50.8% of broilers, 60.6% of turkeys, 65.5% of ducks and 72.7% of geese were positive for IBDV antibodies (Table 1).

The sex distribution of IBDV seropositive samples showed prevalence of 70% among males and 30% among females (Table 2). IBDV seropositive sera showed only the broiler samples reacted with 31.5% middle OD values followed by turkeys with 9.4% middle OD values and all the remaining samples reacted with lower OD values (Figure 1).

Table 1: Distribution of IBDV antibodies among some species of poultry in Maiduguri, Nigeria

Poultry Type	Total number tested	Number (%) Positive
Local chicken	401	134 (33.4)
Layers	52	35 (67.3)
Broilers	313	159 (50.8)
Turkeys	127	77 (60.6)
Ducks	29	19 (65.5)
Geese	22	16 (72.7)
Total	944	440 (46.6)

Table 2: Distribution of IBDV antibodies based on sex among some species of poultry in Maiduguri, Nigeria

Poultry Species	No. tested	Males		Females	
		No. (%) positive	No. tested	No. (%) positive	
Local chicken	251	81 (32.3)	150	53 (35.3)	
Layers	NA *	NA	52	35 (67.3)	
Broilers	313	159 (50.7)	0	0 (0)	
Turkeys	57	51 (89.4)	70	26 (37.1)	
Ducks	15	12 (80)	14	7 (50)	
Geese	7	5 (71.4)	15	11 (73.3)	
Total	643	308 (47.9)	301	132 (43.8)	

Key: NA*= Not Applicable

Discussion

IBDV infection has been reported to be widespread in major poultry producing countries (Cardona *et al.*, 2000), and this study also confirmed the presence of antibodies to IBDV in the study area. The specie/type specific prevalence rate shows that geese recorded the highest prevalence of 72.7%; this is followed by layers and ducks with 67.3% and 65.5% respectively. This higher prevalence might have resulted from natural infection or from close contacts with commercial chickens, since geese are not routinely vaccinated against IBDV. Similarly, the prevalence of 65.5% in ducks in this study is higher than the 19.1% reported by Oluwayelu *et al.* (2007) and 0.0% reported by Mai *et al.* (2004). The higher prevalence recorded here may be attributed to the highly sensitive and specific nature of the diagnostic tool employed (ELISA test) when compared to the less sensitive AGID used by the authors (Oladele *et al.*, 2001). The high prevalence observed among layers (67.3%) in this study may be attributed to vaccination.




Figure 1: Bar chart showing the optical density (OD) values of IBDV antibody positive samples

Key: B= broilers; L= layers; T= turkeys; G= geese; D= ducks; VC= village chickens

In this study, local chickens showed a seroprevalence rate of 33.4%. This is lower than 63.5% reported by Lawal *et al.* (2014). The lower prevalence can be attributed to small sample size (944) in comparison to that analyzed by Lawal *et al.* (2014) (1500). It is however higher than 7.7% prevalence reported by Mbuko *et al.* (2010).

Khan *et al.* (2009) reported a prevalence rate of 7.75% in broiler in the district of Peshawar, Pakistan. This is lower than 50.8% reported in this study. This variation may be because Khan *et al.* (2009) reported their prevalence on the basis of history and detailed post-mortem examination.

The sex distribution of the IBDV antibodies observed in this study showed that males had higher seroprevalence rate (70%) than females (30%) (Table 2). The higher prevalence in males in this study could be due to the fact that male birds, which are mostly

broilers were sampled more than the female birds in this study.

The detection of IBDV antibodies in the sera of apparently healthy, free-roaming turkeys, geese, local chickens and ducks in this study indicates natural exposure to the virus and implicates them as a potential source of infection to commercial chickens since neither vaccination nor other preventive measures are practiced in local chickens, turkeys, ducks and geese in Nigeria.

References

- Abdu PA, Abdullahi SU, Adesiyun AA & Ezeokoli CD (1986). Infectious bursal disease. *World Poultry Science Association Journal*, **42**(3): 219-229.
- Akoma MB & Baba SS (1995). Survey for infectious bursal disease virus antibodies in free and commercial birds in Borno State, Nigeria. *Studies and Researches in Veterinary Medicine*, **3**: 46-48.
- Awolaja AO & Adene (1995). Outbreak of infectious bursal disease in vaccinated poultry flocks. *Tropical Veterinarian*, **13**(182): 37-43.
- Cardona CJ, Oswald WB & Schat KA (2000). Distribution of chicken infectious anemia virus in the reproductive tissue of specific pathogen-free chickens. *Journal of General Virology*, **81**(8): 2067-2075.
- Cheville NF (1967). Studies on the pathogenesis of Gumboro disease in the bursa of fabricius, spleen and thymus of the chicken. *American Journal of Pathology*, **51**(4): 527-551.
- Cosgrove AS (1962). An apparently new disease of chicken- Avian Nephrosis. *Avian Disease*, **6**(3): 385-389.
- El-Yuguda AD & Baba SS (2002). Prevalence of selected viral infections in various age group of village chickens in Borno State, Nigeria. *Nigerian Journal of Animal Production*, **29**(2): 245-250.
- El-Yuguda AD & Baba SS (2004). An outbreak of infectious bursal disease in eight weeks old IBD vaccinated commercial poultry flock in Maiduguri, Nigeria. *Tropical Veterinarian*, **22** (3 and4): 93-98.
- Khan RW, Khan FA, Farid K, Khan I & Tariq M (2009). Prevalence of Infectious bursal Disease in Broiler in District Peshawar. *Journal of Agricultural and Biological Science*. **4**: 1-5.
- Lasher HN & Davis VS (1997). History of infectious bursal disease in the U.S.A- the first two decades. *Avian Disease*, **41**(1): 11-19.
- Lawal JR, Jajere SM, Bello AM, Mustapha M, Wakil Y, Ndahi JJ, Mustapha FB, Paul BT, Gulani IA, Ibrahim UI, Geidam YA, Ambali AG & Waziri I (2014). Prevalence of Infectious Bursal Disease (Gumboro) Antibodies in Village Chickens in Gombe State, Northeastern Nigeria. *International Journal of Poultry Science*, **13** (12): 703-708.
- Lukert PD & Saif YM (1997). Infectious bursal disease. In: Diseases of poultry, Calnek BW, Barnes HJ, Beard CW, McDougald LR and Saif YM. (Eds) 10th Edition Iowa State University Press. Pp 721-735.
- Lukert PD & Saif, YM (2003). Infectious Bursal Disease Virus. In: Diseases of Poultry (YM Saif, HJ Barnes, JR Glison, LR McDougald, DE Swayne, editors). Eleventh edition. Ames, 10: Iowa State University Press. Pp 161-179.
- Mai HM, Ogunsola OD & Obasi OL (2004). Serological survey of the Newcastle disease and infectious bursal disease in local ducks and local guinea fowls in Jos, Plateau state, Nigeria. *Revue d'élevage et de médecine vétérinaire des pays tropicaux*, **57**(1-2): 41-44.
- Mbuko IJ, Musa WI, Ibrahim S, Sa'idu L, Abdu PA, Oladele SB & Kazeem HM (2010). A Retrospective Analysis of Infectious Bursal Disease Diagnosed at Poultry Unit of Ahmadu Bello University, Nigeria. *International Journal of Poultry Science*, **9**(8): 784-790.
- McFerran JB, McNulty MS, McKillop ER, Connor TJ, McCracken RM, Collins PS & Allan GM (1980). Isolation and serological studies of infectious bursal disease virus from fowl, turkey and ducks: Demonstration of a second serotype. *Avian Pathology*, **9**(3): 395-404.

- Oladele OA, Emikpe BO, Oluwayelu OD & Ohore OG (2004). Comparison of Agar gel precipitation test (AGPT) and Enzyme linked immunosorbent assay (ELISA) in the detection of infectious bursal disease virus (IBDV) antibody in village chickens in Oyo State, Nigeria, *Nigerian Veterinary Journal*, **25**(1): 26-29.
- Oluwayelu DO (2010). Diagnosis and Epidemiology of chicken infectious anemia in Africa. *African Journal Biotechnology*, **9**(14): 2043-2049.
- Oluwayelu DO, Emikpe BO, Oladele OA, Ohore OG & Fagbohun OA (2007). Seroprevalence of Infectious Bursal Disease in flocks of indigenous Nigerian ducks (*Anas platyrhynchos*). *Journal of Animal and Veterinary Advances*, **6**(1): 64-67.
- Pankhurst RT (1964). Avian nephrosis (Gumboro disease) in USA broilers: treatment trials. *World's Poultry Science Journal*, **20**: 208-211.
- Tanimura N & Sharma JM (1997). Appearance of T cell in the bursa of Fabricius and cecal tonsils during the acute phase of infectious bursal disease virus infection in chickens. *Avian Disease*, **41**(3): 638-45.
- Waziri M, Kagu A & Monguno AK (2009). Issues in the Geography of Borno State, Published by Adamu Joji publishers No. 123 Mangwarori Street Sabon Titin Mandawari, Kano city, Kano State Nigeria. Pp 9.