



Review Manuscript

An Overview of Maternity healthcare Monitoring in Developing Nations

Godphrey G. Kyambille^{1†}, Aloys N. Mvuma¹ and Dina Machuve²

¹Mbeya University of Science and Technology

²The Nelson Mandela African Institute of Science and Technology

[†]Correspondence author: gkyambille@gmail.com

[†]ORCID: <https://orcid.org/0009-0002-5169-1238>

ABSTRACT

A healthcare monitoring arrangement is essential for frequently monitoring a patient's health status. Specifically, maternal healthcare systems for tracking are utilized to evaluate the clinical status and monitor any abnormal condition changes during all three trimesters. This paper's objective is to conduct an extensive literature review and acknowledge earlier initiatives and studies conducted previously in maternal health care monitoring. This review focuses on accumulating information from earlier work and presents a general overview of previous studies concerning maternal health care monitoring (MHCM). The paper focuses on the maternal healthcare systems in developing countries accessed by pregnant women during the antenatal period. Furthermore, this paper highlights the challenges in the maternal health sector that include inadequate work conditions for health workers, home-based birth practices, long distances to healthcare facilities, and long waiting times in health facilities. The paper indicates the potential opportunities to strengthen maternal healthcare services and allow effective access to maternal healthcare facilities by pregnant women.

ARTICLE INFO

Submitted: **Jan. 10, 2023**

Revised: **Dec. 27, 2023**

Accepted: **Jan. 15, 2024**

Published: **Feb., 2024**

Keywords: *Maternity, Health Services, Hospitalization, patients, prenatal care, Care during pregnant*

INTRODUCTION

The field of health is one of the most crucial fields for saving lives in any nation. Worldwide most hospitals strive to improve their services, including treatment service, delivery systems and environment, to meet patients' expectations. As part of the service improvement initiatives, computers create and maintain access to patients' medical records. In the healthcare field, the World Health Organization (WHO) endorse access to

services for maternal healthcare, family planning and delivery services (Organization, 2005). or. Yet, to target efficient MHCM, important areas need appropriate management and quality evaluation. For example, the time it takes pregnant women to walk into a hospital and attend is the most critical factor that needs proper handling (Shija *et al.*, 2011). In Tanzania, MHCM is conducted manually, requiring pregnant women to visit the health centre monthly physically. The pregnancy health status is recorded

manually in clinic cards and scheduled for the next visit. Furthermore, it is reported that slots of time available for MHCM are not unionized in the way pregnant women can reschedule some other day if missed a clinical appointment checkup or need an urgent clinical appointment.

Moreover, when pregnant women visit the hospital, they are not assured of being seen by doctors when they are in bench queues. On a day, those doctors attend a massive number of pregnant women, as the designation is based on a day, not a time slot. When pregnant women postpone their clinical appointment, there is no mechanism for doctors to notice that patients won't show up for clinical maternal checkups.

Monitoring maternal health care was selected since it encourages distinguishing proof of information gaps and the chance that exists due to emerging technology (Mtei *et al.*, 2014). It is important to follow up on maternal health (Pembe *et al.*, 2010). Currently, the methodology used in maternal healthcare monitoring is manual, allowing pregnant women to attend healthcare facilities physically. Maternal healthcare monitoring for pregnancy is conducted on monthly visits to a health facility until the last trimester, where maternal monitoring is on weekly basis. During the first maternal visit, a pregnant woman is required to fill in a personal clinic card with her personal details.

This literature explores the information about diverse technology mechanisms for monitoring maternal health found from several sources. This paper discusses the finding based on 345 reviewed articles from three databases; Web of Science, PubMed/Medline, WHO Global Health Library, Popline and Google Scholar. The literature review is focused on maternal health care monitoring in Tanzania. The information collected from this literature review will be applied to design a technological solution for MHCM.

The technological solution can help maternal health care improve (Bhat *et al.*,

2011). It will promote better health sector services to patients and doctors by knowing when to attend to the patient and who is responsible at what time. (Mey & Sankaranarayanan, 2013) Show that service waiting time will decrease if technology is applied to health services.

Therefore, the challenge of MHCM application of Information and Communication Technology (ICT) is inevitable. ICT is used in symptom diagnosis and offers efficiency in obtaining results quicker. ICT presently has made tremendous innovation and become accessible both in organizations and healthcare industries.

METHODS AND MATERIALS

This segment briefly clarifies the method followed for identifying related and suitable literature articles for the research work. The first part explains the aim of this literature review; the second part narrates the search process for the literature review, including database selection and search query. The final part defines the study selection procedure.

Goal of the Research

Examining the available research is the goal of the literature review, acknowledging earlier research and researching maternal health care monitoring in underdeveloped nations. The literature review focus on compiling the information obtained from previous work and presents a comprehensive overview of information attained from previous studies concerning MHCM.

Search Process

A literature review was steered in four databases. The literature search was conducted on Web of Science, WHO Global Health Library, PubMed/Medline, Popline and Google Scholar for peer-reviewed articles published up to June 31, 2022, using search terms as shown in Table 1. The search was restricted to review

papers focusing only on monitoring maternal health to meet the criteria. Review articles concentrating on manual

monitoring systems were also included to comprehend the current monitoring of maternal health care services.

Table 1. Searched terms

SN	KEYWORDS	SEARCHED ITEM
1	Maternal care Health	Monitoring, Machine Learning, ICT, Patient Monitoring, Medical Sensor, Internet of Things, Remote Monitoring, Artificial Intelligence, Wearable tech Antenatal care, Prenatal care
2	Developing countries	Sub-Saharan countries, middle-income countries, lower-income countries

RESULTS AND DISCUSSION

Data search process

After literature searches in the databases, reviewed articles were further extracted into their title and abstract if they were related, and related articles were fully reviewed. Figure 2 represents the data search process. A total of 345 reviewed articles in all databases were screened. After screening for title-related, 125 articles were eliminated, 220 articles were

evaluated based on their abstract-related, 98 articles were eliminated, and 77 articles could not be accessed. The final identified reviewed articles that met the criteria were 77. Key issues recognized from the reviewed articles were the information on the sensors used in the devices for capturing data and monitoring symptoms and the analyzed data was utilized to predict by machine learning techniques.




Figure 2: Data search process.

Description of sensors used in the devices for capturing data

The literature review revealed different types of sensors used to find symptoms or health data for numerous diseases. These sensors include heart rate, temperature,

ECG, EEG, gyroscopes, and accelerometers. Information obtained from (Site *et al.*, 2021) accelerometer sensors have been used in more than 23% of studies, gyroscopes sensors have been used in more than 10% of studies, Magnetometer

sensors have been used in more than 5% of studies, Blood glucose meter, Gait sensors have been used more than 3% of the studies, moreover pulse oximeter, temperature sensor, heart rate sensor, humidity sensor and pacemakers have been used more than 1% of the studies. Furthermore, the reviewed article proves that even smartwatches/wearables devices with built-in sensors have been used in more than 14% of the studies.

Zhao *et al.*, 2019 presented an accessory system for automatically tracking of speedometers and methods for learning from machines to monitor activities of the fetus.. Ullah *et al.*, 2021 propose a system for recognizing and tracking physical activity that employs embedded detectors to detect and monitor a vigorous workout throughout pregnancy. The effects of exercising during pregnancy can differ based on the level of exertion, which might whether exacerbate or alleviate health issues.

Musyoka et al., 2019, implemented an ambulating cardiovascular health for an entire day. This system comprises a timepiece featuring hosted on the internet and smartphone functions applications. When the expectant mother's blood pressure is measured, a notification is delivered to the designated caregiver to take immediate action.

Ullah *et al.*, 2021, In the context of pregnancy, provide a wearable sensor-based system for recognizing and tracking physical activity. A body-worn component, as per the advised structure, consists of sensors for temperature, an accelerometer with three axes and a three-axis gyroscope. Raspberry PI historical information is transmitted via Bluetooth Low Energy (BLE) sensors.

Roham *et al.*, 2011 suggest a wireless, portable technology for non-invasively monitoring uterine contractions and fetal heart rate at the same time. A mobile cellular gateway for wide-area connectivity makes up the entire system, a wearable front-end for Doppler ultrasound and

pressure sensors with short-range radio, for remote monitoring and diagnostics, a web server and browser-based user interface are required.

They provide a vital mobile a multi-communication fusion and Android OS-based indications monitoring system that enables expectant mothers to track fetal and maternal data from any location. Moreover, by sending information to the server for processing and remote diagnostics, patients have access to comprehensive care. (Lyu et al., 2013)

Tsai *et al.*, 2006 proposes a tracking the fetal heart rate (FHR) and prenatal recognition of distress portable equipment. The suggested method uses machine vision techniques to gather FHR values and the use of pattern matching to analyze them. once a mobile GSM network detects fetal distress, a medical alarm will instantly alert medical professionals. A pregnant woman can monitor her baby's health state using the suggested approach, and medical professionals can promptly provide resources for treatment.

Queyam *et al.*, 2018, Present a reliable bio tele monitoring of pregnant women using an intra-probe, multiple-parameter system to enable comprehensive the collection of personally identifiable information to track their health throughout pregnancy and labour. Zhang *et al.*, 2018, suggest home-based fetal heartbeat monitoring and identification throughout pregnancy.

(Subalakshmi *et al.*, 2018) Outlines a framework for monitoring physical variations in pregnant women. This framework can spot unforeseen events and alert the patient and doctor. The controller node (CN) is physically connected to the patient in the proposed design to collect all uterine contraction signals, and Monitoring devices sensors are tracked and sent to the access point. A network of wireless body sensors (WBSN) has sensors linked to the patient's body frame that may assess things like blood pressure, pulse rate, and uterine tightness.

Allahem & Sampalli, 2017, Propose a framework for employing sensor networks to monitor pregnant women at risk for early labour. The proposed approach tracks uterine contractions non-invasively using body-worn wireless sensor network and notifies a smartphone if the results are above or below the normal limit.

Chourasia & Tiwari, 2012 Propose a fetal tracker for electronic health utilizing smart mobile devices and wearable network to deliver cutting-edge healthcare in a home setting.

Nitulescu *et al.*, 2015, Presents a framework for employing sensor networks to monitor pregnant women at risk for early labour. The suggested method uses a body-worn wireless sensor network to track uterine contractions non-invasively and alert a smartphone if the readings are above or below typical limits.

Aravind *et al.*, 2017 provided a health monitoring system powered by the Internet of Things (IoT) that continuously detects body pressure with the use of a pressure measurement kit and measures a pregnant woman's heartbeat using a heartbeat sensor. Every time there are variations from the specific number, it sends the information to her doctor through a mobile.

Predictive algorithms using machine learning

Numerous automated learning techniques have been used to predict maternal health status. Supervised and unsupervised learning algorithms are examples of classification of machine learning algorithms. In reviewed articles, both supervising and unstructured, both of which have been employed for various arrangement and regression tasks.

According to Site *et al.*, 2021, a supervised machine-learning algorithm, which supports vector machines, has been used in more than 43 studies to analyze various features. Neural networking has been used 41 times to analyze information extracted from different features.

Mehbodniya *et al.*, 2022 used machine learning (ML) techniques that classify the medical condition as normal, needs guarantee, or pathology to forecast the maternal wellbeing from the cardiocotographic (CTG) information. Using algorithms like support vector machines and random forests, this study determines how several CTG-measured parameters affect the forecasting of the fetus's health. Yakong *et al.*, 2010, K-nearest neighbours and multi-layer perceptron. Furthermore, regression analysis Additionally, the correlation and regression analyses showed how the variables affected fetal health.

Marin *et al.*, 2019 employed a machine learning strategy for early preeclampsia identification and the Viterbi algorithm to identify preeclampsia.

Sharma & Sharma, 2022 suggested employing a novel upgraded binary bat algorithm to assess the health of a fetus. The suggested EBBA can be utilized to classify cardiocotography datasets into normal, suspicious, and pathologic fetal states when using the random forest classification algorithm from machine learning, and it has an efficiency of 96.21%.

Ahmed & Kashem, 2020, employed a modified decision tree algorithm for classification and risk level prediction to create great monitoring and insight into the risk a pregnant woman faces. (Hoffman *et al.*, 2021) propose the development of machine learning and validate in the context of neonatal hospitalization of mothers attributed explicitly to difficulties arising from pregnancy-related hypertension issues.

Challenges of maternal healthcare service in developing countries

Maternal healthcare services have been fundamental objective of The WHO introduced the Millennium Development Objectives (MDGs) and the

Sustainable Development Goals (SDGs) main concerns, (Sachs, 2012).

The difficult working condition of health workers, especially in rural areas, affects their main responsibility of providing good care services for maternal health. Bureaucracy facilitates the unavailability of a specific strategy for health workers' development to enhance the expertise of health professionals in maternal health care issues (Mkoka *et al.*, 2015).

Home-based neonatal care practised by many societies facilitates the challenge of attending maternal healthcare monitoring during pregnancy. Research conducted in the Lindi region reveals that some society allows pregnant women to move to their parents during their last trimester to deliver (Mrisho *et al.*, 2008).

Long distance to maternal health facilities hinders entry to maternal services for health care, particularly in remote locations. Some pregnant women attend only the final trimester due to the distance to a health facility. According to various research, armed conflicts make maternity healthcare services less accessible and raise the rate of maternal mortality (Chukwuma & Ekhat-Mobayode, 2019). Long waiting time to be attended at a maternal health facility is a big challenge for pregnant women. Pregnant women are expected not to consume more time attending the maternal health facility. Inadequate scheduling during maternal healthcare visiting. Normally, scheduling is provided in a range of groups of time and not a slot of time for individual pregnant women.

Some pregnant women are unsatisfied with maternal healthcare services when attending clinical checkups, hence attending maternal healthcare monitoring only during the last trimesters and the first two trimesters remained unmonitored. Only pregnant women with obstetric and required emergency care are attending all three trimesters for maternal healthcare monitoring.

Opportunities to improve maternal healthcare services

Primary healthcare providers are struggling to quicken improvement towards attaining Millennium Development goals on how easily pregnant women can access maternal healthcare services during pregnancy without any difficulties (Basinga *et al.*, 2011). Maternal healthcare service in Tanzania is free in all public health facilities, including antenatal, delivery, and family planning (Magoma *et al.*, 2010; Mpembeni *et al.*, 2007). Even pregnant women from low families access maternal healthcare monitoring for all trimesters for free in any public facility. The presence of private health institutions, which annually produce health professionals, alleviates the scarcity of healthcare personnel in the commercial and public sectors, particularly in maternal healthcare (White *et al.*, 2013). Currently, the Tanzanian government has built health facilities with emergency care in practically every district to facilitate frequent visits and access to maternal healthcare services for expectant women often then get maternal healthcare services. It makes it easier for people to travel large distances to medical facilities, especially in rural areas where pregnant women with obstetric difficulties must receive specialized care in urban healthcare centres, (Ramsey *et al.*, 2013).

Men's participation in reproductive health is inevitable (Amoo *et al.*, 2017). Suppose men are well involved during the pregnancy period. In that case, all complications that pregnant women are expected to face can be reduced as knowledge of reproductive are on both sides of the community (Vermeulen *et al.*, 2016). Men are regarded as financial backers and those who make decisions about when and where women can get maternal health care and services. Pregnancy-related difficulties are seen as a woman's responsibility throughout all trimesters (August *et al.*, 2016). Participation of community people in

reproductive health will make maternal health services more easily accessible (Kilonzo *et al.*, 2001).

Discussion

The literature review examined the process of monitoring maternal health care in developing countries using 77 studies attained over a succession of stages. This study revealed different types of sensors that can be used to obtain health data. This review discovered that different signals could be collected from sensors, including gyroscopes, EEG/ECG, and wearables like smartwatches. The study also identified several traits that can be drawn from those signals.

The article investigates how more studies have utilized supervised and unsupervised machine learning methods for data analysis. The greatest studies use machine learning algorithms to achieve classification and prediction tasks. The study employed multiple machine learning techniques to analyze those characteristics. Several different kinds of tree-based and neural networking algorithms were used to analyze feature vectors. The outcomes attained from this literature review- and the information and knowledge gained from earlier studies will be used for emerging computerized real-time systems monitoring maternal health care. Furthermore, the review was more attentive to features like wearable/sensors essential in designing real-time monitoring systems for maternal health care.

CONCLUSION AND RECOMMENDATION

A review of previously published research revealed a critical need to enhance maternal healthcare monitoring. Some obstacles make it more difficult for pregnant women to get maternal health treatments, such as when they are women with impairments who lack adequate transportation (Ganle *et al.*, 2016). Most of them are expected to have lower access to education and social

and economic issues than those without disabilities (Hosseinpoor *et al.*, 2013). Moreover, during the covid 19, there were movement restrictions and accessing maternal health services was difficult and very challenging (Nguyen *et al.*, 2022; Pant *et al.*, 2020)

All women should have equal access to maternal health services, yet there are differences between rural and urban women in terms of attending and using these services (Yakong *et al.*, 2010). Other articles advise that maternal health services need to be increased in areas where maternal services are limited and accessibility is difficult (Elmusharaf *et al.*, 2015). When better services are provided, the focus is on how satisfied women are with their capacity to access maternity care. (Srivastava *et al.*, 2015). The attitude and behaviour of caregivers during pregnancy can be taken concerning women when the accessibility of maternal care services is conducted smoothly and appreciated by women who attend the services (Ajayi, 2019). It can be used to evaluate how satisfied pregnant women are with the treatment they receive from maternal health care providers. (Camacho *et al.*, 2012); in turn, will assist hospital administrators in stepping up efforts to provide maternal healthcare services, especially to pregnant women who need special care throughout pregnancy (Özkan *et al.*, 2020). The continuum of care enables women and maternal health care professionals to interact more effectively. (Macpherson *et al.*, 2016).

REFERENCES

- Ahmed, M., & Kashem, M. A. (2020). IoT based risk level prediction model for maternal health care in the context of Bangladesh. *2020 2nd International Conference on Sustainable Technologies for Industry 4.0 (STI)*, doi:[10.1109/STI50764.2020.9350320](https://doi.org/10.1109/STI50764.2020.9350320)
- Ajayi, A. I. (2019). "I am alive; my baby is alive": Understanding reasons for satisfaction and dissatisfaction with maternal health care services in the context of user fee removal

- policy in Nigeria. *PloS one*, **14**(12), e0227010. doi: 10.1371/journal.pone.0227010
- Allahem, H., & Sampalli, S. (2017). Framework to monitor pregnant women with a high risk of premature labour using sensor networks. *2017 IFIP/IEEE Symposium on Integrated Network and Service Management (IM)*, doi: [10.23919/INM.2017.7987458](https://doi.org/10.23919/INM.2017.7987458)
- Amoo, E. O., Oni, G. A., Ajayi, M. P., Idowu, A. E., Fadayomi, T. O., & Omideyi, A. K. (2017). Are men's reproductive health problems and sexual behavior predictors of welfare? *American Journal of Men's Health*, **11**(3), 487-497. doi: [10.1177/15579883155988](https://doi.org/10.1177/15579883155988)
- Aravind, P., Kumar, U. N., Sivakumar, V., & Karthikeyan, S. (2017). Pregnancy health care monitoring system. *International Journal on Information Sciences and Computing*, **11**(1).
- August, F., Pembe, A. B., Mpembeni, R., Axemo, P., & Darj, E. (2016). Community health workers can improve male involvement in maternal health: evidence from rural Tanzania. *Global Health Action*, **9**(1), 30064. doi: [10.3402/gha.v9.30064](https://doi.org/10.3402/gha.v9.30064)
- Basinga, P., Gertler, P. J., Binagwaho, A., Soucat, A. L., Sturdy, J., & Vermeersch, C. M. (2011). Effect on maternal and child health services in Rwanda of payment to primary healthcare providers for performance: an impact evaluation. *The lancet*, **377**(9775), 1421-1428. doi: [10.1016/S0140-6736\(11\)60177-3](https://doi.org/10.1016/S0140-6736(11)60177-3)
- Bhat, S., Sidnal, N. S., Malashetty, R. S., & Manvi, S. S. (2011). Intelligent Scheduling in Health Care Domain. *International Journal of Computer Science Issues (IJCSI)*, **8**(5).
- Camacho, F. T., Weisman, C. S., Anderson, R. T., Hillemeier, M. M., Schaefer, E. W., & Paul, I. M. (2012). Development and validation of a scale measuring satisfaction with maternal and newborn health care following childbirth. *Maternal and Child Health journal*, **16**(5), 997-1007. doi: 10.1007/s10995-011-0823-8
- Chourasia, V. S., & Tiwari, A. K. (2012). Implementation of foetal e-health monitoring system through biotelemetry. *International Journal of Electronic Healthcare*, **7**(1), 36-52. doi: [10.1504/IJEH.2012.048668](https://doi.org/10.1504/IJEH.2012.048668)
- Chukwuma, A., & Ekhatior-Mobayode, U. E. (2019). Armed conflict and maternal health care utilization: evidence from the Boko Haram Insurgency in Nigeria. *Social Science & Medicine*, **226**, 104-112. doi: [10.1016/j.socscimed.2019.02.055](https://doi.org/10.1016/j.socscimed.2019.02.055)
- Elmusharaf, K., Byrne, E., & O'Donovan, D. (2015). Strategies to increase demand for maternal health services in resource-limited settings: challenges to be addressed. *BMC Public Health*, **15**(1), 1-10. doi: 10.1186/s12889-015-2222-3
- Ganle, J. K., Otupiri, E., Obeng, B., Edusie, A. K., Ankomah, A., & Adanu, R. (2016). Challenges women with disability face in accessing and using maternal healthcare services in Ghana: a qualitative study. *PloS one*, **11**(6), e0158361. doi: [10.1371/journal.pone.0158361](https://doi.org/10.1371/journal.pone.0158361)
- Hoffman, M. K., Ma, N., & Roberts, A. (2021). A machine learning algorithm for predicting maternal readmission for hypertensive disorders of pregnancy. *American Journal of Obstetrics & Gynecology MFM*, **3**(1), 100250. doi: [10.1016/j.ajogmf.2020.100250](https://doi.org/10.1016/j.ajogmf.2020.100250)
- Hosseinpour, A. R., Stewart Williams, J. A., Gautam, J., Posarac, A., Officer, A., Verdes, E., . . . Chatterji, S. (2013). Socioeconomic inequality in disability among adults: a multicountry study using the World Health Survey. *American Journal of Public Health*, **103**(7), 1278-1286. doi: [10.2105/AJPH.2012.301115](https://doi.org/10.2105/AJPH.2012.301115)
- Kilonzo, A., Kouletio, M., Whitehead, S. J., Curtis, K. M., & McCarthy, B. J. (2001). Improving surveillance for maternal and perinatal health in 2 districts of rural Tanzania. *American Journal of Public Health*, **91**(10), 1636-1640. doi: [10.2105/AJPH.91.10.1636](https://doi.org/10.2105/AJPH.91.10.1636)
- Lyu, P., Peng, M., Lyu, Y., Chen, Y., & Yang, J. (2013). A multi-communication-fusion based mobile monitoring system for maternal and fetal information. *2013 IEEE 15th International Conference on e-Health Networking, Applications and Services (Healthcom 2013)*, doi: [10.1109/HealthCom.2013.6720739](https://doi.org/10.1109/HealthCom.2013.6720739)
- Macpherson, I., Roqué-Sánchez, M. V., Legget, F. O., Fuertes, F., & Segarra, I. (2016). A systematic review of the relationship factor between women and health professionals within the multivariate analysis of maternal satisfaction. *Midwifery*, **41**, 68-78. doi: [10.1016/j.midw.2016.08.003](https://doi.org/10.1016/j.midw.2016.08.003)
- Magoma, M., Requejo, J., Campbell, O. M., Cousens, S., & Filippi, V. (2010). High ANC coverage and low skilled attendance in a rural Tanzanian district: a case for implementing a birth plan intervention. *BMC Pregnancy and Childbirth*, **10**(1), 1-12. doi: 10.1186/1471-2393-10-13
- Marin, I., Pavaloiu, B.-I., Marian, C.-V., Racovita, V., & Goga, N. (2019). Early detection of preeclampsia based on a machine learning

- approach. 2019 E-Health and Bioengineering Conference (EHB), doi: [10.1109/EHB47216.2019.8970025](https://doi.org/10.1109/EHB47216.2019.8970025)
- Mehbodniya, A., Lazar, A. J. P., Webber, J., Sharma, D. K., Jayagopalan, S., Singh, P., . . . Sengan, S. (2022). Fetal health classification from cardiotocographic data using machine learning. *Expert Systems*, **39**(6), e12899. doi: [10.1111/exsy.12899](https://doi.org/10.1111/exsy.12899)
- Mey, Y. S., & Sankaranarayanan, S. (2013). Near field communication based patient appointment. *International Conference on Cloud & Ubiquitous Computing & Emerging Technologies (CUBE)*, 2013. doi: [10.1109/CUBE.2013.27](https://doi.org/10.1109/CUBE.2013.27)
- Mkoka, D. A., Mahiti, G. R., Kiwara, A., Mwangu, M., Goicolea, I., & Hurtig, A.-K. (2015). "Once the government employs you, it forgets you": Health workers' and managers' perspectives on factors influencing working conditions for provision of maternal health care services in a rural district of Tanzania. *Human Resources for Health*, **13**(1), 1-13. DOI doi: [10.1186/s12960-015-0076-5](https://doi.org/10.1186/s12960-015-0076-5)
- Mpembeni, R. N., Killewo, J. Z., Leshabari, M. T., Massawe, S. N., Jahn, A., Mushi, D., & Mwakipa, H. (2007). Use pattern of maternal health services and determinants of skilled care during delivery in Southern Tanzania: implications for achievement of MDG-5 targets. *BMC Pregnancy and Childbirth*, **7**(1), 1-7. DOI doi: [10.1186/1471-2393-7-29](https://doi.org/10.1186/1471-2393-7-29)
- Mrisho, M., Schellenberg, J. A., Mushi, A. K., Obrist, B., Mshinda, H., Tanner, M., & Schellenberg, D. (2008). Understanding home-based neonatal care practice in rural southern Tanzania. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **102**(7), 669-678. doi: [10.1016/j.trstmh.2008.04.029](https://doi.org/10.1016/j.trstmh.2008.04.029)
- Mtei, G., Makawia, S., & Masanja, H. (2014). Monitoring and evaluating progress towards universal health coverage in Tanzania. *PLoS Medicine*, **11**(9), e1001698. doi: [10.1371/journal.pmed.1001698](https://doi.org/10.1371/journal.pmed.1001698)
- Musyoka, F. M., Thiga, M. M., & Muketha, G. M. (2019). A 24-hour ambulatory blood pressure monitoring system for preeclampsia management in antenatal care. *Informatics in Medicine Unlocked*, **16**, 100199. doi: [10.1016/j.imu.2019.100199](https://doi.org/10.1016/j.imu.2019.100199)
- Nguyen, T. V., King, J., Edwards, N., & Dunne, M. P. (2022). "Nothing suitable for us": experiences of women with physical disabilities in accessing maternal healthcare services in Northern Vietnam. *Disability and Rehabilitation*, **44**(4), 573-581. doi: [10.1080/09638288.2020.1773548](https://doi.org/10.1080/09638288.2020.1773548)
- Nitulescu, A., Crisan-Vida, M., Stoicu-Tivadar, L., & Bernad, E. (2015). Integrated wireless sensor network for monitoring pregnant women. *MIE*, doi: [10.3233/978-1-61499-512-8-354](https://doi.org/10.3233/978-1-61499-512-8-354)
- Organization, W. H. (2005). *The World health report: 2005: make every mother and child count*. World Health Organization.
- Özkan, Ş., Chiang, C., Aba, G., & Çelik, Y. (2020). Satisfaction with maternal and birth services: a survey in public hospitals in Turkey. *International Journal of Health Care Quality Assurance*. doi: [10.1108/IJHCQA-08-2019-0146](https://doi.org/10.1108/IJHCQA-08-2019-0146)
- Pant, S., Koirala, S., & Subedi, M. (2020). Access to maternal health services during COVID-19. *Europasian Journal of Medical Sciences*, **2**(2), 46-50. doi: [10.46405/ejms.v2i2.110](https://doi.org/10.46405/ejms.v2i2.110)
- Pembe, A. B., Carlstedt, A., Urassa, D. P., Lindmark, G., Nyström, L., & Darj, E. (2010). Effectiveness of maternal referral system in a rural setting: a case study from Rufiji district, Tanzania. *BMC Health Services Research*, **10**(1), 1-9. doi: [10.1186/1472-6963-10-326](https://doi.org/10.1186/1472-6963-10-326)
- Queyam, A. B., Meena, R. K., Pahuja, S. K., & Singh, D. (2018). An IoT based multi-parameter data acquisition system for efficient bio-telemonitoring of pregnant women at home. *2018 8th International Conference on Cloud Computing, Data Science & Engineering (Confluence)*, doi: [10.1109/CONFLUENCE.2018.8442686](https://doi.org/10.1109/CONFLUENCE.2018.8442686)
- Ramsey, K., Hingora, A., Kante, M., Jackson, E., Exavery, A., Pemba, S., . . . Phillips, J. F. (2013). The Tanzania Connect Project: a cluster-randomized trial of the child survival impact of adding paid community health workers to an existing facility-focused health system. *BMC Health Services Research*, **13**(2), 1-14. doi: [10.1186/1472-6963-13-S2-S6](https://doi.org/10.1186/1472-6963-13-S2-S6)
- Roham, M., Saldivar, E., Raghavan, S., Zurcher, M., Mack, J., & Mehregany, M. (2011). A mobile wearable wireless fetal heart monitoring system. *2011 5th International Symposium on Medical Information and Communication Technology*, doi: [10.1109/ISMICT.2011.5759813](https://doi.org/10.1109/ISMICT.2011.5759813)
- Sachs, J. D. (2012). From millennium development goals to sustainable development goals. *The lancet*, **379**(9832), 2206-2211. doi: [10.1016/S0140-6736\(12\)60685-0](https://doi.org/10.1016/S0140-6736(12)60685-0)
- Sharma, P., & Sharma, K. (2022). Fetal state health monitoring using novel Enhanced Binary

- Bat Algorithm. *Computers and Electrical Engineering*, **101**, 108035. doi: [10.1016/j.compeleceng.2022.108035](https://doi.org/10.1016/j.compeleceng.2022.108035)
- Shija, A. E., Msovela, J., & Mboera, L. E. (2011). Maternal health in fifty years of Tanzania independence: challenges and opportunities of reducing maternal mortality. *Tanzania Journal of Health Research*, **13**(5). doi: 10.4314/thrb.v13i5.5
- Site, A., Nurmi, J., & Lohan, E. S. (2021). Systematic review on machine-learning algorithms used in wearable-based eHealth data analysis. *IEEE Access*. doi: [10.1109/ACCESS.2021.3103268](https://doi.org/10.1109/ACCESS.2021.3103268)
- Srivastava, A., Avan, B. I., Rajbangshi, P., & Bhattacharyya, S. (2015). Determinants of women's satisfaction with maternal health care: a review of literature from developing countries. *BMC Pregnancy and Childbirth*, **15**(1), 1-12. doi: 10.1186/s12884-015-0525-0
- Subalakshmi, S., Murugaboopathi, G., & Senthilkumar, D. (2018). Efficient EER-LEACH protocol for monitoring the activities of pregnant women using wearable body sensor network. *Int. J. Eng. Technol*, **7**(3.27), 111-115.
- Tsai, C.-Y., Chiu, C.-C., & Chao, S.-M. (2006). A real-time mobile system for fetal heart rate monitoring and fetal distress detection. *7th International Conference on Mobile Data Management (MDM'06)*, doi: [10.1109/MDM.2006.14](https://doi.org/10.1109/MDM.2006.14)
- Ullah, F., Iqbal, A., Iqbal, S., Kwak, D., Anwar, H., Khan, A., . . . Kwak, K.-S. (2021). A framework for maternal physical activities and health monitoring using wearable sensors. *Sensors*, **21**(15), 4949. doi: [10.3390/s21154949](https://doi.org/10.3390/s21154949)
- Vermeulen, E., Solnes Miltenburg, A., Barras, J., Maselle, N., Van Elteren, M., & Van Roosmalen, J. (2016). Opportunities for male involvement during pregnancy in Magu district, rural Tanzania. *BMC Pregnancy and Childbirth*, **16**(1), 1-9. doi: 10.1186/s12884-016-0853-8
- White, J., O'Hanlon, B., Chee, G., Malangalila, E., Kimambo, A., Coarasa, J., . . . McKeon, K. (2013). *Private Health Sector Assessment in Tanzania*. World Bank Publications.
- Yakong, V. N., Rush, K. L., Bassett-Smith, J., Bottorff, J. L., & Robinson, C. (2010). Women's experiences of seeking reproductive health care in rural Ghana: challenges for maternal health service utilization. *Journal of Advanced Nursing*, **66**(11), 2431-2441. doi: [10.1111/j.1365-2648.2010.05404.x](https://doi.org/10.1111/j.1365-2648.2010.05404.x)
- Zhang, B., Lebedeva, I., Zhang, H., & Hu, J. (2018). Design for fetal heartbeat detection and monitoring in pregnancy care. *International Conference on Distributed, Ambient, and Pervasive Interactions*, DOI doi: 10.1007/978-3-319-91125-0_13
- Zhao, X., Zeng, X., Koehl, L., Tartare, G., de Jonckheere, J., & Song, K. (2019). An IoT-based wearable system using accelerometers and machine learning for fetal movement monitoring. *2019 IEEE International Conference on Industrial Cyber Physical Systems (ICPS)*, doi: [10.1109/ICPHYS.2019.8780301](https://doi.org/10.1109/ICPHYS.2019.8780301)