

Original Research Article

Effect of Diallylsulphide Supplementation on Wistar Rats Exposed to Mercuric Chloride

S Ansar and HT AIGHOSON

Clinical Laboratory Sciences, College of Applied Medical Science, King Saud University, Riyadh, Saudi Arabia

*For correspondence: **Email:** sansar@ksu.edu.sa; **Tel:** +966-118052970

Received: 8 September 2015

Revised accepted: 18 December 2015

Abstract

Purpose: To evaluate the protective potential of diallylsulphide (DAS) against mercury-induced oxidative stress and antioxidant enzymatic alterations in spleen of rats.

Methods: Rats were randomly divided into 4 groups of 6 rats each and exposed to mercuric chloride ($HgCl_2$) (50 mg/kg, i.p.) and/or diallylsulphide (200 mg/kg/b.w) by gavage. Oxidative stress was evaluated in spleen by antioxidant markers, viz, lipid peroxidation (LPO), superoxide dismutase (SOD), reduced glutathione (GSH), glutathione peroxidase (GPx) and catalase (CAT). Histomorphological changes in the spleen of rats were also compared between groups.

Results: Oral administration of DAS at a concentration of 50 mg/kg daily showed a significant increase in GSH and GPx ($p < 0.05$), SOD and CAT ($p < 0.05$), as well as decreased LPO ($p < 0.05$) level in the spleen of rats as compared with $HgCl_2$ treated rats. Histopathology of spleen also showed that administration of DAS reduced the damage generated by $HgCl_2$ treatment.

Conclusion: The results suggest that DAS may effectively normalize impaired antioxidant status in $HgCl_2$ -induced oxidative stress. DAS has a protective effect against lipid peroxidation by scavenging free radicals and is thus capable of ameliorating mercury-induced changes in the spleen of adult Wistar rats.

Keywords: Diallylsulphide, Metal toxicity, Antioxidants, Spleen, Mercury

Tropical Journal of Pharmaceutical Research is indexed by Science Citation Index (SciSearch), Scopus, International Pharmaceutical Abstract, Chemical Abstracts, Embase, Index Copernicus, EBSCO, African Index Medicus, JournalSeek, Journal Citation Reports/Science Edition, Directory of Open Access Journals (DOAJ), African Journal Online, Bioline International, Open-J-Gate and Pharmacy Abstracts

INTRODUCTION

Mercury toxicity has led to many health problems in the world. Mercury results in a variety of adverse neurological, renal, respiratory, immune, dermatological, reproductive and developmental disorders [1]. Its wide industry-related effects on humans have been well documented [2]. Mercuric chloride from industries, when released into water, causes morbidity and mortality in humans consuming sea food, including fish [3,4]. Exposure to mercury contamination could be from water, soil and air and can disturb the metabolic pathways [5]. It has been shown that kidneys contain the largest residues of mercury,

followed by liver, spleen, intestine, lymph node, skeletal muscles, lungs, heart, brain, and the intensity of the cytotoxic changes in the various organs was proportional to the amount of mercury accumulated [6].

The spleen is a secondary lymphoid organ present in all vertebrates in which the parenchyma is divided into white pulp (WP) and red pulp (RP) distinguished by different colors. The WP contains dense and highly organized accumulations of B and T lymphocytes around arterioles, while most of the RP consists of blood-filled spaces. These RP spaces are composed of two structures, the arterioles and

veins forming the splenic sinuses [7]. Earlier it has been reported that sub-chronic exposure to mercuric chloride resulted in a transient decrease of the lymphoid cell ratio which affected the incidence of splenic T-cell subsets [7-9].

The protective effects of garlic, and its constituents, have been attributed to presence of organosulphur compounds like diallyl sulphide (DAS), diallyl disulphide (DADS), ajoene, allixin, allyl mercaptans and allyl methylsulphides [10-12]. The DADS is one of the major volatile degradative compounds of garlic. Many studies on animals showed its protective effects against chemically induced toxicity and carcinogenesis [13,14].

Therefore, the present study was designed to study the antioxidant and protective effects of DAS against HgCl₂ toxicity.

EXPERIMENTAL

Animal groups

Male Wistar rats weighing approximately 180 - 200 g were used. All the animals used in this study were placed in cages in an air conditioned room maintained 12 h light/dark cycle. The study was approved by the Ethical Committee of King Saud University (CAMS 28-36/7) and was carried out in compliance with the declarations of National Research Council [15].

Experimental protocol

Thirty male Wistar rats were randomly divided into 5 groups (6 rats in each group). Group I received saline injection intraperitoneally (0.85 % NaCl) at a dose of 10 ml/kg body weight. Group II received a single intraperitoneal injection of mercuric chloride at a dose of 50 mg/kg bodyweight. Group III and IV received pretreatment with DAS i.p. once a day for 7 days at a dose of 200 mg/kg body weight. After the last treatment with DAS, the rats of group IV received a single intraperitoneal injection of HgCl₂ at a dose level of 50 mg/kg body weight. The single doses of DAS and mercury selected were based on a published dosage regimen that has been shown to be biologically and therapeutically active in experimental systems. After 24 h of last administration, the animals were sacrificed by cervical dislocation and the spleen tissue was harvested for biochemical and histopathological studies.

The spleen of rats were washed in ice-cold 50 mM Tris-HCl, pH 7.4 and homogenized in ice-cold medium that contained 50 mM Tris-HCl, pH

7.4. The homogenates were centrifuged at 3000 rpm for 10 min at 4 °C. The supernatants were used for the various biochemical determinations. The total protein content of the homogenized spleen was determined using bovine serum albumin as a standard [16].

Assay of oxidative stress markers

MDA content was assayed using the thiobarbituric acid (TBA) test as described by Ohkawa [17]. After incubation of homogenate with TBA at 95 °C, MDA reacts to form a pink colored complex. Absorbance was measured spectrophotometrically at 532 nm to determine the MDA content. The specific activity is expressed as nmol/mg protein.

For measurement of glutathione (GSH), an aliquot of 1.0 ml of renal PMS (10 % w/v) was precipitated with 1.0 ml of sulphosalicylic acid (4 % w/v). The samples were kept at 4 °C for at least 1 h and then subjected to centrifugation at 1200 g for 15 min at 4 °C. The assay mixture contained 0.1 ml filtered aliquot, 2.7 ml phosphate buffer (0.1 M, pH 7.4) and 0.2 ml DTNB (40 mg/10 ml of phosphate buffer 0.1 M, pH 7.4) in a total volume of 3.0 ml. The yellow color developed was read immediately at 412 nm on a spectrophotometer. The amount of glutathione was expressed as mmol/g tissue [18]. Total SOD activity was determined according to the method described by Marklund and Marklund [19] by assaying the autooxidation and illumination of pyrogallol at 440 nm for 3 min. One unit of SOD activity was calculated as the amount of protein that caused 50 % pyrogallol autooxidation inhibition. The SOD activity is expressed as IU/mg protein.

Before determination of the CAT activity, homogenates were diluted 1:9 with 1 % (v/v) Triton X-100. CAT activity was measured according to the method described by Aebi [20] by assaying the hydrolysis of H₂O₂ and the resulting decrease in absorbance at 240 nm over a 3 min period at 25 °C. CAT activity is expressed as IU/mg protein.

GPx activity was measured using H₂O₂ as substrate according to the method described by Paglia and Valentine [21]. The reaction was monitored indirectly as the oxidation rate of NADPH at 240 nm for 3 min. Enzyme activity was expressed as U/mg protein.

Histological examinations

Spleen tissues from rats were fixed in 10 % formaldehyde. Fixed tissues were dehydrated in

serial ethanol series, trimmed, embedded in paraffin, sectioned into 2- μ m sections and stained with hematoxylin and eosin (H&E). Morphological examination was conducted under a light microscope (Nikon Eclipse E600).

Statistical analysis

The results are expressed as mean \pm standard error of the mean (SEM). One-way ANOVA was applied to test for the significance of the biochemical data for the different groups. Significance level was set at $p < 0.05$.

RESULTS

HgCl₂ administration caused a significant ($p < 0.05$) increase in the level of LPO in brain compared to the control group. DAS pretreated group caused a significant ($p < 0.05$) reduction in LPO levels increased by HgCl₂ administration as compared to control as shown in Fig. 1.

The HgCl₂-induced oxidative stress was indicated by a significant reduction ($p < 0.05$) in GSH contents of spleen tissue of HgCl₂-treated

rats in comparison with the control group. This reduction in GSH contents was attenuated by pretreatment with DAS as shown in Fig. 2.

HgCl₂ administration led to modulation of antioxidant enzymes relative to the control rats. It also led to decrease in SOD, CAT, and GPx activities in spleen homogenates significantly ($p < 0.05$) compared to the control (Table 1). On the other hand DAS pretreatment elevated the activities of SOD, CAT, and GPx significantly ($p < 0.05$) compared to the HgCl₂ group.

Histopathological examination of spleen showed changes in the rats administered with mercury chloride with or without DAS as shown in Fig. 3. The result from the control (A) showed normal spleen tissue showed normal arrangement of cells, white pulp (WP) and red pulp (RP). While the spleen of the rats in the HgCl₂ treated group (B) showed changes in the spleen characterized by swelling, and monocular infiltration. While DAS pretreated group (C) and group (D) showed normalcy to architecture of the spleen which are similar to the control group (A).




Fig 1: Effects of DAS on lipid peroxidation formation treated with mercuric chloride. Values are means \pm SEM. a Significant change at $p < 0.05$ with respect to the control group. b Significant change at $p < 0.05$ with respect to Hg group




Figure 2: Effects of DAS on glutathione depletion. Values are mean \pm SEM. (a) Significant change at $p < 0.05$ with respect to the control group. (b) Significant change at $p < 0.05$ with respect to Hg group

Table 1: Effect of DAS on Hg-induced antioxidant status of rat spleen

Parameter	EXPERIMENTAL GROUPS			
	Control (I)	Hg (II)	DAS (III)	DAS + HgCl ₂ (IV)
SOD(U/mg prot)	250.31 ± 25.21	198.31 ± 22.34*	245.12 ± 18.32	225.12 ± 23.24**
CAT (IU/mg prot)	0.28 ± 0.034	0.20 ± 0.022*	0.29 ± 0.031	0.25 ± 0.018**
GPx (U/mg prot)	0.32 ± 0.012	0.23 ± 0.019 *	0.30 ± 0.021	0.29 ± 0.342**

SOD = superoxide dismutase, CAT = catalase, GPx = glutathione peroxidase; ***p* < 0.05 compared with control group; **p* < 0.05 compared with Hg-exposed group




Figure 3: Effect of DAS on spleen morphology: control (A) showing normal arrangement of cells, white pulp (WP) and red pulp (RP), HgCl₂ (B), DAS (C) DAS + HgCl₂ (D). Scale bar: 50 μ, H& E staining

DISCUSSION

The present study shows the ameliorative effects of DAS on the spleen of experimental rats treated with mercury chloride. It is well known that heavy metals are widely distributed in the environment and some of them can cause physiological, biochemical and histological disorders. The degree of toxic manifestation of different metals depend on dose, duration, route of administration and other physiological factors has been shown in many past studies [22-24].

The mercury usually deposits in the liver, kidneys, nerve cells, heart and other skeletal muscles, whereas in chronic condition its deposition is observed predominantly in hard tissues. In this study, DAS has been shown to ameliorate the toxic effects of mercury chloride on the biochemical parameters and plays important role in preventing oxidative damage to the spleen. Histological examination revealed that in the mercury treated group, spleen showed increased infiltration and vacuolization of the cells. Non treated spleen sections showed normal spleen structure composed of white pulp and red pulp, besides fibrous capsule which covered the spleen. White pulp consists mainly of B lymphocytes arranged into two zones, marginal zone (outer rim of loose lymphocytes

that contains macrophages and mantle zone (inner rim of lymphocytes).

In rats treated with mercuric chloride, there was significant decrease in the levels of GSH and SOD in spleen tissue. Glutathione, a tripeptide present in the majority of cells, is responsible for hydrophilic xenobiotics conjugation. GSH serves many vital physiological functions including protection of cells from reactive oxygen species (ROS), detoxification of exogenous compounds, and amino acid transport. Sulphydryl group of glutathione is essential for its antioxidant activity against some forms of ROS in cells. Much of the pathology is associated with the decrease in intracellular GSH concentration. Therefore, GSH concentration is important for survival of the cells. MDA is the most abundant individual aldehyde resulting from lipid peroxidation breakdown in biological systems and is commonly used as an indirect index of lipid peroxidation. In this study, lipid peroxidation levels increased by HgCl₂ administration as compared to control and DAS pretreated group caused a significant reduction as compared to mercury treated group.

The most important protective mechanism for free radical scavenging and superoxide dismutases (SODs) belongs to a family of antioxidant enzymes that catalyze the

dismutation of superoxide to yield hydrogen peroxide and oxygen. SOD is essentially a protective enzyme which scavenges the superoxide ions produced as cellular by-products during oxidative stress. Its decreased activity can lead to adverse effects because superoxide anions are extremely toxic and may accumulate in the cells.

Supplementation of DAS to the mercuric chloride-treated groups ameliorated malondialdehyde, glutathione and antioxidant enzymes activities. This could be explained by the major role of natural antioxidants in inhibiting lipid peroxidation and in protecting the wholeness and functioning of tissues and cells. Antioxidants may play an important role in abating some health hazards of heavy metals in connection with an interaction of physiological free radicals. It has also been known that garlic and its related compounds have inhibitory effects on chemical carcinogenesis and mutagenesis. These compounds may inhibit the activation of other carcinogens at low efficiency and the induction of glutathione -S-transferase and phase II enzymes may also play a role. Henceforth, it could be interpreted that the tissue protection against mercury may be due to the free radical scavenger effect of DAS.

CONCLUSION

The findings of this study suggest that mercuric chloride intoxication causes reactive oxygen species generation which in turn induces biochemical and histological alterations in rats. Administration of DAS proved to be beneficial in attenuating the mercuric chloride-induced oxidative toxicity.

ACKNOWLEDGEMENT

This work was supported by a grant from Research Center of the Center for Female Scientific and Medical Colleges', Deanship of Scientific Research, King Saud University, Riyadh, Saudi Arabia.

REFERENCES

- Risher JF, Amler SN. Mercury exposure. Evaluation and intervention the inappropriate use of chelating agents in the diagnosis and treatment of putative mercury poisoning. *Neurotoxicol* 2005; 26(4): 691-699.
- Kurl S, Tuomainen TP, Laukkanen JA, Nyyssonen K, Lakka T, Sivenius J, Salonen JT. Plasma vitamin C modifies the association between hypertension and risk of stroke. *J Cereb Circ* 2002; 33(6): 1568-1573.
- Son HY, Lee S, Park SB, Kim MS, Choi EJ, Singh TS, Bae Y, Kwack SJ, Kang TS, Shin HI et al. Toxic effects of mercuric sulfide on immune organs in mice. *Immunopharmacol Immunotoxicol* 2010; 32(2): 277-283.
- Magos L, Clarkson TW. Overview of the clinical toxicity of mercury. *Ann Clin Biochem* 2006; 43(4): 257-268.
- Bersenyi A, Fekete S, Hullar I, Kadar I, Szilagyi M, Glavits R, Kulcsar M, Mezes M, Zoldag L. Study of the soil-plant (carrot)-animal cycle of nutritive and hazardous minerals in a rabbit model. *Acta Vet Hung* 1999; 47(2): 181-190.
- Pathak SK, Bhowmik MK. The chronic toxicity of inorganic mercury in goats: clinical signs, toxicopathological changes and residual concentrations. *Vet Res Commun* 1998; 22(2): 131-138.
- Brunet S, Guertin F, Flipo D, Fournier M, Krzystyniak K. Cytometric profiles of bone marrow and spleen lymphoid cells after mercury exposure in mice. *Int J Immunopharmacol* 1993; 15(7): 811-819.
- Badano MN, Camicia GL, Lombardi G, Maglioco A, Cabrera G, Costa H, Meiss RP, Piazzon I, Nepomnaschy I. B-cell lymphopoiesis is regulated by cathepsin. *PloS one* 2013; 8(4): e61347.
- Verma S, Waldschmidt TJ. Characterization of splenic CD21hi T2 B cells. *Immunol Res* 2007; 39(1-3): 240-248.
- Shukla Y, Taneja P. Anticarcinogenic effect of black tea on pulmonary tumors in Swiss albino mice. *Cancer Lett* 2002; 176(2): 137-141.
- Shukla Y, Arora A, Taneja P. Antimutagenic potential of curcumin on chromosomal aberrations in Wistar rats. *Mutat Res* 2002; 515(1-2): 197-202.
- Shukla Y, Arora A, Singh A. Antitumorigenic potential of diallyl sulfide in Ehrlich ascites tumor bearing mice. *Biomed Environ Sci* 2002; 15(1): 41-47.
- Ansar S, Iqbal M, AlJameil N. Diallyl sulphide, a component of garlic, abrogates ferric nitrilotriacetate-induced oxidative stress and renal damage in rats. *Hum Exp Toxicol* 2014; 33(12): 1209-1216.
- Chen GW, Chung JG, Ho HC, Lin JG. Effects of the garlic compounds diallyl sulphide and diallyl disulphide on arylamine N-acetyltransferase activity in *Klebsiella pneumoniae*. *J. Appl. Toxicol* 1999; 19(2): 75-81.
- Derrell C. Guide for the care and use of laboratory animals. Institute of Laboratory Animal Resources. National Academy Press, Washington DC, USA 1996.
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the Folin phenol reagent. *J Biol Chem* 1951; 193(1): 265-275.
- Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal biochem* 1979; 95(2): 351-358.
- Jollow DJ, Mitchell JR, Zampaglione N, Gillete JR. Bromobenzene induced liver necrosis: protective role of glutathione and evidence for 3, 4 bromobenzeneoxide as the hepatotoxic intermediate. *Pharmacol* 1974; 11: 151-169.

19. Marklund, S., Marklund, G. Involvement of the superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase. *Eur J Biochem* 1974; 47: 469-474.
20. Aebi H. Catalase *in vitro*. *Methods Enzymol* 1984; 105: 121-126.
21. Paglia DE, Valentine WN. Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. *J Lab Clin Med* 1967; 70(1): 158-169.
22. Jomova K, Vondrakova D, Lawson M, Valko M. Metals, oxidative stress and neurodegenerative disorders. *Mol Cell Biochem* 2010; 345(1-2): 91-104.
23. Su L, Wang M, Yin ST, Wang HL, Chen L, Sun LG, Ruan DY. The interaction of selenium and mercury in the accumulations and oxidative stress of rat tissues. *Ecotoxicol Environ Saf* 2008; 70(3): 483-489.
24. Rao MV, Chhunchha B. Protective role of melatonin against the mercury induced oxidative stress in the rat thyroid. *Food Chem Toxicol* 2010; 48(1): 7-10.