

Original Research Article

Molecular detection of *Brucella melitensis* in sheep and goat milk in Iran

Amir Shakerian^{1,2*}, Permal Deo¹, Ebrahim Rahimi², Ali-Reza Shahjavan³ and Faham Khamesipour⁴

¹School of Pharmacy and Medical Sciences, University of South Australia, City East Campus, Adelaide SA, 5001, Australia, ²Department of Food Hygiene, Faculty of Veterinary Medicine, ³Faculty of Veterinary Medicine, Shahrekord Branch, Islamic Azad University, Shahrekord, ⁴Cellular and Molecular Research Center, Sabzevar University of Medical Sciences, Sabzevar, Iran

*For correspondence: **Email:** amshakerian@iaushk.ac.ir; amshakerian@yahoo.com; **Tel.:** +98-383 3361045

Received: 28 January 2016

Revised accepted: 12 April 2016

Abstract

Purpose: To detect *Brucella melitensis* in the milk of reared sheep and goats from Isfahan and Shahrekord regions, Iran.

Methods: A total of 225 milk samples (sheep = 125; goat = 100) were collected from Isfahan and Shahrekord regions, Central Iran. Polymerase chain reaction (PCR) was used to detect the presence of *B. melitensis* in the milk following standard procedures.

Results: From 225 milk samples, 20 (8.9 %) were positive for *B. melitensis*. Out of 125 sheep milk, 12 (9.6 %) had *B. melitensis*, and of these, 8 (66.6 %) were milk collected from Shahrekord and 4 (33.3 %) from Isfahan region. On the other hand, out of 100 goat milk samples, 18 (18 %) were positive for *B. melitensis*, out of which 10 (55.5 %) were from Shahrekord and 8 (44.4 %) from Isfahan.

Conclusion: The findings show that *B. melitensis* is present in a significant proportion of caprine and ovine milk in a section of Iran.

Keywords: *Brucella melitensis*, Milk, Polymerase chain reaction, Sheep, Goat

Tropical Journal of Pharmaceutical Research is indexed by Science Citation Index (SciSearch), Scopus, International Pharmaceutical Abstract, Chemical Abstracts, Embase, Index Copernicus, EBSCO, African Index Medicus, JournalSeek, Journal Citation Reports/Science Edition, Directory of Open Access Journals (DOAJ), African Journal Online, Bioline International, Open-J-Gate and Pharmacy Abstracts

INTRODUCTION

Brucellosis in sheep and goats is caused by *Brucella melitensis*, one of the most virulent species of *Brucella* [1-2]. Brucellosis is a widespread zoonosis, especially in the Mediterranean and Middle-East regions of the world where it constitutes occupational and public hazard [3-6]. *Brucella melitensis* is transmitted via ingestion of contaminated meat and milk and contact with contaminated individuals or fomites. Transmission of *B. melitensis* among caprine and ovine herds is rapid, and the disease being systemic in nature affects many organs and tissues [1,3,7]. Once

the acute period elapses, symptoms of brucellosis becomes non-pathognomonic, and the incriminating organism then becomes chronically located in both supramammary lymph nodes and mammary glands of 80 % of infected animals [8].

Interest have been aroused to keep brucellosis under control in endemic regions, because of the economic and health impact of brucellosis [9,10]. It is necessary to identify new endemic regions and to implement strict eradication programs beyond national borders. There is increasing reports on the incidence of brucellosis [8,9]. Studies have revealed that in a few years to

come, there is likely to be geometric increase in the incidence of brucellosis [8,9].

Diagnosis of brucellosis using fluid samples (such as milk and blood samples) is the cornerstone for the control and eradication of *Brucella* infection [11]. Traditional diagnostic methods such as culturing and serological tests (such as milk ring test) are often used for diagnosis of brucellosis. Culturing of *Brucella* species is difficult due to its fastidious nature and coupled with laborious biochemical tests for its identification [12]. The serologic method only detects the presence of *Brucella* antibody in serum of infected individual, it often gives false positive/negative reactions, cross-reactions between *Brucella* antigen is common including cross-reactions with other bacteria including *Yersinia enterocolitica*, *Campylobacter fetus*, *Vibrio cholera*, *Bordetella bronchiseptica* and *Salmonella* species [13,14], it is not specific, and does not distinguish active and non-active infection following post-treatment antibody responses [7,15,16]. Thus, these methods do not detect any *Brucella* species. Further, these methods are limited because they are time-consuming, low sensitivity when there is low amount of living *Brucella* organisms in the blood and the risk of laboratory personnel being infected following inhalation of aerosol droplets [13,17]. This makes treatment of the infection difficult and moreso limits efforts being made to reduce the prevalence of *Brucella melitensis* infection.

However, molecular detection using polymerase chain reaction (PCR)-based test has proved to be fast (> 4 h) and sensitive for the diagnosis of brucellosis [18,19]. Another advantage is that PCR is capable of detecting few *Brucella* cells or the minutest *Brucella* gene copies in samples [20]. The aim of this study was to detect molecularly *B. melitensis* in the milk of sheep and goats reared in Iran.

EXPERIMENTAL

Milk samples

This study was a cross-sectional conducted. A total of 225 apparently healthy animals (125 sheep and 100 goats) were sampled randomly. Milk samples consisting of 125 samples from Lori-Bakhtiary sheep breeds and 100 samples from traditional goat breeds, were collected in Isfahan and Shahrekord areas in central Iran. Sampled animals were members of flocks with a history of abortion. The milk samples were

collected aseptically in a sterile cup with a lid and aseptically transported to the Islamic Azad University of Shahrekord laboratory within 10 minutes of collection. Samples were divided into 0.5 mL of sterile 2-mL Eppendorf tubes, and kept frozen (-20 °C) until used [7].

DNA extraction and PCR

Brucella melitensis DNA was extracted from milk by the method of Leal-Klevazas *et al* [20]. Briefly, frozen milk samples were thawed at room temperature and 400 µL of lysis solution (100 mM Tris-HCl (pH 8), 100 mM NaCl, 1 % Sodium dodecyl sulfate (SDS), 2 % Triton-x100) and 10 µL of proteinase K (10 mg/mL) were added to 400 µL of the fatty top layer of each milk sample. The contents were incubated at 50 °C for 30 min. Thereafter, 400 µL of saturated phenol (liquid phenol containing 0.1 % 8-hydroxyquinoline, saturated and stabilized with 10 mM Tris-HCl (pH 8) and 0.2 % of 2-mercaptoethanol) were added, mixed thoroughly and centrifuged at 8000 ×g for 5 min. The aqueous layer was transferred to a fresh tube and an equal volume of chloroform: isoamyl alcohol (24:1) was added, mixed thoroughly and centrifuged as described above.

The upper layer was again transferred to a fresh tube and an equal volume of 7.5 M ammonium acetate was added and mixed thoroughly. The contents were kept on ice for 5 min, centrifuged at 8,000 × g for 5 min and the aqueous content was transferred to a fresh tube. Two volumes of 95 % ethanol were added, mixed and the tubes were stored at -20 °C for 12 h. DNA was recovered by the final centrifugation as described above, the pellets were rinsed with 1 ml of 70 % ethanol, dried and resuspended in 30 µL TE buffer (Appli Chem, Darmstadt, Germany). In addition, a commercial DNA extraction kit (Dneasy® Tissue Kit, Qiagen, Hilden, Germany) was also used in the study. For this purpose, 25 µg of the fatty top layer were used as the initial extraction material. Subsequent extraction stages were applied according to the manufacturer's recommendations. Extracted DNA was stored at -20 °C until processed.

Synthetic oligonucleotide design

The *B. melitensis*-specific primers used were previously described by Bricker and Halling [21]. The sequences of the primers were 5'-AAATCGCGTCCTTGCTGGTCT GA-3' (*B. melitensis*-specific primer) and 5'-TGCCGATCACTTAAGGGCCTTCAT-3' (IS711-specific primer from Sina Gene, Iran).

DNA amplification and detection of PCR products

PCR was carried out in a total volume of 50 μ l, using 10 mM Tris-HCl (pH 9), 3 mM MgCl₂, 50 mM KCl, 0.1 % Triton-x100, 200 mM of each of the four deoxynucleotide triphosphates (Lavora, Tellow, Germany), 0.4 mM of each primer (50 pmol), 2 IU of Taq polymerase (Fermentas, Opelstrasse 9, Leon-Rot, Germany) and 2 μ l template. The amplification was performed in a DNA thermal cycler (Thermo, Px2 Thermal Cycler, USA) as follows: initial denaturation step at 94 °C for 4 min, and 35 cycles of 94 °C for 1 min, 60 °C for 1 min and 72 °C for 1 min. The final incubation was at 72 °C for 5 min [20,21].

Amplification products were resolved in a 1.5 % (w/v) agarose gel containing 1xTBE buffer (100 mM Tris-HCl (pH 8), 90 mM boric acid and 1 mM Na₂EDTA) and stained with ethidium bromide (0.5 μ g/mL) and evaluated by a computerized image analysis system (Spectronics Co., GI-5000, England). A visible band of appropriate size (731 bp) was considered as a positive reaction for *B. melitensis*. A positive control (based on DNA from *B. melitensis* 16 M) and a negative control (DNases and RNases free water, AppliChem) were included in all the tests. To check the reliability of the results and to detect any external contamination, all samples were processed in duplicate.

Determination of detection limit of PCR for inoculated milk

B. melitensis 16 M strain was grown on trypticase soy agar (TSA, Merck) at 37 °C for 48 h. A single colony was removed from TSA, placed in trypticase soy broth (Merck) and incubated at 37 °C for 48 h. Thereafter, the culture was prepared in sterile saline and 10-fold dilutions (from 10⁻¹ to 10⁻¹⁰) were made. From these dilutions, 0.1 ml suspension was inoculated onto TSA plates and incubated at 37 °C for 48 h and the colonies present were then enumerated. The number of organisms in the dilutions was estimated spectro-photometrically at 623 nm and the concentration of the original *B. melitensis* culture was estimated as 1.7 \times 10⁹ CFU/ml (OD 0.18). To assess the limit of detection of the PCR assays, 10 raw milk

samples collected from Brucella-free sheep from the Research Farm of Faculty of Veterinary Medicine, Islamic Azad University Shahrekord Branch, Iran, were artificially contaminated with a known decreasing number of pure *B. melitensis* 16 M strain. The final concentrations of the organism in milk was 1.7 \times 10⁸, 1.7 \times 10⁷, 1.7 \times 10⁶, 1.7 \times 10⁵, 1.7 \times 10⁴, 1.7 \times 10³, 1.7 \times 10² and 1.7 \times 10¹ CFU/mL. *B. melitensis* DNA was extracted from all dilutions of milk, and processed by PCR as described earlier. The final concentrations of organisms were verified by plating onto TSA.

Statistical analysis

All statistical analyses were performed at 95 % confidence interval (CI) using Win Episcopo version 2.0 programme. The level of significance was set at $p < 0.05$.

RESULTS

Out of 225 milk samples, 20 (8.9 %) were positive for the PCR of *B. melitensis*. Out of 125 sheep milk, 12 (9.6 %) were positive for PCR of *B. melitensis*. Of these, 8 (66.6 %) were milk collected from Sharekhord while 4 (33.3 %) were from Isfahan region. Out of 100 goat milk, 18 (18 %) were positive for the PCR of *B. melitensis*. Of these 10 (55.5 %) were from Sharekhord while 8 (44.4 %) were from Isfahan (Table 1 and Figure 1).




Figure 1: *Brucella melitensis* PCR products obtained from local isolate. Lane M: Marker (GeneRuler 100bp DNA Ladder, Fermentas); Lane 1: Negative Control; Lane 2-5: Local isolate from sheep and goats milk farms in current study

Table 1: PCR data for *Brucella melitensis* in two regions of Iran

Milk type	No. of samples	Total no. positive (%)	No. positive (%) in Shahrekord Region	No. positive (%) in Isfahan Region
Sheep milk	125	12 (9.6 %)	8 (66.6 %)	4 (33.3 %)
Goats milk	100	18 (18 %)	10 (55.5 %)	8 (44.4 %)

No significant differences ($p > 0.05$) were found between sheep and goats milk samples positive for *B. melitensis* in Isfahan and Shahrekord, Iran. A positive PCR result on the ethidium bromide stained agarose gel was detected with different aliquots containing *B. melitensis* at a density of at least 1.7×10^3 - 1.7×10^4 CFU/ml in milk (Figure 2).




Figure 2: Detection limit of *B. melitensis* 16 M strain in inoculated milk by PCR assay. Lane 1: Marker (GeneRuler 100bp DNA Ladder, Fermentas); Lanes 2-8: containing decreasing number of *B. melitensis* 16 M per ml of milk. Lane 2: 1.7×10^8 ; Lane 3: 1.7×10^7 ; Lane 4: 1.7×10^6 ; Lane 5: 1.7×10^5 ; Lane 6: 1.7×10^4 ; Lane 7: 1.7×10^3 ; Lane 8: 1.7×10^2 cfu/ml; Lane 9: control positive (based on DNA from *B. melitensis* 16 M strain); Lane 10: control negative (DNases and RNases free water, AppliChem)

DISCUSSION

PCR assay is a specific and sensitive choice for the detection of different bacterial agents [2,22,23]. In this study, 9.6 % (12/125) of ovine and 18 % (18/100) of caprine milk samples were positive for PCR of *B. melitensis* biovar 3. This suggests that sheep and goat infection by *B. melitensis* in Iran is relatively lower when compared with other regions. For instance, Erdenlig and Sen [24] reported 88.5 % *B. melitensis* biovar 3 among 78 *Brucella* isolates from different regions of Turkey [26]. In Central Anatolia region of Turkey, 94.8 % among 39 *Brucella* isolates from sheep were *B. melitensis* biovar 3 [25].

It has been reported that the detection limit of the PCR in milk samples range from 10 bacteria/ml [26], 1000 CFU/ml [11], 2.8×10^4 CFU/ml [18] to 4.2×10^4 CFU/ml [28]. In this study, 1.7×10^3 - 1.7×10^4 CFU/ml of *B. melitensis* 16 M strain was detected. The lower detection rate recorded in the current study may be attributed to the DNA extraction procedures.

The extraction procedure used in this study had successfully been employed in the detection of *B. melitensis* DNA in sheep/goat milk [26]. In a PCR study by Hamdy and Amin [27], 39 milk samples were collected from 21 sheep and 18 goats. In agreement with our results, Leal-Klevezas *et al* [20] also detected as positive a higher positive number of milk samples by PCR assay when compared to bacteriological culture methods [20].

The PCR results achieved in this study are in agreement with the results obtained in previous studies, i.e. less variable than the results of bacteriology or serology [3,20,24]. Deficient isolation techniques or the stage of infection may explain the superiority of the PCR assay to isolation methods. Moreover, PCR assay detects minutest traces of genetic material in samples, while culture methods detect only viable organisms [6].

The specificity of the primers used in the current study has been evaluated with a variety of microorganisms that have a close antigenic relationship with *Brucella* which causes false-positive results in serology, and the absence of amplification with DNA of these species has shown the primers to be specific for *B. melitensis* biovars 1, 2 and 3 [11,28]. In this study, *B. melitensis* DNA was detected in sheep and goat milk samples by PCR. Gupta *et al* [19] revealed that the sensitivity and specificity of PCR in detecting the presence of *B. melitensis* in goat milk were 90 and 100 %, respectively [19]; and our results showed almost the same sensitivity and specificity.

The determination of *B. melitensis* from sheep and goat milk samples is important because raw ovine/caprine milk is used in the production of traditional cheese in Isfahan and Shahrekord, Iran [6].

CONCLUSION

The findings of this study show that a sizeable percentage of sheep and goat reared in Iran are infected with *B. melitensis* biovar 3 which is excreted in their milk. Consumption of traditional cheese may be a route for transmission of brucellosis to humans. Control of brucellosis in animals will subsequently result in decreased incidence of the disease in humans. To effect rapid and accurate diagnosis of *Brucella* load/status of caprine/ovine milk for human consumption is paramount for public health.

ACKNOWLEDGEMENT

The authors would like to thank Islamic Azad University, Shahrekord Branch for financially supporting this research through Project 12875.

CONFLICT OF INTEREST

No conflict of interest associated with this work.

CONTRIBUTION OF AUTHORS

We declare that this work was done by the authors named in this article and all liabilities pertaining to claims relating to the content of this article will be borne by the authors.

REFERENCES

1. Khamesipour F, Doosti A, Taheri H. Molecular Detection of *Brucella* spp. in the Semen, Testis and Blood Samples of Cattle and Sheep. *J Pure Appl Microbio.* 2013; 7: 495-500.
2. Taktaz-Hafshejani T, Khamesipour F, Heydari A, Katsande S. Molecular prevalence of *Brucella abortus*, *Actinomyces pyogenes* and *Mycobacterium tuberculosis* in reproductive organs of apparently healthy rams slaughtered in Iran. *Rev Med Vet.* 2015; 166: 132-137.
3. Leyla G, Kadri G, Umran O. Comparison of polymerase chain reaction and bacteriological culture for the diagnosis of sheep brucellosis using aborted fetus samples. *Vet Microbiol.* 2003; 93: 53-61.
4. Khamesipour F, Rahimi E, Shakerian A, Doosti A, Momtaz H. Molecular study of the prevalence of *Brucella abortus* and *Brucella melitensis* in the blood and lymph nodes samples of slaughtered camels by Polymerase Chain Reaction (PCR) in Iran. *Acta Vet.* 2014; 64: 245-256.
5. Safarpour Dehkordi F, Khamesipour F, Momeni M. *Brucella abortus* and *Brucella melitensis* in Iranian Bovine and Buffalo Semen Samples: The First Clinical Trial on Seasonal, Senile and Geographical Distribution Using Culture, Conventional and Real-time Polymerase Chain Reaction Assays. *Kafkas Univ Vet Fak Derg.* 2014; 20 (6): 821-828. DOI: 10.9775/kvfd.2014.10827.
6. Shakerian A, Karim G, Sharifzadeh A, Sadeghy M. The Survey on the contamination of ewes fresh white cheese non pasteurized with *Brucella mellitensis*, *Escherichia coli* and *Staphylococcus aureus* in Shahrekord, Iran. *Iranian J Vet Sci.* 2005; 2(4): 275-282.
7. Mohamed NS, Stephen MB, Nammalwar S. *Brucella*: A pathogen without classic virulence genes. *Vet Microbiol.* 2008; 129: 1-14.
8. Delrue RM, Deschamps C, Leonard S, Nijssens C, Danese I, Schaus JM, Bonnot S, Ferooz J, Tibor, A, De Bolle X, Letesson JJ. A quorum sensing regulator controls expression of both the type IV secretion system and the flagellar apparatus of *Brucella melitensis*. *Cell Microbiol.* 2005; 7: 1151-1161.
9. Comerci DJ, Martinez-Lorenzo MJ, Sieira R, Gorvel JP, Ugalde RA. Essential role of the *VirB* machinery in the maturation of the *Brucella abortus*-containing vacuole. *Cell Microbiol.* 2001; 3: 159-168.
10. Foulongne V, Bourg G, Cazevieuille C, Charachon S, O'Callaghan D. Identification of *Brucella suis* genes affecting intracellular survival in an in vitro human macrophage infection model by signature-tagged transposon mutagenesis. *Infect Immun.* 2000; 68: 1297-1303.
11. Seleem MN, Boyle SM, Sriranganathan N. Brucellosis: A re-emerging zoonosis. *Vet Microbiol.* 2010; 140: 392-398.
12. Stack JA, Harrison M, Perrett LL. Evaluation of a selective medium for *Brucella* isolation using natamycin. *J Applied Microbiol.* 2002; 92: 724-728.
13. OIE: Office International des Epizooties. *Manual of Diagnostic Tests and Vaccines for Terrestrial Animals 2009 OIE (Ed.) Caprine and Ovine Brucellosis (Excluding Brucella ovis).* OIE Manual. 2009.
14. Alton GG, Jones LM, Angus RD, Verger JM. *Techniques for the brucellosis laboratory.* Institute National de la Recherche Agronomique, Paris, France, 13-61, 1988.
15. Kara R, Akkaya L. Investigation of *Brucella abortus* and *Brucella melitensis* at Cheeses in Afyonkarahisar, Turkey. *Br J Dairy Sci.* 2013; 3(1): 5-8.
16. Lavigne JP, Patey G, Sangari FJ, Bourg G, Ramuz M, O'Callaghan D. Identification of a new virulence factor, *BvfA*, in *Brucella suis*. *Infect Immun.* 2005; 73: 5524-5529.
17. Nagalingam M, Shome R, Balamurugan V, Shome B, NarayanaRao K, Vivekananda V, Isloor S, Prabhudas K. Molecular typing of *Brucella* species isolates from livestock and human. *Trop Anim Health Produc.* 2012; 44: 5-9.
18. Romero C, Lopez-Goni I. Improved method for purification of bacterial DNA from bovine milk for detection of *Brucella* spp. by PCR. *Appl Environ Microbiol.* 1999; 65: 3735-3737.
19. Gupta VK, Verma DK, Rout PK, Singh SV, Vihan VS. Polymerase chain reaction (PCR) for detection of *Brucella melitensis* in goat milk. *Small Rumin Res.* 2006; 65: 79-84.
20. Leal-Klevezas DS, Martinez-Vazquez IO, Garcia-Cantu J, Lopez-Merino A, Martinez-Soriano JP. Use of polymerase chain reaction to detect *Brucella abortus* biovar 1 in infected goats. *Vet Microbiol.* 2000; 75: 91-97.
21. Bricker BJ, Halling SM. Differentiation of *Brucella abortus* bv. 1, 2 and 4, *Brucella melitensis*, *Brucella ovis* and *Brucella suis* bv. 1 by PCR. *J Clin Microbiol.* 1994; 32: 2660-2666.
22. Khamesipour F, Doosti A, Emadi MF, Awosile B. Detection of *Brucella* sp. and *Leptospira* sp. in dogs using conventional polymerase chain reaction. *Bull Vet Inst Pulawy.* 2014; 58: 527-531.

23. Khamesipour F, Doosti A, Rahimi E. Molecular study of brucellosis in camels by the use of TaqMan Real-time PCR. *Acta Microbiol Immunol Hung.* 2015; 62 (4): 411–424.
24. Erdenlig S, Sen A. Isolation and biotyping of *Brucella* species from aborted ewes fetuses. *J Pendik Vet Microbiol.* 2000; 31: 31-42.
25. Guler L, Gunduz K, Ok U. Comparison of polymerase chain reaction and bacteriological culture for the diagnosis of sheep brucellosis using aborted fetus samples. *Vet Microbiol.* 2003; 93: 53-61.
26. Ilhan Z, Solmaz H, Aksakal A, Gulhan T, Ekin IH, Boynukara B: Detection of *Brucella melitensis* DNA in the milk of sheep after abortion by PCR assay. *Arch Med Vet.* 2008; 40: 141-146.
27. Hamdy MER, Amin AS. Detection of *Brucella* species in the milk of infected cattle, sheep, goats and camels by PCR. *Vet J.* 2002; 163: 299-305.
28. Falenski A, Mayer-Scholl A, Filter M, Gollner C, Appel B, Nockler K. Survival of *Brucella* spp. in mineral water, milk and yogurt. *Int J Food Microbiol.* 2011; 145: 326-330.