

## Research Article

# Antioxidant and antimicrobial activities of *Heracleum nepalense* D Don root

Suvakanta Dash<sup>1\*</sup>, Lila Kanta Nath<sup>1</sup>, Satish Bhise<sup>2</sup> and Nihar Bhuyan<sup>1</sup>

<sup>1</sup>Himalayan Pharmacy Institute, Majhitar, Rangpo, East Sikkim-737132, India. <sup>2</sup>Govt College of Pharmacy, Vidyanagar, Satara, Karad- 415124, India.

## Abstract

**Purpose:** The aim of the present study was to investigate antioxidant and antimicrobial effects of the methanol extract of *Heracleum nepalense* D. Don roots.

**Method:** The antimicrobial effect was determined by agar dilution and disc diffusion method. The free radical scavenging potential was studied by using different antioxidants models of screening using vitamin E (5mM) as standard.

**Results:** The crude methanol extract of *H. nepalense* root was found to be active against both Gram-positive and Gram-negative organisms. The ethyl acetate soluble fraction of the extract showed similar activity against these organisms. Similarly, the methanol extract at 1000  $\mu\text{g. ml}^{-1}$  and the ethyl acetate fraction at 50  $\mu\text{g. ml}^{-1}$  exhibited significant antioxidant activity in ferrous sulphate induced lipid peroxidation, 1,1- diphenyl- 2-picryl hydrazyl (DPPH), Hydroxyl radical and Superoxide scavenging models.

**Conclusions:** The study confirms the possible antioxidant and antimicrobial potentiality of the plant extract. Presence of flavonoid alone or in combination with its other components could be responsible for the activity.

**Keywords:** *Heracleum nepalense*, Lipid peroxidation, Superoxide scavenging, DPPH assay, Antimicrobial effect, Flavonoid.

---

\*Corresponding Author, E-mail: [sdash777@sify.com](mailto:sdash777@sify.com)

## Introduction

Reactive oxygen species (ROS) such as superoxides, peroxides and hydroxyl radicals are known to play an important role and have been identified as major contributors to all cell and tissue damage in many disease conditions<sup>1</sup>. In living organisms, various ROS can form in different ways. Normal aerobic respiration stimulates polymorphonuclear leukocytes and peroxisomes appear to be the main endogenous sources of most of the oxidants produced by cells. Exogenous sources of ROS include tobacco smoke, alcohol, pesticides, certain pollutants and microbial infections<sup>2-3</sup>.

In recent years, there has been increased incidence of antibiotic resistance in pathogenic organisms and the persistence of pathogens in immune compromised individuals is of great concern<sup>4</sup>. Therefore, actions must be taken to reduce this problem such as controlling the use of antibiotics, carrying out research to better understand the genetic mechanisms of resistance and continuing investigations aimed at the development of drugs from natural sources.

The recent years have witnessed resurgence of interest in herbal drugs globally as more people are turning to the use of herbal medicinal products in health care. About 80% of individuals from developing countries use traditional medicine, which involves compounds derived from medicinal plants. It is high time the hidden wonders of plant molecules were revived with the modern tools of target-based screening to develop newer advanced generation antioxidants and antimicrobials with novel modes of action. Antioxidants play an important role in inhibiting and scavenging free radicals, thus providing protection to human body against infection and degenerative diseases.

*Heracleum nepalense* D.Don (Apiaceae) is a small shrub occurring in Nepal and Sikkim<sup>5</sup>. The plant is used in veterinary medicine. It exhibits stimulant property and increases blood pressure in goats<sup>6</sup>. The roots of the plant are used in folk

medicine as digestive, carminative and anti-diarrhoeal (Authors personal experience). The roots of the plant are reported to have coumarins<sup>7</sup> and steroids<sup>8</sup>. The present study was aimed at evaluating antioxidant and antimicrobial properties of *Heracleum nepalense*.

## Experimental

### Plant materials and phytochemical screening

The fresh dried roots of *Heracleum nepalense* (HNSE) were collected from the southern district of Sikkim. The plant was authenticated by Botanical Survey of India, Gangtok, Sikkim. The voucher specimen was preserved in our laboratory for future reference. The dried roots were powdered and stored in a well-closed container. 1 kg of powder (40 mesh size) was extracted by cold percolation with 3 liters of 70% v/v methanol in a percolator for 72 h at room temperature<sup>9</sup>. The residue was removed by filtration. The extract was then evaporated to dryness under reduced pressure in a rotary evaporator at 42-45°C. The concentrated extract of leaves was kept in a dessicator for further use.

The preliminary phytochemical tests of the root extract were done by Pollock and Steverson method<sup>10</sup>. Concentrated methanol extract was suspended in hot distilled water, cooled and the blast precipitate was filtered off. The filtrate (aqueous solution) was fractionated by extracting it successively with petroleum ether, ethyl acetate and acetone. The ethyl acetate fraction of the crude extract on purification yielded one major fraction A (flavonoid) with some fatty substances.

### Chemicals

Thiobarbituric acid was obtained from Loba Chemie, India. 1,1-Diphenyl-2-picryl hydrazyl (DPPH), NADH and nitroblue tetrazolium (NBT) were obtained from Sigma chemicals, St. Louis, USA. Deoxy ribose was obtained from Merck India. Dimethyl sulphoxide, ethylene diamine tetra acetic acid, ferrous sulphate, trichloroacetic acid, hydrogen peroxide, ascorbic acid, mannitol, potassium dihydrogen phosphate, potassium

hydroxide, deoxy ribose, phenazine methosulphate were of analytical grade and were obtained from Ranbaxy fine chemicals.

### Determination of antioxidant activity

#### Assay of lipid peroxidation

The extent of lipid peroxidation in goat liver homogenate was measured *in vitro* in terms of formation of thiobarbituric acid reactive substances (TBARS) by using standard method<sup>11</sup> with minor modifications<sup>12</sup>. Goat liver was purchased from local slutter house. Its lobes were dried between blotting papers (to remove excess blood) and were cut into small pieces with a heavy-duty blade. They were then homogenized in glass-Teflon homogenizing tube in cold phosphate buffer saline (pH 7.4). It was centrifuged at 2000 rpm for 10 min and supernatant was diluted with phosphate buffer saline up to final concentration of protein 0.8-1.5 mg/0.1 ml. Protein concentration was measured by using standard method of Lowery et.al<sup>13</sup>. To study the comparative response, the experiment was divided into nine groups. Liver homogenate (5%, 3ml) was aliquoted to nine different 35mm glass petri dishes. The first two groups were treated as control and standard where buffer and vitamin E were added. In the third to seventh group, different concentrations of methanol extract (200-1000  $\mu\text{g}\cdot\text{ml}^{-1}$ ) and fraction A (25, 50  $\mu\text{g}\cdot\text{ml}^{-1}$ ) were added. Lipid peroxidation was initiated by adding 100 $\mu\text{l}$  of 15mM ferrous sulphate solution to 3ml of liver homogenate<sup>14</sup>. After 30 minutes, 100  $\mu\text{l}$  of this reaction mixture was taken in a tube containing 1.5 ml of 10% Trichloro acetic acid. After 10 minutes, tubes were centrifuged and supernatant was separated and mixed with 1.5ml of 0.67% thiobarbituric acid. The mixture was heated in a water bath at 85°C for 30 min and in a boiling water bath to complete the reaction. The intensity of pink colored complex formed was measured at 535nm in a spectrophotometer (Shimadzu model 1601). The percentage of inhibition of lipid peroxidation was calculated by comparing the results of the test with those of controls as per the following formula:

Inhibition (%) = [(control - test)  $\times$  100] / control----  
--- Eqn 1.

### DPPH radical scavenging activity

DPPH scavenging activity was measured by spectrophotometric method<sup>14</sup>. To a methanolic solution of DPPH (100  $\mu\text{M}$ , 2.95ml), 0.05 ml of methanol extract as well as fraction A (25, 50  $\mu\text{g}\cdot\text{ml}^{-1}$ ) and Standard compound (vitamin E) were added at different concentrations. Equal amount (0.05 ml) of methanol was added to a control. Absorbance was recorded at 517nm at regular intervals of 1 to 5 min. The percentage of scavenging was calculated by comparing the control and test samples with the Eqn 1.

### Hydroxyl radical scavenging activity

Hydroxyl radical scavenging activity was measured by studying the competition between deoxy ribose and test compounds for hydroxyl radical generated by the  $\text{Fe}^{3+}$  - ascorbate - EDTA -  $\text{H}_2\text{O}_2$  system (Fenton reaction) according to the method of Kunchandy & Rao<sup>15</sup>. The reaction mixture containing, a final volume of 1.0 ml, 100  $\mu\text{l}$  2-deoxy-ribose, 500  $\mu\text{l}$  of the various concentrations of the methanol extract as well as fraction A (25, 50  $\mu\text{g}\cdot\text{ml}^{-1}$ ) and standard compound (Mannitol 50 mM) in  $\text{KH}_2\text{PO}_4$ - KOH buffer (20mM, pH 7.4), 200  $\mu\text{l}$  1.04 mM  $\text{H}_2\text{O}_2$  and 100  $\mu\text{l}$  1.0mM ascorbic acid was incubated at 37°C for 1 hour. One milliliter 1% trichloroacetic acid was added to each test tube and incubated at 100°C for 20 min. After cooling to room temperature, absorbance was measured at 532nm against a control preparation containing deoxyribose and buffer. Percent inhibition was determined by comparing the results of the test and control samples with the above mentioned Eqn 1.

### Superoxide scavenging activity

The superoxide scavenging activity of methanol extract was determined by the method described by Nishimik et al<sup>16</sup>, with slight modification. About 1.0 ml NBT solution containing 156  $\mu\text{M}$  NBT dissolved in 1.0 ml 100 mM phosphate buffer, pH 7.4, 1.0 ml NADH solution containing 468  $\mu\text{M}$  NADH dissolved in 1.0 ml 100mM phosphate buffer, pH 7.4, and 0.1 ml of various concentration of the methanol extract as well as fraction A (25, 50  $\mu\text{g}\cdot\text{ml}^{-1}$ ) and standard compound (vitamin E) were mixed and the reaction was started by adding 100  $\mu\text{l}$  of

phenazine methosulfate solution containing 60 $\mu$ M phenazine methosulphate in 100 mM phosphate buffer, pH 7.4. The reaction mixture was incubated at 25 °C for 5 min and absorbance at 560 nm was measured against control sample. Percent inhibition was determined by comparing the results of the test and control samples with the above mentioned Eqn 1.

### **Antimicrobial activity**

#### **Bacteria**

A total of 257 bacterial strains belonging to different genera were tested in this study. The test organisms were obtained from Department of Bacteriology, Calcutta School of Tropical Medicine and Institute of Microbial Technology (IMTECH), Chandigarh, India. Three multiresistant *Staphylococcus* strains (*Staphylococcus aureus* ML 275, *Staphylococcus aureus* NCTC 8530 and *Staphylococcus epidermidis* 865) were kindly provided by Prof. (Mrs.) Sujata Ghosh Dastidar, Department of Pharmacy, Jadavpur University, and Kolkata, India. They were aseptically isolated and identified by the Barrow and Feltham's method and preserved in the freeze-dried state<sup>17</sup>. Gram-positive strains were grown in nutrient broth (NB, Oxoid brand) and Gram-negative bacteria were grown in peptone water (PW, Oxoid brand, bacteriological peptone plus NaCl 0.5%) for 18 h before use.

#### **Determination of antimicrobial activity**

Sensitivity tests were performed by disc diffusion method<sup>18</sup>. The nutrient agar plates (Oxoid brand), containing an inoculum size of 10<sup>5</sup>-10<sup>6</sup> cfu.ml<sup>-1</sup> of bacteria were used. Previously prepared crude methanol extract (Concentration 128-2000  $\mu$ g.ml<sup>-1</sup>) and fraction A (Concentration 128-2000  $\mu$ g.ml<sup>-1</sup>) discs were placed aseptically on sensitivity plates. The discs containing methanol and known antibiotics (Amoxycillin and Gentamicin) served as negative and positive controls. All the plates were then incubated at 37°C $\pm$ 2°C for 18 h. The sensitivity was recorded by measuring the clear zone of inhibition on agar plate around the discs.

The MICs were determined by the standard agar dilution method<sup>19</sup>. All the test compounds were dissolved in methanol. These were then individually added at each final concentrations of 0-2000  $\mu$ g.ml<sup>-1</sup>, to molten agar (Oxoid brand), mixed thoroughly, pH adjusted to 7.2 to 7.4 and poured into sterile Petri dishes. Bacterial cell suspensions were spot inoculated on the plates using a bacterial planter (10  $\mu$ l). The final number of cfu inoculated onto the agar plates was 10<sup>4</sup> for all strains. The inoculated plates were then incubated at 37°C $\pm$ 2°C for 18 h. The lowest concentration of the plate, which did not show any visible growth after incubation, was considered as MIC. The agar plate containing only methanol and Amoxycillin was served as negative and positive control.

#### **Statistical analysis**

Data are reported as the mean  $\pm$  SD of three measurements. Statistical analysis was performed by the student *t*-test and by ANOVA. IC<sub>50</sub> values for all the above experiments were determined by linear regression method. A *p*-value less than 0.05 was considered as indicative of significance.

### **Results and Discussion**

#### **Antioxidant activity**


##### *Assay of lipid peroxidation*

The results presented in Table-1 showed that the methanol extract of the HNSE inhibited FeSO<sub>4</sub> induced lipid peroxidation in a dose dependent manner. The extract at 1000  $\mu$ g.ml<sup>-1</sup> exhibited maximum inhibition (69.25  $\pm$  1.21%) of lipid peroxidation, on the other hand fraction A at 50  $\mu$ g.ml<sup>-1</sup> concentrations showed (72.38  $\pm$  1.9%) inhibition, nearly equal to the inhibition produced by vitamin E (Fig 1). The IC<sub>50</sub> value was found to be 747.5  $\pm$  3.16  $\mu$ g.ml<sup>-1</sup>. The inhibition could be caused by the absence of ferryl-perferryl complex or by changing the ratio of Fe<sup>3+</sup> / Fe<sup>2+</sup> or by reducing the rate of conversion of ferrous to ferric or by chelating the iron itself or combination thereof<sup>20</sup>.

#### **DPPH scavenging activity**

DPPH is a stable free radical that can accept an electron or hydrogen radical to become a stable dimagnetic molecule. Due to its odd electron, the

methanolic solution of DPPH shows a strong



**Fig 1:** Inhibition (%) of lipid peroxidation by different concentrations of methanol extract of *H. nepalense*, Ethyl acetate fraction (FA) and Vitamin E (Vit E).

absorption at 517 nm. DPPH radical reacts with suitable reducing agents and then electrons become paired off and the solution loses colour stoichiometrically with the number of electrons taken up<sup>21</sup>. Such reactivity has been widely used to test the ability of compound/plant extracts to act as free radical scavengers. Reduction of the DPPH radicals can be observed by the decrease in absorbance at 517 nm. The DPPH scavenging capacity of the extract was found to be  $72.38 \pm 3.92\%$  at  $1000 \mu\text{g}\cdot\text{ml}^{-1}$ . The fraction A at  $50 \mu\text{g}\cdot\text{ml}^{-1}$  doses, on the other hand, exhibited  $76.38 \pm 5.12\%$  inhibition compared with  $80.46 \pm 4.62\%$  for the standard drug vitamin E at 5mM (Table 1). The  $\text{IC}_{50}$  value was found to be  $6.5 \text{ mg}\cdot\text{ml}^{-1}$ . The activity was also dependent on time (Fig 2).


#### Hydroxyl radical scavenging activity

Hydroxyl radicals are the major active oxygen species causing lipid peroxidation and enormous biological damage<sup>22</sup>. Ferric-EDTA was incubated with  $\text{H}_2\text{O}_2$  and ascorbic acid at pH 7.4. Hydroxyl radicals were formed in free solution and were detected by their ability to degrade 2-deoxy-2-ribose into fragments that formed a pink chromogen upon heating with TBA at low pH<sup>23</sup>. When the test compounds were added to the reaction mixture they removed hydroxyl radicals from the sugar and prevented their degradation. The extract of HNSE significantly inhibited

( $80.38 \pm 2.28\%$ ) degradation of deoxy-ribose mediated by hydroxyl radicals at the dose of  $1000 \mu\text{g}\cdot\text{ml}^{-1}$  (Table 1), compared to that of a known scavenger mannitol (50mM). The concentration of the methanol extract needed for 50% inhibition was  $615.57 \mu\text{g}\cdot\text{ml}^{-1}$ . The fraction A at  $50 \mu\text{g}\cdot\text{ml}^{-1}$  on the other hand, exhibited ( $78.68 \pm 2.62\%$ ) inhibition compared with ( $89.64 \pm 2.84\%$ ) for the standard mannitol.

#### Superoxide scavenging activity

Superoxide radical  $\text{O}_2^-$  is a highly toxic specie, which is generated by numerous biological and photochemical reactions. Both aerobic and anaerobic organisms possess superoxide dismutase enzymes, which catalyse the



**Figure 2:** DPPH scavenging activity of *H. nepalense*. 10mg/ml (□), 7.5mg/ml (▲), 5mg/ml (■) and 2.5 mg/ml (△).

breakdown of superoxide radical<sup>24</sup>. Reduced phenazine methosulfate assay was used to measure the superoxide dismutase activity of HNSE. The results presented in Table 1 showed that the scavenging capacity of the extract was  $60.57 \pm 2.34\%$  at  $1000 \mu\text{g}\cdot\text{ml}^{-1}$ . The fraction at  $25 \mu\text{g}\cdot\text{ml}^{-1}$  exhibited  $68.36 \pm 2.73\%$  inhibition of superoxide radicals.  $\text{IC}_{50}$  was found to be  $8.9 \text{ mg}\cdot\text{ml}^{-1}$ . Inhibition was proportional to the amount of the extract added.

#### Antimicrobial activity

The methanol extract of HNSE roots exhibited a significant *in vitro* antimicrobial activity against 257 Gram-positive and Gram-negative bacteria including multiresistant *Staphylococcus* (MRSC)

strains. All the three reference strains of bacteria were found to be sensitive within  $1000 \mu\text{g.ml}^{-1}$ . The preliminary biological screening of the fractions showed that the ethyl acetate part was more active than other fractions (Data not shown). The results of the antimicrobial spectrum of the roots extract presented in Table 2 showed that out of 257 bacteria, the growth of 197 isolates were inhibited at a concentration of  $128 - 512 \mu\text{g.ml}^{-1}$ . 57 isolates were resistant up to  $1000 \mu\text{g.ml}^{-1}$ , while the remaining 03 isolates were resistant up to  $>2000 \mu\text{g.ml}^{-1}$ , the highest concentration tested. The MICs tests revealed that 63 out of 75 Gram-positive bacteria were

$128 \mu\text{g.ml}^{-1}$ , while they were resistant to the two test antibiotics. Thus, ethyl acetate fraction may become clinically relevant, particularly for antibiotic-resistant strains. However, the activity has to be studied using isolated pure compounds from the fraction to which the more resistant strains were susceptible in order to confirm these findings.

#### Acknowledgement

The authors are thankful to the All India Council for Technical Education, Government of India, New Delhi and Prof (Dr) S.G.Dastidar, Division of Microbiology, Jadavpur University, Kolkata,

**Table: 1** Effect of *H.nepalense* on different antioxidant models.

Concentration ( $\mu\text{g/ml}$ )	Percentage of Inhibition (%)			
	Lipid Peroxidation	DPPH	Hydroxyl radical	Superoxide radical
1000	$69.25 \pm 1.21$	$72.38 \pm 3.92$	$80.38 \pm 2.28$	$60.57 \pm 2.34$
800	$51.74 \pm 1.92$	$69.42 \pm 3.86$	$79.54 \pm 1.24$	$52.27 \pm 3.18$
600	$36.36 \pm 1.84$	$45.52 \pm 3.73$	$65.37 \pm 2.26$	$40.26 \pm 2.16$
400	$26.57 \pm 3.8$	$36.46 \pm 2.32$	$46.28 \pm 1.89$	$24.12 \pm 1.38$
200	$16.08 \pm 4.3$	$8.47 \pm 1.83$	$32.14 \pm 2.24$	$12.23 \pm 1.42$
Vitamin E (5mM)	$73.42 \pm 2.3$	$80.46 \pm 4.62$	NT	$68.36 \pm 2.73$
Mannitol (50mM)	NT	NT	$89.64 \pm 4.62$	NT
Fraction A				
25	$70.15 \pm 1.64$	$71.32 \pm 1.85$	$72.35 \pm 2.93$	$68.36 \pm 2.73$
50	$72.38 \pm 1.9$	$76.38 \pm 5.12$	$79.68 \pm 2.62$	$68.24 \pm 1.86$
$\text{IC}_{50}$ ( $\mu\text{g/ml}$ )	$747.5 \pm 3.16$	$600.52 \pm 3.46$	$615.57 \pm 2.16$	$891.9 \pm 14.42$

Values are mean  $\pm$  S.E.M of 3 replicates. NT: Not tested.

sensitive between 128 and 256  $\mu\text{g/ml}$  (zone diameter 10–16 mm); while out of 179 Gram-negative isolates, 120 were sensitive between 256–512  $\mu\text{g.ml}^{-1}$  (zone diameter 10-14 mm). Hence, it appears that the antimicrobial activity of the methanol extracts was directed both against Gram-positive and Gram-negative bacteria. The ethyl acetate fraction (fraction A) was also tested for antimicrobial activity. The result revealed that all the isolates were sensitive at 128–256  $\mu\text{g.ml}^{-1}$  (Table 3). It was interesting to note that all the MRSC were susceptible to fraction A at a concentration of

India for providing the research facility.

#### References

- Halliwell B. Free radicals, antioxidants and human diseases: Curiosity, cause or consequences. *Lancet*, 1994; 344: 721-724.
- Robinson EE, Maxwell SRJ, Thorpe GHG. An investigation of antioxidant activity of black tea using enhanced chemiluminescence. *Free Radi Res*, 1997; 26: 291-302.
- Davis KJA. Oxidative stress: the paradox of aerobic life. *Biochem Soci Sympos*, 1994; 61: 1-34.

4. Austin DJ, Kristinsson KG, Anderson RM. The relationship between the volume of antimicrobial consumption in human communities and the frequency of resistance, In: *Proceedings of the National Academy of Sciences of the United States of America* 96, 1999, 1152.
5. Gurung G. The medicinal plants of Sikkim Himalaya, Subash publication, Sikkim, 1999, pp 156-158.
6. The wealth of India, Vol-3, National Institute of Science Communication, CSIR, New Delhi, pp 254-255.
7. Gupta BD, Banarjee SK, Handa KL. Coumarins from *H.nepalense* and *H.wallaichi*. *Phytochemistry*, 1975; 14: 2533-2538.
8. Sun HD, Lin ZQ, Nij FD. Study of the Chinese drugs of umbelliferae II. Chemical constituents of the roots of *H.nepalense* D. Don. *Yunnan Zhiwu Yanjiu*, 1980; 2: 224-227.
9. Chattapadhaya D, Maiti K, Kundu AP, Chakrabarty MS, Bhadra R, Mandal SC, Mandal AB. Antimicrobial activity *Alstonia macrophylla*. *J.Ethnopharmacol*, 2001; 77: 49-56.
10. Pollock JRA, Stevens R. *Dictionary of organic compounds*, Eyre and Spottish woode, London, 1965, 143.
11. Ohkawa H, Oshishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid. *Anal Biochem*, 1979; 95: 351- 358.
12. Pandey S, Sharma M, Chaturvedi P, Tripathi YB. Protective effect of *R. cordifolia* on lipid peroxide formation in isolated liver homogenate. *Exp Biol*, 1995; 193: 265- 269.
13. Lowery OH, Rosenbrough NJ, Farr AL, Randall RJ. Protein estimation with Folin phenol reagent. *Biol Chem*, 1951; 193: 265-275.
14. Sreejayan N, Rao MNA. Free radical scavenging by curcuminoids. *J Pharm Pharmacol*, 1997; 49: 105-109.
15. Kunchandy E, Rao MNA. Oxygen radical scavenging activity of curcumin. *Int J Pharmacog*, 1990; 58: 237-240.
16. Nishimiki M, Rao NA, Appaji N, Yagi K. The occurrence of superoxide anion in the reaction of reduced phenazine methosulphate and molecular oxygen. *Biochem Biophys Res Comm*, 1972; 46: 849-854.
17. Barrow GI, Feltham RKA. *Manual for the identification of medical bacteria*. (Cambridge university press, Cambridge) 1993, 168.
18. Dash SK, Dastidar SG, Chakraborty AN. Antimicrobial property of Promazine hydrochloride. *Ind J Exp Biol*, 1977; 15: 197-199.
19. Dastidar SG, Chaudhary A, Annadurai S, Roy S, Mookhrjee M, Chakraborty AN. *In vitro* and *in vivo* antimicrobial action of Fluphenazine. *J Chemother*, 1995; 7: 201-207.
20. Braughler JM, Duncan CA, Chase IR. The involvement of iron in lipid peroxidation. Importance of ferrous to ferric ratio in initiation. *J Biol Chem*, 1986; 261: 10282-89.
21. Blois. Antioxidant determinations by the use of stable free radical. *Nature*, 1958; 26: 1199-1206.
22. Aurand LW, Boonmen NH, Gidding GG. Superoxide and singlet oxygen in milk lipid peroxidation. *J Diar Sci*; 1977:363-369.
23. Aruoma OI, Laughton MJ, Halliwell B. Carnosine, homocarnosine and anserine: could they act as antioxidants *in vivo*? *Biochem J*, 1989; 264: 863-869.
24. Govindarajan R, Vijaykumar M, Rawat AKS, Mehotra S. Free radical scavenging potential of *Picrorhiza kurroa* Royle ex Benth. *Ind J Exp Biol*, 2003; 41: 875-879.