

Original Research Article

Anti-proliferation and apoptosis effects of *Camellia nitidissima* C. W. Chi extract on A549 lung cancer cells

Zhi-hui Zhang^{1*}, Pei Li²

¹Cancer Center, Integrated Hospital of Traditional Chinese Medicine, Southern Medical University, 510315, ²Cancer Center, Nanfang Hospital/The First School of Clinical Medicine, Southern Medical University, 510515, Guangzhou, China

*For correspondence: **Email:** zhangzhahui020@126.com; **Tel:** +86 020-61650124

Sent for review: 19 August 2017

Revised accepted: 21 February 2018

Abstract

Purpose: To investigate the effect of *Camellia nitidissima* C. W. Chi extract (CNCE) on apoptosis and proliferation in A549 human lung cancer cells.

Methods: Inverted microscope was used to examine morphological changes in A549 cells after exposure to CNCE. Trypan blue staining of living cells was applied to construct the cell growth curve after treatment with varying concentrations of CNCE. The influence of CNCE on cell proliferation, apoptosis and cell cycle were determined by MTT assay. The protein expressions of key apoptosis-related enzymes were evaluated by immuno-cytochemical method.

Results: CNCE inhibited the growth of A549 lung cancer cells at the concentration range of 20 - 160 µg/mL. Flow cytometry showed that CNCE induced apoptosis in the A549 cells. The proportion of cells in G0/G1-phase increased significantly ($p < 0.01$), while the proportion of cells in S-phase and G2/M-phase decreased correspondingly, indicating that the cells were in G0/G1-phase arrest. Cell cycle arrest and apoptosis-inducing effect gradually increased with increase in CNCE concentration. With increasing concentrations of CNCE, there were significant increases in the expressions of caspase-3 ($p < 0.05$), caspase-8 ($p < 0.01$) and caspase-9 ($p < 0.05$), and significant decreases in Ki-67 ($p < 0.01$) and p21 ras protein ($p < 0.01$).

Conclusion: CNCE exerts significant inhibitory effect on the proliferation of A549 lung cancer cells, and therefore can potentially be developed for the treatment of lung cancer.

Keywords: *Camellia nitidissima*, Antitumor activity, Lung cancer, Apoptosis, Cell cycle arrest, Caspase

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited.

Tropical Journal of Pharmaceutical Research is indexed by Science Citation Index (SciSearch), Scopus, International Pharmaceutical Abstract, Chemical Abstracts, Embase, Index Copernicus, EBSCO, African Index Medicus, JournalSeek, Journal Citation Reports/Science Edition, Directory of Open Access Journals (DOAJ), African Journal Online, Bioline International, Open-J-Gate and Pharmacy Abstracts

INTRODUCTION

The morbidity and mortality from lung cancer account for 13 % and 19.4 % respectively, worldwide [1]. Lung cancer is classified into small cell lung carcinoma (SCLC) and non-small cell lung carcinoma (NSCLC). The latter (NSCLC) accounts for most (about 80 %) lung

tumors. At present, early-stage NSCLC is treated by surgical intervention, while mid-late NSCLC is treated with radiotherapy and chemotherapy. Although effective for some patients in the short run, radiotherapy and chemotherapy are associated with adverse effects, toxicity, high recurrence of tumor and poor prognosis [2]. Therefore, it is necessary to reduce the side effects of chemotherapy drugs by searching for

more effective anticancer drugs with lower toxicity. This has been a major challenge to medical science.

Camellia nitidissima C. W. Chi is a very valuable plant in Guangxi of China. Studies have shown that *Camellia nitidissima* C. W. Chi has anti-aging, blood pressure-lowering, hypoglycemic and immunity-enhancing properties [3]. It also has antioxidant activity [4], and exerts cytotoxic effects on various malignant tumors [5,6]. This study was carried out to investigate the effect of *Camellia nitidissima* C. W. Chi extract on proliferation and apoptosis in A549 human lung cancer cells.

EXPERIMENTAL

Materials

Human lung cancer cell line (A549) was purchased from the Institute of Basic Medical Sciences, Chinese Academy of Medical Sciences. *Camellia nitidissima* C. W. Chi was obtained from Guangxi, China. RPMI1640 culture medium was a product of GIBCO Company, USA). Calf serum and phosphate buffer solution (PBS, American hyclone); dimethylthiozol-2-yl-2,5-diphenyl tetrazolium bromide (MTT), dimethyl sulfoxide (DMSO), propidium iodide (PI) and trypsin were obtained from *Sigma-Aldrich* (America). Caspase-3, caspase-8, caspase-9, Ki-67, bcl-2 p21 ras monoclonal antibodies, and SP generic kits were purchased from American NeoMarkers (San Francisco, USA) .

Samples of *Camellia nitidissima* C. W. Chi were collected from Guiyang City, Guangxi Province, China in September 2016. Taxonomic identification of the plant was done by Professor Ye Hu of Southern Medical University in China. A voucher specimen (no. CNCE 20160912) was deposited in the herbarium of College of Pharmacy, Southern Medical University, China for future reference.

Instrumentation

The instruments used were 96-well plates (Corning Company, Wuhan, China); automatic CO₂ incubator (DuPont Company of America Napoc6100); 450 enzyme standard instrument (American BioRad); inverted microscope (Olympus CX41-32C02, Japan); 4 % hope blue mother liquor (Japanese BigWood Industrial Co., Ltd.) and LKB- ultrathin sections machine (Sweden). Others were YZ-1450 laminar flow ultranet work station (Suzhou Purification Equipment Company, Suzhou, China); elite flow cytometer (Coulter, USA); blood cell counting

plate; vacuum drying oven (DZF-6021 type, Shanghai Fine Macro Experiment Equipment Limited), and ultrasonic cleaner CQ25-12 type (Shanghai Scientz Biotechnology Research Institute).

Preparation of CNCE

Dry leaves of *Camellia nitidissima* C. W. Chi (50 g) were ground and extracted three times by refluxing with 95 % ethanol, each time for 2 h. The extracts were pooled and concentrated under reduced pressure to obtain CNCE.

Cell culture

The A549 cells were cultured in RPMI-1640 medium (Gibco, Rockville, MD) containing 10 % fetal bovine serum (v/v) (FBS; Hyclone, Logan, UT) and 1% Penicillin Streptomycin (100 U/mL and 100 µg/mL, respectively) at 37 °C in a humidified atmosphere containing 5 % CO₂, with 0.25 % trypsin digestion batches. The logarithmic phase of the cells were used for the study.

Drug sensitivity test

Cells at the logarithmic growth phase (in RPMI1640 medium) at a concentration of 1.0×10^5 cells/mL of cell suspension, were seeded in 48-well plates and cultured at 37 °C for 24 h, until the cells became adherent. CNCE at different final concentrations (30, 60, 120, 240 µg/mL) were added to different wells, and the wells were for different periods (24, 48 and 72 h) at 37 °C. A well containing 20 µg/mL cisplatin served as positive control, while another well containing saline in place of CNCE served as negative control.

Cell morphology assay

A suspension of A549 cells at the logarithmic growth phase in RPMI 164 medium was adjusted to a concentration of 1.0×10^5 cells/mL in a 48-well culture plate, and cultured for 24 h at 37 °C to achieve adherence. The wells were grouped into two. One group (experimental group) received different concentrations of CNCE (62.5, 125 or 250 µg/mL) in the culture medium, while the other group (control group) received 0.04 % DMSO in place of CNCE. The cells were observed under inverted microscope after 24, 48 and 72 h for changes in cell size, cell membrane and nucleus.

Growth curve and doubling time (TD) of A549 lung cancer cells

The A549 cells at logarithmic growth phase at a

concentration of 1×10^4 cells/mL were seeded in 24-well plates. One milliliter was taken from each of three wells at the same time point daily, and digested. The number of cells per well was counted using a hemocytometer, and the mean number of cells was determined. Counting was done for 8 days. A growth curve was plotted with mean cell count as vertical axis, and time as horizontal axis. Doubling time (TD) for the human lung cancer A549 cells was calculated using Eq 1.

$$TD = t \{ \log 2 / (\log N_t - \log N_o) \} \dots\dots\dots (1)$$

where t is time interval, while N_o and N_t are the initial mean count and mean count after time t , respectively.

MTT assay

A suspension of the A549 cells (0.2 mL) at logarithmic phase was seeded in a 96-well plate with each well containing 5×10^3 cells, and cultured overnight for cell adhesion. In the experimental group, the original culture medium was drained, and replaced with different concentrations of freshly-prepared CNCE (one concentration in 24 wells, and 6 wells per group). The wells of the control group medium contained 0.04 % DMSO, while wells without cells served as blank control.

All wells were incubated for 24, 48, 72 and 96 h at 37 °C in a humid atmosphere containing 5 % CO₂. The supernatant was discarded, and 20 µL of MTT solution (5 g/L) was added to each well and incubated at 37 °C for 4 h. Then, the supernatant was carefully removed by suction, and 150 µL DMSO was added to each well to dissolve the crystals. The absorbance of each solution was read at 490 nm against blank control, and the results were used for calculating the growth inhibition rate.

FCM assay

A suspension of 549 human lung cancer cells (5.0×10^4 cells/mL) at logarithmic growth phase was seeded in 24-well plates (100 µL in each well), and cultured in RPMI 1640 medium for 24 h for adherence. Thereafter, CNCE was added to a final concentration of 40 µg/mL. Wells in the control group contained an equivalent volume of normal saline. The plates were cultured at 37 °C in an atmosphere of 5 % CO₂. At 24 and 48 h, the cells in each group were collected (cell number $> 1 \times 10^6$), washed twice in cold PBS (pH 7.2 - 7.4), and centrifuged at 1000 rpm for 5 min. Then, 1 mL of cold precipitation PBS (pH

7.2 - 7.4) was added to the cell suspension, followed by 3 mL of pre-cooled 70 % ethanol.

The cell suspension was kept overnight at 4 °C, and then centrifuged at 1000 rpm for 5 min. The supernatant was discarded; and the cells were washed twice in 1 mL of cold PBS (pH 7.2 - 7.4), and centrifuged at 1000 rpm for 5 min. The cell count was adjusted to 1×10^6 / mL, and 200 µL NaseA was added, followed by incubation in a 37 °C water bath for 30 min. Iodide C ingot (PI) staining fluid 800 µL was added, mixed and kept at 4 °C in the dark for 30 min. DNA single parameter analysis was carried out to determine cell distribution and the proportion of cells in G0/D1, S, and G2/M phases, as a function of time. The results were analyzed using Modifit LT software.

Immunocytochemistry staining

The A549 cells at logarithmic phase were digested with 0.25 % trypsin, and 2×10^5 cells were seeded into a 6-well plate and cultured overnight for cell adhesion. The culture medium was removed, and 3mL of CNCE was added to different cells to final concentrations of 30, 60, 120 and 240 µg/mL. Each CNCE concentration was replicated in six parallel wells. In the control wells, CNCE was replaced with an equivalent volume of 0.04 % DMSO. The plates were incubated for 24 and 48 h; the glass slides were taken out, dried and fixed with 4 % paraformaldehyde for 10 min at 4 °C according to instructions contained in the operation manual of S-P kit. The plates were observed for appearance of clear brown granules which is indicative of positive cells.

The presence of Bcl-2 and p21 ras in cytoplasm is indicative of positive expression; Ki67 tan is indicative of positive expression in the nucleus, caspase-9 is an index of positive expression in the cytoplasm and the nucleus; while caspase-3 and caspase-8 indicate positive expression in the cytoplasm or cytoplasm and nuclei. They are expressed less in cells without changes in color, or lightly-colored cells. The cells in the CNCE plates and control plates were observed at high magnification ($\times 400$) in 5 fields, and positive cells were counted and expressed as percentage of the total number of cells.

Statistical analysis

All data are expressed as mean \pm standard error of mean (SEM), and processed using SPSS 16.0 (SPSS Inc., Illinois, Chicago, USA). They were analyzed by one-way analysis of variance

(ANOVA) followed by Dunnett's t-test. $P < 0.05$ was considered statistically significant.

RESULTS

Effect of CNCE on proliferation of A549 cells

CNCE at different doses significantly inhibited the proliferation of A549 cells after 48 h when compared with control group ($p < 0.05$). The inhibitory effect was dose-dependent. These results are shown in Table 1.

Table 1: Inhibitory effect of CNCE on the proliferation of A549 cells

Group	Dose ($\mu\text{g/mL}$)	Absorbance (250 nm)	Inhibition (%)
-ve control	—	0.365 \pm 0.003	—
+ve control	20	0.047 \pm 0.002 ²	91.35 \pm 0.19
CNCE	30	0.189 \pm 0.002 ¹	49.42 \pm 0.22
	60	0.186 \pm 0.004 ¹	52.16 \pm 0.31
	120	0.176 \pm 0.001 ²	53.24 \pm 0.32
	240	0.128 \pm 0.004 ²	62.18 \pm 0.20

-ve = negative, +ve = positive, values are mean \pm SEM (n = 4); ¹ $p < 0.05$, ² $p < 0.01$ relative to control

Effect of CNCE on cell morphology

Under the inverted microscope, the control group of lung cancer A549 cells grew densely, with overlapping clusters, as a long fusiform or polygon, without direction and with clear cellular edges. On the other hand, A549 cells treated with CNCE at concentrations of 10 and 20 $\mu\text{g/mL}$ showed reduction in number of cells, and no obvious morphological changes. When the concentration of CNCE reached 40 $\mu\text{g/mL}$, the cell volume shrank, becoming smaller; and the cells were fewer and separated, with many of floating particles in the culture medium. At concentrations of 160 and 320 $\mu\text{g/mL}$ of CNCE, there were very clear and obvious changes in cell morphology, and evidence of cell disruption and higher cell death.

Growth curve and TD of A549 human lung cancer cells

The growth curve of A549 cells is shown in Figure 1. Cell proliferation was initially slow within the first 48 hours, but thereafter rose rapidly and peaked after 6 days, followed by a decline on the 7th and 8th days. The value of TD obtained was 20.

Inhibitory activity of CNCE

Results from MTT assay showed that CNCE exerted a concentration and time-dependent

inhibitory effects on the growth of human lung cancer A549 cells (Table 2). The inhibitory effect was highest at a concentration of 320 $\mu\text{g/mL}$. In addition, for each CNCE concentration, the inhibitory effect increased with time and became maximum at 96 h.




Figure 1: Growth curve of A549 human lung cancer cells

Table 2: Inhibitory effects of CNCE on A549 lung cancer cell growth

Group	Dose ($\mu\text{g/mL}$)	24h	48h	72h	96h
Control	—	—	—	—	—
CNCE	8	1.28	2.31	2.38	2.48
CNCE	16	5.36	12.34	19.37	23.31
CNCE	32	24.47	27.24	35.24	41.24
CNCE	64	32.24	38.46	49.37	57.33
CNCE	128	42.12	57.14	68.37	75.46
CNCE	256	43.14	58.76	70.64	78.25

Effect of CNCE on A549 cell apoptosis and cell cycle

CNCE at a concentration of 40 $\mu\text{g/mL}$ brought about a CNCEst of cell cycle at G0/G1 phase in 24 and 48 h, and blocked the transformation from G1 phase to S phase (Table 3). With increase in G0 - G1 phase transition time, there was reduction in the number of cells in the S phase. The increases in percentage of cells in the G0/G1 phase, and decreases in cells in S phase, were concentration-dependent and time-dependent ($p < 0.01$).

Table 3: Effect of CNCE on cell cycle distribution of A549 cells at 24 and 48 h

Group	G0/G1 (%)	S (%)	G2/M (%)
Control	53.42 \pm 3.87	42.46 \pm 3.34	8.15 \pm 1.07
CNCE 24 h	58.35 \pm 3.24 [~]	35.54 \pm 4.07 [~]	5.34 \pm 1.46 [~]
CNCE 48 h	63.75 \pm 4.34 [~]	23.84 \pm 5.46 [~]	3.47 \pm 1.28 [~]

Values are mean \pm SEM (n = 8); $p < 0.01$ compared to control

Effect of CNCE on key enzymes of apoptosis and protein expression levels of apoptosis-related genes

Immunocytochemistry results showed that CNCE at 40 µg/mL brought about significant increases in the expressions of apoptosis-related proteins caspase-3, caspase-8 and caspase-9 in 24 h, when compared with control ($p < 0.01$, Table 4).

Table 4: Effect of CNCE on the expressions of caspase-3, caspase-8 and caspase-9

Group	Caspase-3 (%)	Caspase-8 (%)	Caspase-9 (%)
Control	16.24±3.27	18.16±2.24	27.43±3.24
CNCE	52.34±4.42 [*]	48.62±4.43 [*]	58.43±5.42 [*]

Values are expressed as mean ± SEM; $p < 0.01$ compared with control

Visible light microscopy showed Ki-67 protein staining in the nucleus; and BCL-2 and p21 ras protein in the cytoplasm. There were more positive cells in the control group, but at a concentration of 40 µg / mL, there were significant decreases in the number of positive cells ($p < 0.01$). The level of protein expression was significantly higher in the CNCE group than in the control group ($p < 0.01$, Table 5).

Table 5: Effect of CNCE on the expressions of Bcl-2, Ki-67 and P21 ras protein

Group	Bcl-2 (%)	Ki-67 (%)	p21 ras (%)
Control	75.26±6.45	78.43±4.98	54.51±6.12
CNCE	43.21±4.46 [*]	39.23±5.04 [*]	28.24±4.27 [*]

^{*} $P < 0.01$ compared with control group

DISCUSSION

The results obtained in this study show that CNCE has significant time- and concentration-dependent inhibitory effects on the proliferation of A549 cells. The extract also exerted strong apoptotic effects on the A549 cells, as seen from the morphological changes in the cells. The dynamics of tumor cell proliferation are due to loss of control of cell cycle, resulting in uncontrolled G1 / S and G2 / M transformations. The cells were mainly in the active DNA synthesis S phase, which led to abnormal cell proliferation [5]. Thus blockage of the G1/S phase and G2/M phase can effectively control the tumor cell cycle and inhibit tumor proliferation.

In this study, results from cell cycle analysis showed that CNCE reduced the number of A549 human lung cancer cells in G1, S and G2/M phases, leading to stagnation of the cell cycle and restraining of cell proliferation. Increased

concentrations of aqueous fractions of CNCE brought about enhanced effects on cell cycle aCNCEst. This indicates that CNCE causes lung cancer cell cycle aCNCEst by regulation and inhibition of tumor cell mitosis.

Ki-67 antigen is a more positive marker of nuclear proliferation, and Ki-67 may be used as a biomarker for determining high risk individuals in pre-cancer population [7-9]. Results from Ki-67 antigen assay showed that CNCE significantly decreased the expression of Ki-67 in A549 cells. This implies that a large number of the A549 cancer cells were in the stationary phase of cell cycle. In effect, CNCE may contain active principles that regulate the gene which inhibits cancer cell mitosis, thereby inhibiting the growth of tumor cells.

P21 ras cancer gene is involved in transmission of intracellular information which regulates the cell cycle, and it is the "initiation factor" in the occurrence of tumors [10-13]. The expression of p21ras antigen was also significantly decreased in the CNCE-treated A549 cells, in a time- and concentration-dependent manner, when compared with untreated controls. Again, this is an indication that most of the cancer cells were in the S phase of the cell proliferation cycle. This supports the presence of anti-proliferative and mitosis-inhibiting agents in CNCE.

Caspases are proteolytic enzymes which constitute key components of the apoptosis system. They are a family of cysteine proteases involved in regulation of apoptosis, which is considered to be a key to cancer cell death [14-17]. In this study, CNCE treatment led to significantly lowered expression of Bcl-2, and significant increases in the expression of caspase-9. These results demonstrate that the anti-proliferative effects CNCE also involve the mitochondrial apoptotic pathway.

Caspase-3 is a pro-apoptosis caspase. Its expression in the CNCE-treated A549 human lung cancer cells was significantly higher than in the untreated control cells, indicating that CNCE promotes apoptosis in these cells. Thus, CNCE induced apoptosis by inhibiting the signal transduction pathway enzymes to achieve elevated expressions of the pro-apoptosis caspases 3, 8 and 9; as well as the apoptosis-related gene Bcl-2.

CONCLUSION

The findings of this study reveal that CNCE has significant effect on the inhibition of proliferation and induction of apoptosis in human lung cancer

A549 cells. These findings suggest that the plant material might be a potent source of drugs for the treatment of lung cancer patients.

DECLARATIONS

Acknowledgement

The authors thank Professor Ping Wang, Department of Pharmacy of Zhongshan University for his assistance with data analysis.

Conflict of interest

The authors declare that there is no conflict of interest associated with this work.

Contribution of authors

We declare that this work was done by the authors named in this article and all liabilities pertaining to claims relating to the content of this article will be borne by the authors. Zhi-hui Zhang designed all the experiment and revised the paper. Pei Li performed the experiment and wrote the manuscript.

REFERENCES

1. Wagland R, Brindle L, Ewings S, James E, Moore M, Rivas C. Promoting help-seeking in response to symptoms amongst primary care patients at high risk of lung cancer. A mixed method study. *PLoS One* 2016; 11: e0165677.
2. Biaoxue R, Hua L, Wenlong G, Shuanying Y. Increased serum amyloid A as potential diagnostic marker for lung cancer: A meta-analysis based on nine studies. *BMC Cancer* 2016; 16: 836-838.
3. Amoori N, Mirzaei M, Cheraghi M. Incidence of cancers in Kuzestan province of Iran: Trend from 2004 to 2008. *Asian Pac J Cancer Prev* 2014; 15: 8345-8349.
4. Bunn Jr PA, Thatcher N. Systemic treatment for advanced (stage IIIb/IV) non-small cell lung cancer: more treatment options; more things to consider. Conclusion. *Oncologist* 2008; 37: 46-48.
5. Thorn CF, Oshiro C, Marsh S, Hernandez-Boussard T, McLeod H, Klein TE, Altman RB. Doxorubicin pathways: pharmacodynamics and adverse effects. *Pharmacogenet Genomics* 2011; 21: 440-446.
6. Hanusova V, Bousova I, Skalova L. Possibilities to increase the effectiveness of doxorubicin in cancer cells killing. *Drug Metab Rev* 2011; 43: 540-557.
7. Kurniawan NA, Chaudhuri PK, Lim CT. Concentric Gel System to Study the Biophysical Role of Matrix Microenvironment on 3D Cell Migration. *J Vis Exp* 2015; 22: e52735.
8. Alfano M, Nebuloni M, Allevi R, Zerbi P, Longhi E, Lucianò R, Locatelli I. Linearized texture of three-dimensional extracellular matrix is mandatory for bladder cancer cell invasion. *Sci Rep* 2016; 6: 36128-36130.
9. Liu D, Guo H, Li Y, Xu X, Yang K, Bai Y. Association between polymorphisms in the promoter regions of matrix metalloproteinases (MMPs) and risk of cancer metastasis: A meta-analysis. *PLoS One* 2012; 7: e31251.
10. Huang TH, Chiu YH, Chan YL, Chiu YH, Wang H, Huang KC. Prophylactic administration of fucoidan represses cancer metastasis by inhibiting vascular endothelial growth factor (VEGF) and matrix metalloproteinases (MMPs) in lewis tumor-bearing mice. *Mar Drugs* 2015; 13: 1882-1900.
11. Hassan ZK, Elamin MH, Daghestani MH, Omer SA, Al-Olayan EM, Elobeid MA. Oleuropein induces anti-metastatic effects in breast cancer. *Asian Pac J Cancer Prev* 2012; 13: 4555-4559.
12. Ma H, Liu B, Feng D, Xie H, Li R, Yuchi Y, Wang H, Wang J. Pinus massoniana bark extract selectively induces apoptosis in human hepatoma cells, possibly through caspase-dependent pathways. *Int J Mol Med* 2010; 25: 751-759.
13. Cui Y, Xie H, Wang J. Potential biomedical properties of Pinus massoniana bark extract. *Phytother Res* 2005; 19: 34-38.
14. Liu J, Bai J, Jiang G, Li X, Wang J, Wu D, Owusu L, Zhang E. Anti-tumor effect of Pinus massoniana bark proanthocyanidins on ovarian cancer through induction of cell apoptosis and inhibition of cell migration. *PLoS One* 2015; 10: e0142157.
15. Cai Q, Lin J, Wei L, Zhang L, Wang L, Zhan Y, Zeng J, Xu W, Shen A. Hedyotis diffusa Willd inhibits colorectal cancer growth in vivo via inhibition of STAT3 signaling pathway. *Int J Mol Sci* 2012; 13: 6117-6128.
16. Zhang JH, Feng DR, Ma HL, Liu B, Wang HB, Xie H, Li RD. Antitumor effects of Pinus massoniana bark extract in murine sarcoma S180 both in vitro and in vivo. *Am J Chin Med* 2012; 40: 861-875.
17. Wu DC, Li S, Yang DQ, Cui YY. Effects of Pinus massoniana bark extract on the adhesion and migration capabilities of HeLa cells. *Fitoterapia* 2011; 82: 1202-1205.