

Original Research Article

Determination of radical scavenging activities of some pyrimidine derivatives

Oluwole S Aremu^{1*}, Lebogang Katata-Seru¹, Olukayode O Aremu², Constance R Sewani-Rusike², Neil Koobanally³

¹Department of Chemistry, Faculty of Natural and Agriculture Science, North West University, Mafikeng campus, Private bag X2046, Mmabatho 2745, ²Department of Human Biology, Faculty of Health Sciences, Walter Sisulu University, Mthatha 5117, South Africa; ³School of Chemistry and Physics, Westville campus, University of KwaZulu-Natal, Durban 4000,

*For correspondence: **Email:** 23859539@nwu.ac.za; **Tel:** +27-842905180

Sent for review: 15 February 2018

Revised accepted: 16 October 2019

Abstract

Purpose: To synthesize some pyrimidine derivatives and investigate their radical scavenging activities.

Methods: A series of newly pyranopyrimidines derivatives and dithiopyridopyrimidinediones were synthesized by condensation of barbituric acid, malononitrile and different substituted benzaldehydes reacted with 1,4-Diazabicyclo[2.2.2] octane (DABCO) as a base. Compounds P1-7 (series 1), S1-11 (series 2) Scheme 1 and 6-amino-2-thiouracil with aromatic aldehydes in glacial acetic acid under reflux J1-13 (series 3) Scheme 2. ¹H & ¹³C NMR, CHN, GC-MS and IR were used to characterize the compounds and were also screened for radical scavenging activity using 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging activity ($IC_{50} = 50 \mu\text{g/ml}$).

Results: The potency of radical scavenging activity was ranked as series 1 > series 3 > series 2. Compounds P3, J4, S10, P5, P7 with inhibitory concentration at 50 % level (IC_{50}) of 12, 40, 48, 50, and 54 $\mu\text{g/ml}$, respectively, showed radical scavenging activity equal or more potent than the standard antioxidant, ascorbic acid ($IC_{50} = 50 \mu\text{g/ml}$).

Conclusion: Series of newly pyranopyrimidines and dithiopyridopyrimidinediones derivatives have been successfully synthesized, and they demonstrate good radical scavenging activity.

Keywords: Pyranopyrimidine, Dipyrimidines, Anti-oxidant, DPPH, Ascorbic acid, Radical scavenging

This is an Open Access article that uses a fund-ing model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited.

Tropical Journal of Pharmaceutical Research is indexed by Science Citation Index (SciSearch), Scopus, International Pharmaceutical Abstract, Chemical Abstracts, Embase, Index Copernicus, EBSCO, African Index Medicus, JournalSeek, Journal Citation Reports/Science Edition, Directory of Open Access Journals (DOAJ), African Journal Online, Bioline International, Open-J-Gate and Pharmacy Abstracts

INTRODUCTION

Pyrimidine is a six-membered heterocyclic aromatic organic compound that contains two nitrogen atoms at positions 1 and 3. It is known to possess some pharmacological activities, such as antiviral, anticancer [1-3], antimicrobial [4-6], anti-inflammatory [7] and antimalarial [8,9]. Molecules containing a pyrimidines scaffold is

known to have pharmacological properties [10]. Some heterocyclic derivatives of pyrimidines such as pyranopyrimidines and dithiopyridopyrimidinediones derivatives contain pyrimidine, pyrano, pyrido and aryl moieties [11].

The pyranopyrimidines and dithiopyridopyrimidinediones derivatives have been previously

synthesized by various methods as previously reported. [12].

Reactive oxygen species (ROS), such as superoxide radical anions, hydrogen peroxide and hydroxyl radical are free radicals continuously produced during normal metabolic pathways and in pathology [13]. Excessive production of ROS is deleterious to cellular proteins and lipids due to oxidative damage. The body's endogenous antioxidant system acts as a ROS scavenger to protect cells and macromolecules from oxidative damage. Disturbances of anti-oxidant status are seen in diabetes, hypertension, obesity, infertility, Alzheimer's disease, Parkinson's disease and cancer [14]. As such, supplementing the endogenous antioxidant system is being investigated as a possible therapy in prevention, as well as limiting progression and development of complications, of chronic diseases. In view of this, we report on the investigation of the antioxidant properties of the synthesized pyrimidine compounds using 1, 1-diphenyl-2-picrylhydrazyl (DPPH) and free radical scavenging assay.

EXPERIMENTAL

Materials

All chemicals and solvents used in this study are of analytical grade and were purchased from Sigma Aldrich via Capital Laboratories, South Africa. All organic solvents were redistilled and dried according to standard procedures. Nuclear magnetic resonance (NMR) spectra were recorded using Bruker Avance III 400 MHz spectrometer at room temperature, with chemical shifts (δ) recorded against the internal standard, tetramethylsilane (TMS). IR spectra were recorded on a Perkin Elmer Spectrum 100 ATR-FTIR spectrometer. For GC-MS analyses, the samples were analyzed on an Agilent GC-MSD apparatus equipped with DB-5SIL MS (30m \times 0.25mm i.d., 0.25 μ m film thickness) fused-silica capillary column. Helium (at 2mL min⁻¹) was used as a carrier gas. MS was operated in EI mode at 70 eV. Optical rotation was recorded using a polarimeter (Model 341, Perkin Elmer Inc, USA). Melting points were recorded on an Ernst Leitz Wetzlar microhot stage melting point apparatus.

General procedure for the synthesis of pyranopyrimidione derivatives (P1-7, S1-11)

Reported procedure [12]. was used in synthesizing this library of compounds by reacting barbituric acid, benzaldehyde

derivatives (ArCHO), malononitrile (1.0 mmol each) and 10% DABCO in a clean 50ml round bottom flask containing 20ml of aqueous ethanol (Scheme 1). The reaction mixture was stirred for a period of 60 minutes. The progress of the reaction was monitored by TLC. For most of this synthesis, the products were filtered, recrystallized in ethanol and dried under vacuum pressure, with good yields of the final products.

General procedure for the synthesis of dithiopyrimidinediones derivatives (J1-13)

A solution of 6-amino-2-thiouracil 4 (3 mmol) in glacial acetic acid (15 ml) and 0.5 equiv. of the appropriate aromatic aldehyde was heated under reflux for 4 hours. The reaction mixture was diluted with water and then allowed to cool to room temperature. The crude product was collected and recrystallized from a suitable solvent. Details have been reported in previously published articles from our group [15].

Evaluation of antioxidant properties

The free radical scavenging activity of the synthesized compounds was determined by 1, 1-diphenyl-2-picrylhydrazyl (DPPH) method, as described by Brand William et al (1995) [16]. In this protocol, DPPH• acts as a stable free radical with a purple colour. In the presence of an antioxidant that donates an electron, the purple colour of DPPH• decays. This change can be monitored spectrophotometrically and used to assess radical scavenging activity, which equates to antioxidant activity of compounds. For this study, compounds were dissolved in a minimum volume of DMSO and diluted in ethanol to 1mg/ml. To 1ml of compound or ascorbic acid, the standard (500, 250, 125 and 62.5 μ g/ml) in a test tube, 3 ml of 0.5 mM ethanolic solution of DPPH was added. The mixture was vortexed vigorously and incubated for 30 minutes in the dark at room temperature. Absorbance was read spectrophotometrically at 517 nm and inhibition of DPPH free-radical scavenging activity (D) was computed as in Eq 1.

$$D \{(Ac As)/Ac\}100 \dots\dots\dots (1)$$

where Ac and As are the absorbance of control and test sample, respectively.

RESULTS

Chemistry


Various pyranopyrimidiones derivatives P1-7 and S1-11) (Scheme 1) have been reported in the

literature [12]. Scheme 1 showed the good yields of the synthesized final products, whereas, Scheme 2 revealed dithiopyridopyrimidines derivatives (J1-13) synthesized, as reported in the literature [15]. Hence, the formation of the title compounds was confirmed and further established by ^{13}C NMR and mass spectroscopic studies, which agree with the molecular formula.


Antioxidant activity

Radical scavenging activity of compounds is shown in Tables 1 to 4. Better radical scavenging activity was observed for series 1 than for series 2 compounds, with three (27%) showing no activity at the lowest concentration of 62.5 $\mu\text{g/ml}$, contrary to series 2 that showed three (27%) with no activity at all and five (45%) showing no activity at the lowest concentration of 62.5 $\mu\text{g/ml}$. Although series 3 compounds showed radical scavenging activity at all concentrations, the IC_{50} values were very high compared to that in series 1. Thus, potency in radical scavenging activity is ranked as series 1 > series 3 > series 2 (Table 1). Overall, when compared to ascorbic acid (IC_{50}

= 50 $\mu\text{g/ml}$), a known and potent antioxidant compound, five compounds have equal or better radical scavenging activity: 3, C4, B10, 5, 7 with IC_{50} of 12, 40, 48, 50, and 54 $\mu\text{g/ml}$ respectively (Table 2 and Figure 1).



Scheme 1: Synthesis of pyranopyrimidiones (P1-7, S1-12) (oxygenated and halogenated compounds)




Scheme 2: Synthesis of dithiopyridopyrimidiones derivatives (J1-13)

Table 1: Radical scavenging activity of synthesized compounds

Compound	DPPH inhibition (%)					IC_{50} ($\mu\text{g/ml}$)
	500 $\mu\text{g/ml}$	250 $\mu\text{g/ml}$	125 $\mu\text{g/ml}$	62.5 $\mu\text{g/ml}$		
P1	42	28	18	11	595	
P2	66	52	32	17	240	
P3	79	79	78	78	12	
P4	47	31	17	-	532	
P5	78	77	73	62	50	
P6	87	54	26	17	230	
P7	81	82	77	57	54	
S1	60	45	28	16	225	
S2	42	25	18	-	595	
S3	28	19	12	-	893	
S4	50	32	17	-	500	
S5	-	-	-	-	1000	
S6	45	26	-	-	555	
S7	-	-	-	-	1000	
S8	-	-	-	-	1000	
S9	25	11	-	-	893	
S10	28	28	28	18	900	
S11	88	88	77	65	48	
J1	87	65	47	33	133	
J2	91	89	69	51	61	
J3	57	44	34	29	440	
J4	91	91	89	78	40	
J5	65	50	35	28	250	
J6	82	69	46	39	136	
J7	54	38	31	26	460	
J8	59	46	32	25	420	
J9	52	44	34	27	480	
J10	63	46	33	24	270	
J11	50	40	35	27	500	
J12	52	39	30	23	480	
JP13	64	44	33	21	280	
Ascorbic acid	96	96	92	65	50	

Table 2: All compounds DPPH inhibition activity ranked (highest to lowest) comparing with AA

Compound	DPPH inhibition (%)				
	500 µg/ml	250 µg/ml	125 µg/ml	62.5 µg/ml	IC ₅₀ (µg/ml)
P3	79	79	78	78	12
J4	91	91	89	78	40
S10	88	88	77	65	48
P5	78	77	73	62	50
Ascorbic acid	96	96	92	65	50
P7	81	82	77	57	54
J2	91	89	69	51	61
J1	87	65	47	33	133
J6	82	69	46	39	136
S1	60	45	28	16	225
P4	57	54	38	11	230
P2	66	52	32	17	240
J5	65	50	35	28	250
J8	68	50	41	33	250
J4	56	41	28	17	446
J10	53	44	35	26	472
S4	50	32	17	-	500
J11	50	40	35	27	500
B7	45	26	-	-	555
P1	42	28	18	11	595
S2	42	25	18	-	595
J9	42	39	30	27	595
P6	41	22	12	-	610
S3	28	19	12	-	893
S9	25	11	-	-	893
S5	-	-	-	-	1000
S7	-	-	-	-	1000
S8	-	-	-	-	1000

**Figure 1:** Potential inhibition capacities of the most potent synthesized compounds (*rank order*: P3 > J4 > S10 > P5 > P7)

DISCUSSION

A series of new pyranopyrimidiones and dithiopyridopyrimidiones derivatives were synthesized and evaluated for radical scavenging activity.

Pyrimidines, also known as the breakdown products of uric acid, play a vital role in the medicinal chemistry, due to their structures and composition, and act as the key templates associated with the development of different therapeutic agents [16,20]. Several literatures have reported on antioxidant activities of some pyrimidine derivatives [17,18]. In the present study, five compounds have equal or better

radical scavenging activity when compared to ascorbic acid (IC₅₀ = 50µg/ml), a known and potent antioxidant compound. This may be due to the available NH and SH group present in the synthesized pyrimidines [19].

CONCLUSION

A series of pyranopyrimidiones and dithiopyridopyrimidiones derivatives have successfully been synthesized and they exhibit radical scavenging activity in the rank order of series 1 > series 3 > series 2.

DECLARATIONS

Acknowledgement

The authors acknowledge the School of Chemistry and Physics, University of KwaZulu-Natal, the Department of Chemistry at North West University, Mafikeng Campus, and Walter Sisulu University Research Department for financial support during this work.

Conflict of interest

No conflict of interest is associated with this work.

Contribution of authors

We declare that this work was done by the author(s) mentioned in this article and all liabilities pertaining to claims relating to the content of this article will be borne by the authors. Oluwole S. Aremu and Neil Koorbanally devised and designed the study, Oluwole S. Aremu carried out the laboratory work on the chemistry aspect, Olukayode O. Aremu and Constance R. Sewani-Rusike investigated the antioxidant properties of the synthesized pyrimidines, Lebogang Katata-Seru edited the raw data. The manuscript was proof-read by all the authors and approved for publication.

Open Access

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited.

REFERENCES

- Okesli A, Khosla C, Bassik MC. Human pyrimidine nucleotide biosynthesis as a target for antiviral chemotherapy. *Curr. Opin. Biotechnol.* 2017; 48: 127-134.
- Mohan SB, Ravi Kumar BV, Dinda SC, Naik D, Prabhu Seenivasan S, Kumar V, Rana DN, Brahmshatriya PS. Microwave assisted synthesis, molecular docking and antitubercular activity of 1, 2, 3, 4-tetrahydro-pyrimidin-5-carbonitrile derivatives. *Bioorg. Med. Chem. Lett.* 2010; 22(24): 7539-7542.
- El-Gazzar AR, Hussein HA, Hafez HN. Synthesis and biological evaluation of thieno [2, 3-d] pyrimidine derivatives for anti-inflammatory, analgesic and ulcerogenic activity. *Acta Pharm* 2007; 57(4): 395-411.
- Sherif AF, Hayam MA, Heba A. Synthesis and biological evaluation of some novel polysubstituted pyrimidine derivatives as potential antimicrobial and anticancer agents. *Arch. Pharm.* 2009; 342(5): 299-310.
- Hilmy KM, Khalifa MM, Hawata MA, Keshk RM, el-Torgman AA. Synthesis of new pyrrole [2,3-d] pyrimidine derivatives as antibacterial and antifungal agents. *Eur. J. Med. Chem.* 2010; 45(11): 5243-5250.
- Verbitskiy EV, Baskakova SA, Kungurov NV, Kravchenko MA, Skornyakov SN, Pervova MG, Rusinov GL, Chupakhin ON, Charushin VN. Synthesis and biological evaluation of novel 5-aryl-4-(5-nitrofuranyl)-pyrimidines as potential anti-bacterial agents. *Bioorg. Med. Chem. Lett* 2017; 27(13): 3003-3006.
- Nofal ZM, Fahmy HH, Zarea ES, El-Eraky W. Synthesis of new pyrimidine derivatives with evaluation of their anti-inflammatory and analgesic activities. *Acta. Pol. Pharm.* 2009; 68(4): 507-517.
- Azeredo LFS, Coutinho JP, Jabor VA, Feliciano PR, Nonato MC, Kaiser CR, Menezes CMS, Hammes AS, Caffarena ER, Hoelz LV. Evaluation of 7-arylaminopyrazolo [1, 5-a] pyrimidines as anti-*Plasmodium falciparum*, antimalarial, and Pf-dihydroorotate dehydrogenase inhibitors. *Eur. J. Med. Chem.* 2017; 126: 72-83.
- Sun L, Jie W, Zhang L, Luo M, Sun D. Synthesis and antifungal activities of some novel pyrimidine derivatives. *Molecules* 2011; 16: 5618-5628.
- Mohamed MS, Awad SM, Sayed AI. Synthesis of certain pyrimidine derivatives as antimicrobial agents and anti-inflammatory agents. *Molecules* 2010; 15(3): 1882-1890.
- Mohana KN, Kumar BNP, Mallesha L. Synthesis and biological activity of some pyrimidine derivatives, *Drug Invent Today.* 2013; 5: 216-22.
- Oluwole S Aremu, Gopaul K, Kadam P, Singh M, Mocktar C, Singh PA, Koorbanally N. Synthesis, Characterization, Anticancer and Antibacterial Activity of Some Novel Pyrano [2, 3-d] pyrimidinone Carbonitrile Derivatives. *Anticancer Agents Med. Chem.* 2017, 17 (5), 719-725.
- Castro L, Freeman BA. Reactive oxygen species in human health and disease. *Nutr. J.* 2001; 17(2), 161-165.
- Halliwell B. Free radicals, antioxidants, and human disease: curiosity, cause, or consequence? *Lancet* 1994; 344: 721-724.
- Aremu OS, Singh M, Singh P, Koorbanally NA. Facile one-pot synthesis of novel Pyridodipyrimidine derivatives: Their Anticancer activity against HeLa human cervical cancer cell line. *Bioorg. Chem Let* 2015 (In press).
- Brand-Williams W, Cuvelier ME, Berset C and Lebensm. W. Use of a free radical method to evaluate antioxidant activity. *Technol.* 1995; 28: 25-32
- Himaja M, Rout PK, Karigar AA, Poppy D, Munirajashekhhar D. Synthesis and antioxidant activity of 4,6 - diphenylpyrimidine- 2- amine & its derivatives. *Univ J Pharm* 2012; 01(01): 72-77.
- Bano T, Kumar N, Dudhe R. Free radical scavenging properties of pyrimidine derivatives. *Org Med Chem Lett* 2012; 2(1): 34.
- Dansena H, Dhongade HJ, Chandrakar K. Pharmacological Potentials of Pyrimidine Derivative: A Review. *Asian J Pharm Clin Res* 2015; 8(4): 171-177.
- Ghomi AS, Ali M. An efficient route to the synthesis of pyrimidine-2-ones under ultrasound irradiation. *Dig J Nanomater Biostruct* 2010; 5(2): 303-306.