

Eclampsia and abnormal QTc

*¹S. A. Isezuo and ²B. A. Ekele

Departments of Medicine¹ and ²Obstetrics and Gynaecology
Usman Danfodiyo University Teaching Hospital
Sokoto, Nigeria.

E-mail: sisezuo@skannet.com

Summary

Background: Abnormalities of calcium and magnesium metabolism are known risk factors for QT interval prolongation and have been reported in eclampsia.

Objective: To determine if eclampsia is associated with QT abnormalities

Methods: In a prospective study, the QT interval corrected for heart rate (QTc), serum calcium, magnesium and potassium were compared among 30 intrapartum eclamptics and 30 age, parity and gestational age-matched women with uncomplicated pregnancy

Results: Primigravidae made up 93.3% of patients. Mean age was 19.5 ± 4.2 years Blood pressure was significantly higher among patients than controls (163.0 ± 34.7mmHg versus 120.4 ± 18.6mmHg systolic, p<0.05) and (104.7 ± 15.2mmHg versus 79.6 ± 10.7mmHg, p<0.05). Heart rate ranged from 76 to 163 beats per minute (bpm) and 65 to 112bpm among patients and controls respectively. The corresponding QTc were 390-572 and 390-460 respectively. Compared to the controls, patients had significantly higher mean heart rate (121.1 ± 24.9bpm versus 89.3 ± 12.9bpm, p<0.05) and higher QTc (470.4 ± 42.4 versus 436.3 ± 17.7; p<0.05). Compared to the controls, the eclamptics also had higher frequency of sinus tachycardia (90% versus, 13.3%) Odd Ratio (OR) = 29.57; 95% Confidence Interval (CI) = 5.87-174.08, p=0.00), abnormal QTc (46.7% versus 6.6%, OR = 9.2; 95% CI = 1.61-68.48, p=0.01) and T-axis deviation (26.6% versus 3.3%, OR=10.55, 95% CI=1.2-244.3, p=0.03)

Serum calcium level was significantly lower among patients than controls (2.0 ± 0.4mmol/L versus 2.3 ± 0.2mmol/L, p<0.05). The eclamptics had higher frequency of hypocalcaemia than the controls (40% versus 6.6%; OR = 14, 95% CI = 1.58 - 316.9, p=0.01). Serum magnesium, potassium, urea and creatinine levels were similar between the two groups.

Conclusion: Eclampsia is associated with abnormal QTc that may be related to maternal hypocalcaemia. Correction of electrolyte abnormalities and cautions about agents that have potential for QT prolongation are vital in the care of eclamptics.

Keywords: Eclampsia, Abnormal QTc, Hypocalcaemia.

Résumé

Introduction: Les anomalies du métabolisme du cal-

cium et magnésium sont des facteurs de risque connus pour la prolongation intervalle QT et ont été rapportée en éclampsie.

Objectif: Déterminer si l'éclampsie est associée avec des anomalies QT.

Methodes: D'après une étude en perspective, l'intervalle QT, corrigé pour la pulsation du Coeur (Otc) sérum calcium, magnésium et potassium ont été comparé parmi 30 éclamptiques intrapartum et âgé de 30, parité et groupe d'âge des femmes gestationelle avec grossesse sans complications.

Résultats: Primigravidae composé de 93,3% de patients, Age moyen était 19,5±4,2 ans. Tension artérielle était remarquablement élevée parmi des patients plus que chez des contrôle (163,0±37,7mmHG contre 120,4±18,6mmHg systolique, P<0,05) et (104,7±15,2mmHg contre 79,6±10,7mmHg, P<0,05). Pulsation du (Coeur varie de 76 à 163 par minute (bpm) et 65 à 1/2 bpm parmi des patients et des contrôles respectivement. Les QTc correspondant étaient 390 - 572 et 390 - 460 respectivement. Par rapport à des contrôles, des patients avaient la pulsation du Coeur moyenne remarquablement élevée (121,1 ± 24,9 bpm contre 89,3 ± 12,9bpm, P<0,05) et QTS plus élevé (470,4 ± 42,4, contre 436,3 ± 17,7, P <0,05) par rapport aux contrôles, l'éclamptique aussi avait la fréquence plus élevée de la tachycardie sinusale (90% contre 13,3; proportion bizarre (ou) = 29,57, 95% intervalle de confiance (CI) = 5,87 - 174,08, P=0,00) QTc anormal (46,7% contre 6,6%, OU=10,55, 95% CI = 1,2-244,3, P=0,03.

Niveau du sérum calcium était remarquablement en baisse chez des patients plus que chez des contrôles (2,0±0,4mmol/L contre 2,3 ±0,2mmol/L: P<0,05). L'éclamptique avait une fréquence plus élevée d'hypocalcémie plus que chez des contrôles (40% contrôle 6,6% OU+14,95% CI= 1,58-3169, P=0,01). Sérum magnésium potassium, urée et niveau créatine était pareil entre les deux groupes.

Conclusion: Eclampsie est liée avec QTc anormal qui pourrait être en rapport avec l'hypocalcémie maternelle. La correction d'anormalité électrolyte et prudence en matière des agents qui ont un potentiel pour la prolongation QT sont majeurs dans la prise en charge des éclamptiques.

Introduction

Eclampsia is a syndrome characterized by generalized convulsion, elevated blood pressure and proteinuria

*Correspondence

with or without oedema occurring after 20 weeks of gestation. Its occurrence varies from 1 in 100 to 1 in 1700 in developing countries where it is also an important cause of maternal mortality^{1,2}. Its pathogenesis is still incompletely understood. Abnormalities of calcium and magnesium metabolism, particularly, hypocalcaemia probably play important roles in the pathogenesis of pregnancy induced hypertension and eclampsia^{3,4}. Thus, calcium supplementation has been found useful in the prevention of pre-eclampsia⁵, while treatment and prevention of eclampsia with magnesium sulphate has been established⁶⁻⁸. QT interval represents the total time from the onset of ventricular depolarisation to the completion of repolarisation. Prolonged QT interval is a disorder of cardiac repolarisation caused by alteration in the transmembrane cation contents. This reduces net ventricular repolarising current, thereby amplifying the intrinsic spatial dispersion of repolarisation, a prime step for development of re-entry phenomenon. The net results are QT interval prolongation, abnormal T-waves, generation of early after-potential and development of arrhythmias^{9,10}.

Hypocalcaemia, a reported occurrence in eclampsia^{3,4} is an independent risk factor for QT prolongation¹¹. Some drugs including chlorpromazine, an antipsychotic/anticonvulsant agent and quinolones used in treating infection in eclampsia have also been shown to prolong QT^{12,13}. The main significance of QT interval prolongation is the development of *torsade de pointes*, a potentially lethal ventricular polymorphic tachyarrhythmia that may cause sudden cardiac death^{9,10}. Cardiac arrest and heart failure have been recently reported in about 0.1% of patients with eclampsia⁶.

In spite of the presence of the potential risk factors for QT prolongation in eclampsia, information on QT in this condition is lacking. In addition to its therapeutic implications, such information may provide insight into some of the causes of high mortality in eclampsia. We therefore determined the QT among eclamptic patients.

Methodology

Thirty consecutive patients with intra-partum eclampsia admitted into the labour ward of Usman Danfodiyo University Teaching Hospital, Sokoto, Nigeria between August 2001 and July 2002 served as the subjects. Information about age, tribe, parity, gestational age, history of diabetes and medications used prior to admission were obtained. Blood pressure was measured using standard mercury sphygmomanometre (Accosson), size 150cm length and 35cm breadth. Three readings were taken after about 15 minutes of rest. The average of the last two readings was taken as the blood pressure. Systolic blood pressure ≥ 140 mmHg and/or diastolic blood pressure ≥ 90 mmHg were required to make a diagnosis of systemic hypertension¹⁴. Clinical evaluation was carried out to exclude structural or primary heart disease. The diagnosis of intra-partum eclampsia was based on the presence of generalised convulsions, systemic hypertension, proteinuria of at least 2+ by dipstick and established labour

with or without oedema.

A standard 12-lead electrocardiogram (ECG) was recorded during the intrapartum period using automated ECG machine (EKG 2000, Bionet Co. Ltd). Lead placements were as recommended by the American Heart Association¹⁵. ECG paper speed was set at 25mm per second while calibration was set at 10mm per mV. About 5 cycles were recorded per ECG lead. In order to prevent observer's error, the same investigating Cardiologist verified all measurements manually. The leads with the most distinct Q and T waves were utilised for manual measurements. QT interval was considered as the total time from the beginning of Q wave to the end of T wave. It was corrected for heart rate using Bazett's formula: $QTc = QT / \sqrt{RR}$. $QTc > 460$ was considered prolonged. T-axis outside 0° - 90° was considered as T-axis deviation.

About 10mls of blood was withdrawn for the determination of serum calcium, magnesium and glucose using Banguski, Xyldye Blue calorimetric and glucose oxidase tests respectively. Serum potassium, urea and creatinine were also determined. Serum calcium < 2.2 mmol/L, serum magnesium < 0.8 mmol/L and serum potassium < 3.5 mmol/L constituted hypocalcaemia, hypomagnesium and hypokalaemia respectively.

Patients on admission prior to hospitalisation were excluded from the study. This is because records of such medications were usually not available to determine whether they could potentially prolong QT interval. Patients on medication that could cause QT prolongation (chlorpromazine, quinine, halofantrine, artemeter and quinolones) prior to ECG recording as well as those with diabetes mellitus and structural heart disease were also excluded.

Age, parity, tribe and gestational age-matched healthy individuals with normal pregnancy in established labour served as control.

Statistical analysis

Data entry and analysis were done using Epi-Info software. Values are presented as mean \pm SD. Continuous variables were compared using Student's t-test

Table 1 Baseline clinical and personal data of patients and controls

Characteristics		Patients N = 30	Controls N = 30
Tribes (N (%))	Hausa/Fulani	24(80.0)	25(83.3)
	Ibo	1(3.3)	2(6.7)
	Yoruba	2(6.7)	1(3.3)
	Others	3(10.0)	2(6.7)
Parity N (%)	Primigravida	28(93.3)	30(100.0)
	Multigravida	2(6.7)	0(0)
Gestational age > 38 weeks N(%)		30(100.0)	30(100.0)
Age (years)		19.5 \pm 4.2	20.2 \pm 2.7
*Systolic blood pressure (mmHg)		163.0 \pm 24.7	124.8 \pm 18.1
*Diastolic blood pressure (mmHg)		104.7 \pm 15.02	79.6 \pm 10.7

* $p < 0.05$




Fig. 1. A 22 year old eclamptic with abnormal QTc (QTc = 500)

while discrete variables were compared using chi-square test with Yate's correction. Fisher – Exact test was applied where appropriate. A p-value <0.05 was considered statistically significant.

Results

The baseline personal and clinical data of the study group controls are shown in Table 1. 80% of the patients belonged to the Hausa and Fulani tribes of Nigeria. The primigravidae made up 93.3% of patients.

The biochemical parameters of patients and controls are compared in Table 2. The mean serum potassium, urea and creatinine were similar between the two groups. Serum calcium levels was significantly lower among patients than controls (2.0 ± 0.4 mmol/L, range 1.8–2.6 mmol/L) versus (2.3 ± 0.2 mmol/L, range 2.1–2.8 mmol/L) ($t=3.5$

$df=58$ $p<0.05$). The eclamptics had higher frequency of hypocalcaemia than the controls (40% versus 6.6%; OR =14, 95% CI = 1.58–316.68, $p=0.01$). The mean serum magnesium level did not suffer significantly among patients and controls (0.9 ± 0.2 mmol/L, range: 0.7 – 1.7 mmol/L) versus (0.85 ± 0.3 mmol/L, range 0.5–1.8 mmol/L) ($t=0.8$ $df=58$, $p>0.05$).

The ECG parameters of patients and controls are shown in Table 3. Heart rate ranged from 76 to 163 beats per minute (bpm) and 65 to 112 bpm among patients and controls respectively. The corrected QT interval (QTc)

Table 2 Comparison of the biochemical parameters of patients and controls

Parameters	Patients N = 30	Controls N = 30
*Serum calcium (mmol/L)	2.0 ± 0.04	2.3 ± 0.2
Serum magnesium (mmol/L)	0.9 ± 0.2	0.9 ± 0.3
Serum potassium (mmol/L)	4.2 ± 0.7	4.4 ± 0.7
Serum urea (mmol/L)	4.6 ± 1.6	4.0 ± 1.1
Serum creatinine (mg%)	1.0 ± 0.1	0.9 ± 0.1
*Hypocalcaemia N (%)	12(40)	2 (6.7)
Hypomagnesaemia N (%)	2(6.7)	2(6.7)

*p <0.05.

Table 3 Comparison of the ECG parameters of patients and controls

Parameters	Patients N = 30	Controls N = 30
*Heart rate (beat/min)	121.1 ± 24.9	89.3 ± 12.9
*RR interval (ms)	13.1 ± 2.9	17.6 ± 3.3
QT interval (ms)	336.7 ± 44.5	351.4 ± 26.9
*QTc	470.4 ± 42.4	436.3 ± 18.7
*Abnormal QTc N(%)	14(46.7)	2 (6.7)
*Sinus tachycardia N(%)	27 (90.0)	7 (23.)
*T-wave axis deviation N (%)	8 (26.7)	1 (3.3)
Left atrial enlargement N(%)	5 (16.7)	0
Left axis deviation N (%)	2 (6.6)	0
Left ventricular hypertrophy N (%)	1 (3.3)	0
Premature ventricular contraction N(%)	1 (3.3)	0

*p <0.05.

values were 390-572 among the patients and 390 – 460 among the controls. Only 5 (16.7%) patients had QTc greater than 500. Compared to the controls, patients had significantly higher mean heart rate (121.1 ± 24.9 bpm versus 89.3 ± 12.9 bpm) ($t=6.2$, $p<0.05$), higher frequency of sinus tachycardia (OR = 29.57, 95% CI = 5.87 – 174.08, $p=0.00$) and significantly higher QTc (470.4 ± 42.4 versus 436.3 ± 17.7 ; $t=4.1$; $p<0.05$). The frequency of abnormal QTc was higher among the study group than controls (46.7% versus 6.6%, OR=9.19, 95% CI=1.60–68.48, $p=0.01$). T-axis deviation (-18 to 110) was present in 8 (26.6%) eclamptics and 1 (3.3%) woman with healthy pregnancy (OR = 10.55, 95% CI = 1.17–241.26, $p=0.03$).

An ECG tracing illustrating an eclamptic patient with abnormal QTc is shown in Figure 1.

Discussion

This study shows that eclampsia is associated with hypocalcaemia, abnormal QTc and T-axis deviation. Serum magnesium did not differ significantly among patients and controls. A relationship between low maternal serum calcium and pregnancy-related hypertension and eclampsia has been widely demonstrated^{3,4}. A high level of maternal parathyroid hormone has been suggested as the pathogenetic basis of hypocalcaemia in eclampsia³. Most of our patients, like in the previous reports from Nigeria¹⁶, were young primigravidae. The dominance of Hausa and Fulani tribes in the current report was due to the predominance of these tribes in Northwestern Nigeria where this study was carried out. Early marriage and child bearing are also frequent among these tribes.

The current report is to our knowledge the first on QT in eclampsia. Hypocalcaemia is a known cause of QT interval prolongation¹¹ and may play important roles in the genesis of abnormal QTc in eclampsia^{3,4}. Eclamptic patients as shown in this report were more likely to have hypocalcaemia than women with normal pregnancy. Hormonal changes may be a contributing factor to abnormal QT in eclampsia. Clinical and experimental studies have shown that female gender is associated with higher risk of QT interval prolongation than men. This has been attributed to a probable sex steroid-related specific regulation of ionic channels (calcium, potassium) such that estrogen facilitates QT interval response to agents or conditions that prolong QT interval¹⁷. Thus, the high estrogen level during pregnancy may therefore increase QT sensitivity to hypocalcaemia.

Alteration of trans-membrane cation contents is the major mechanism underlying the disorder of cardiac repolarisation. The net results of this disorder include reduced net ventricular repolarising current, amplified intrinsic spatial dispersion of repolarisation and development of QT interval prolongation and T-wave abnormality^{9,10}. The main significance of QT interval prolongation is however the development of arrhythmias. This largely accounts for the negative outcome of diseases associated with QT interval prolongation. Though an associa-

tion between abnormal QTc and mortality has been identified in patients with cardiac disease, its significance in patients without cardiac disease is unclear¹⁸.

Agents and conditions that prolong QT interval without increasing the transmural dispersion of repolarisation or early depolarisation-induced extrasystole do not cause arrhythmia. Sertindole, an antipsychotic agent, for example has low arrhythmia potential despite QT interval prolongation because it has no significant effect on ventricular dispersion of repolarisation¹⁹. Furthermore, QT interval prolongation less than 500 is unlikely to induce arrhythmias¹⁸. A few of our patients had T-axis deviation. This is an independent predictor of cardiovascular event, though more in men than women²⁰.

In conclusion, eclampsia is associated with prolonged ventricular repolarisation evidenced by abnormal QTc measured on the surface ECG. This may be related to maternal hypocalcaemia. Correction of electrolyte abnormalities and cautious use of agents that prolong QT interval are advocated in eclampsia.

References

1. Crowther CA. Eclampsia in Harare Maternity Hospital. *South Afr Med J* 1985; 68: 927 – 929.
2. Duley R. maternal mortality associated with hypertensive disorders in pregnancy in Africa, Asia, Latin America and the Caribbean. *Br J Obstet Gynaecol* 1992; 99: 547 – 553.
3. Villanueva LA, Figueroa A, Villanueva S. Blood concentrations of calcium and magnesium in women with severe pre-eclampsia. *ginecol Obstet Mex* 2002; 69: 277 – 281.
4. Males NO, Shuriden ZM. Does serum calcium in pre-eclampsia and uncomplicated pregnancy differ? *Saudi Med J* 2001; 22: 868 – 871.
5. Ehrenberg A. Non-medical prevention of pre-eclampsia. *Acta Obstetrica et gynaecologica Scandinavica* 1997; 164: 108 – 110.
6. The MAGPIE Trial Collaborative Group. Do women with pre-eclampsia and their babies benefit from magnesium sulphate? The Magpie Trial: a randomized placebo-controlled trial. *Lancet* 2002; 359: 1877 – 1890.
7. The Eclampsia Trial Collaborative Group. Which anti-convulsant for women? Evidence from the 4 Collaborative Eclampsia Trial. *Lancet* 1995, 345: 1455 – 1463.
8. Adewole IF, Oladokun A, Okewale AI, Omigbodun AO, Afolabi A, Ekele BA et al. Magnesium sulphate for treatment of Eclampsia: the Nigerian experience. *Afr J Med Med Sci* 2000; 29: 239 – 241.
9. Antzelevitch C, Shimizu W. Cellular mechanisms underlying the long QT syndrome. *Curr Opin Cardiol* 2002 Jan; 17: 433 – 51.
10. El-Shrif N. Mechanism of ventricular arrhythmias in the

- long QT syndrome on the hermeneutics. *J Cardiovasc Electrophysiol* 2001; 12: 973 – 976.
11. Soni CL, Kumhar MR, Gupta BK, Singh VB, Srimali L, Nayak KC et al. Prognostic implication of hypocalcemia and QTc interval in malaria. *Indian J Malariol* 2000; 37: 61 – 67.
 12. Piepho RW. Cardiovascular effects of antipsychotics used in bipolar illness. *J Clin Psychiatry* 2002; 63: Suppl 4: 20 – 23.
 13. Frothingham. Rates of torsades de pontes associated with ciprofloxacin, ofloxacin, levofloxacin, gatifloxacin, and moxifloxacin. *Pharmacotherapy* 2001; 21: 1468 – 72.
 14. World Health Organisation/.International Society of Hypertension. guideline for the management of hypertension. *J hypertens* 1999; 17: 151 – 183.
 15. American Heart Association committee. Report for standardisation of leads and specification for instruments in electrocardiography and vectorcardiography. *Circulation* 1975; 52: 11 – 31.
 16. Oladokun O, Okewale AI, Adewole IF, Babarinsa AI. Evaluation of cases of eclampsia in UCH, Ibadan over a 10 year period. *West Afr J Med* 2000; 19: 192 – 194.
 17. Schwartz JB. The electrocardiographic QT interval and its prolongation in response to medications: differences between men and women. *J Gend Specif Med* 2000; 3: 25 – 8.
 18. Bednar MM, Harrigan EP, Anziano RJ, Camm AJ, Ruskin JN. The QT interval. *Prog Cardiovasc Dis* 2001, 43: 1 – 45.
 19. Eckardt L, Breithardt G, Haverkamp W. Electrophysiologic characterisation of the antipsychotic drug sertindole in a rabbit heart model of torsade de pointes: low torsadogenic potential despite QT interval prolongation. *J Pharmacol Exp Ther* 2002; 300: 64 – 71.
 20. Rautaharju PM, Nelson JC, Kronmal RA, Zhang ZM, Robbins J, Gottdiener JS. Usefulness of T-axis deviation as an independent risk indicator for incident cardiac events in older men and women free from coronary heart disease (the Cardiovascular health Study). *Am J Cardiol* 2001; 88: 118 – 123.