

## SHORT COMMUNICATION

# Rapid Visual Assessment of Fish Communities on Selected Reefs in the Bazaruto Archipelago

Jade Q. Maggs<sup>1</sup>, Camilla Floros<sup>1</sup>, Marcos A.M. Pereira<sup>2</sup>  
and Michael H. Schleyer<sup>1</sup>

<sup>1</sup>Oceanographic Research Institute, P.O. Box 10712, Marine Parade, 4056 South Africa; <sup>2</sup>Associação para Investigação Costeira e Marinha (AICM), P.O. Box 2046, Maputo, Mozambique.

**Keywords:** Bazaruto Archipelago, Mozambique, ichthyofauna, fish surveys, underwater visual census, coral reefs.

**Abstract**— Rapid visual censuses were conducted of fish on eight coral reefs in the Bazaruto Archipelago, Mozambique, in 2007. SCUBA and snorkelling were used for the censuses in depths between 1-20 m, yielding an inventory of 249 fish species belonging to 50 families. This is intended to serve as a baseline for more detailed studies and monitoring programmes in the future. Although fewer species were recorded relative to other studies conducted in the Western Indian Ocean, the trophic structure on Bazaruto's reefs proved typical for the region, indicating a relative measure of reef health. However, other regional studies were not directly comparable, differing in habitat, duration of sampling effort and methodology. This highlighted the need for a long-term monitoring programme specifically adapted for the Bazaruto reef types to provide a basis for their sound management and conservation.

---

## INTRODUCTION

The Bazaruto Archipelago consists of five islands and is located approximately 20 km off the coast of

Mozambique in the Western Indian Ocean (WIO). The seas around the archipelago are rich in marine life and provide an important source of protein to the local community

(Everett *et al.*, 2008; Reina, 1998). The ecological importance of the archipelago first received attention in 1971 with three of the islands, Benguera, Magaruque and Bangue, receiving national park status (area of protection ~600 km<sup>2</sup>); Bazaruto and Santa Carolina were only designated as 'special surveillance zones' (Reina, 1998). After many years, the Bazaruto Archipelago National Park (BANP) was proclaimed in 2001, protecting all five islands. The BANP was extended in 2003 to include the Cabo de São Sebastião peninsula in the south and now covers 1430 km<sup>2</sup>. Mozambique's Ministry of Tourism is responsible for management of the BANP.

The coral and rocky reefs in the archipelago provide habitat for a wealth of biodiversity, making it a popular tourist destination (Schleyer & Celliers 2005). Visiting SCUBA divers and recreational anglers bring in valuable revenue to the area and, although fishing is allowed in the BANP, it is regulated by means of permits and no-take zones. All recreational fishing requires a permit and is mostly boat-based, emanating from a number of resorts scattered through the islands. A number of seasonal fishing competitions are hosted by the various resorts, bringing in foreign anglers.

The artisanal fishery is the main economic activity for more than 70% of the local population (Everett *et al.*, 2008). Artisanal fishermen harvest fish using dhows, pirogues, approved

beach seines, gamboa traps and spearguns. Mozambican citizens living outside the BANP are only allowed to use handlines inside the park and are allowed to use beach-seines in a small area south of Magaruque. Gill-netting is prohibited and no industrial or semi-industrial fishing operations are allowed in the BANP. The no-take zones are Two-mile Reef, Lighthouse Reef, Santa Carolina and small rocky outcrops on the inside and outside of Bazaruto Island (Fig. 1). However, only Two-mile Reef, Lighthouse Reef and Santa Carolina enjoy strong compliance. There are no seasonal restrictions on fishing in the BANP and recreational SCUBA diving and snorkelling is allowed in the no-take zones. A new management plan is being developed for the Park but, at the time of writing, had not yet been implemented.

The livelihoods of the local communities depend, to a great extent, on marine and coastal resources (Everett *et al.*, 2008). An increasing population characterises many coastal communities in the WIO, placing pressure on such resources (Lindén *et al.*, 2002). Effective management is thus required, the success of which depends on monitoring programmes (Obura *et al.*, 2002). These have been seen as a priority in the WIO region since the 1998 mass coral bleaching event (Lindén *et al.*, 2002). Monitoring of a resource depends on a thorough knowledge of the biodiversity of an area. Species inventories comprise

a fundamental baseline in this regard, providing a foundation for an understanding of ecological processes and the effects of biodiversity loss on ecosystem function (Bellwood & Hughes 2001; Gillibrand *et al.*, 2007). Baseline fish community data are sparse for the WIO region but those published include fish inventories for the Glorieuses Islands (Durville *et al.*, 2003), Mayotte (Chabanet, 2002), Andavadoaka (Gillibrand *et al.*, 2007), Juan De Nova (Chabanet & Durville 2005), Tuléar (Harmelin-Vivien, 1979) Geyser and Zéléé (Chabanet *et al.*, 2002), Réunion (Chabanet, 1994), Sodwana Bay (Chater *et al.*, 1993; 1995) and Bassas da India (van der Elst & Chater 2001).

Pereira (2000) prepared a general checklist of reef-associated fishes for Mozambique, Benayahu & Schleyer (1996) and Schleyer & Celliers (2005) compiled coral inventories for the Bazaruto reefs, Motta *et al.*, (2002) and Rodrigues *et al.*, (2000) quantified fish communities at two sites off Bazaruto and van der Elst & Afonso (2008) compiled a fish inventory based on work undertaken at Bazaruto in the late 1980s. However, there appears to have been no further ichthyofaunal research in the Bazaruto Archipelago. This paper, therefore, presents a recent and more comprehensive inventory of the fish community as a precursor to further ecological studies on the Bazaruto reefs and highlights the need for long-term monitoring.

## MATERIALS AND METHODS

### Study Area

The Bazaruto Archipelago is a chain of four islands, Bazaruto, Benguera, Magaruque and Bangué, with a fifth island, Santa Carolina, lying on the inside of the island chain (Fig. 1). The reefs around the archipelago are diverse and include rocky patch reefs, rocky massifs, fringing and barrier coral reefs and deeply submerged coral reefs (Schleyer & Maggs 2008). The fish community assessment was undertaken on all these reef types in depths from 1-20 m. A range of reef habitats within these reef types was sampled with varying coral cover and topographic complexity.

Twelve-mile Reef is a submerged sandstone coral reef roughly twelve nautical miles (18 km) north of Bazaruto's northern point in the open sea. It is open to fishing and SCUBA diving but is relatively inaccessible for artisanal fisherman using traditional pirogues and dhows. However, recreational fishers and SCUBA divers, using large boats with outboard engines, are able to access this reef. Relative to the other reefs sampled in this study, Twelve-mile Reef lies furthest from the waters enclosed between the islands and the mainland. The reef was sampled at depths between 15-20 m.

Tubarão, Garoupa and Kingfish Reefs are sedimented rocky patch reefs which are open to fishing and



Fig. 1. Map of the Bazaruto Archipelago, Mozambique. Study sites are indicated by (◆).

SCUBA diving. Garoupa lies nine km north of Bazaruto and was sampled at 16-20 m. Tubarão lies 19 km north-east of Inhassoro and was sampled at 13-18 m. Kingfish Reef is 13.5 km east of Inhassoro and was sampled at 6-11 m. High turbidity is common on these reefs.

Lighthouse Reef is a fringing coral reef located on the north-eastern tip of Bazaruto. It is a no-take zone and is closed to fishing (including artisanal fishing) but open to SCUBA diving and snorkelling. Only the inner lagoon was sampled and the depth ranged between 1-3 m.

Two-mile Reef is a barrier coral reef which lies four kilometres out to sea between Bazaruto and Benguera Islands. This reef is also a no-take zone being closed to fishing (including artisanal fishing). Although fishing is prohibited, Two-mile Reef is subjected to diver pressure and anchor damage from visiting small craft, recreational dive operators and tourists. All reef habitats at Two-mile Reef were sampled at depths between 1-18 m.

Amphitheatre and Camel's Hump are submerged rocky massifs located two kilometres seaward of Cabo de São Sebastião. They are open to fishing and SCUBA diving but turbidity is high on these reefs. Sampling at Camel's Hump ranged between 13-16 m in depth and at Amphitheatre, between 14-19 m.

## Data Collection

Two fish surveys were conducted, one in February 2007 and one in November 2007. Surveys were undertaken in the late morning on a low to outgoing spring tide. Fish communities were sampled on the reefs using an underwater visual census technique adapted from Samoilys (1997) in which three divers recorded the presence of fish species on slates. A combination of diving methods was used with SCUBA being used for deeper locations and snorkelling for shallow inner lagoons. Divers conducted a 45 minute timed swim following a random path. Although the underwater visual census method is known to underestimate small and cryptic species, it was employed because it provides a means of sampling the fish community with little disturbance (Fowler, 1987; Harmelin-Vivien *et al.*, 1985). Identification of species was confirmed after sampling using appropriate reference books (King, 1996; King & Fraser 2001; Lieske & Myers 1999; Smith & Heemstra 1986).

## Trophic Categorisation

Fish species were assigned to one of ten trophic categories based on classifications by Harmelin-Vivien (1979); Hiatt & Strasberg (1960); Hobson (1974) and Myers (1999) as cited by Chabanet & Durville

(2005); Durville *et al.*, (2003) and Gillibrand *et al.*, (2007). These studies used eight categories: herbivores, omnivores, browsers of sessile invertebrates, diurnal carnivores, nocturnal carnivores, piscivores, diurnal planktivores and nocturnal planktivores. In the present study, some of these categories were consolidated, viz. general carnivores and general planktivores, as the diel preference of some species was unknown (Froese & Pauly 2009; Heemstra & Heemstra 2004; King, 1996; King & Fraser 2001 and Smith & Heemstra 1986).

## RESULTS

### Species Richness

A total of 249 species belonging to 50 families were recorded (Table 1), of which six were cartilaginous fishes in four families and the remaining 243 species were bony fish in 46 families. The top five families according to species count were the Labridae (37 species), Acanthuridae (22 species), Chaetodontidae (22 species), Pomacentridae (17 species) and Serranidae (13 species). These five families contributed 45% to the species diversity (Table 2). Overall, 19 families were represented by only one species.

Fish families and species were not evenly distributed among all the reefs (Table 2). The top three reefs according to species richness were Two-mile Reef with 197 species, Lighthouse Reef with 103 species, and Garoupa

**Table 2. Summary of the number of species and families recorded on the Bazaruto reefs. Shaded reefs are no-take zones where fishing (including artisanal fishing) is prohibited, but SCUBA diving and snorkelling are allowed. Reefs are ordered according to increasing latitude.**

| Reef             | No. of Species | No. of Families |
|------------------|----------------|-----------------|
| Twelve-mile Reef | 59             | 21              |
| Tubarão          | 44             | 20              |
| Garoupa          | 101            | 31              |
| Kingfish         | 84             | 31              |
| Lighthouse Reef  | 103            | 25              |
| Two-mile Reef    | 197            | 43              |
| Camel's Hump     | 45             | 21              |
| Amphitheatre     | 50             | 22              |

with 101 species. Two-mile Reef also had the highest number of fish families (43) but Lighthouse Reef, despite having the second highest number of species, had relatively few families (25). Conversely, Garoupa had a relatively high number of fish families (31). All the other reefs had relatively few fish families and species; only Kingfish was better represented by 84 species in 31 families.

### Trophic Structure

When all the carnivorous categories were grouped (i.e. all groups except herbivores and omnivores), they constituted 76% of the species composition (Fig. 2). Herbivores, (mostly acanthurids) and omnivores (mostly pomacentrids) each accounted for 12% of the species composition. The largest group, diurnal carnivores (27%), was dominated by labrids




Figure 2. Overall trophic structure of Bazaruto reef fish communities.

and the nocturnal carnivores (17%) were dominated by larger lutjanids, lethrinids and serranids. Chaetodons accounted for the majority of browsers of sessile invertebrates. None of the other categories was dominated by any specific family. The piscivores, contributing 5% to the species composition, comprised mostly larger predators such as *Carcharhinus amblyrhynchos*, *Aprion virescens* and *Scomberomorus commerson*.

## DISCUSSION

### Species Richness

This study yielded 249 fish species in 50 families, a lower tally than other studies in the region. Durville *et al.*, (2003) recorded 332 fish species in 57 families at the Glorieuses Islands, while Chabanet & Durville (2005) listed 299 species in 55 families for Juan De Nova. Further south, Gillibrand *et al.*, (2007) counted 334 species of fish in

**Table 1. Species list of the Bazaruto Archipelago on a per reef basis (depth 1-20 m). Presence is indicated by (●). H, herbivores; O, omnivores; BSI, browsers of sessile invertebrates; DC, Diurnal carnivores; NC, Nocturnal carnivores; PI, Piscivores; DP, Diurnal planktivores; NP, Nocturnal planktivores; C, General carnivores; PL, General planktivores. Shaded reefs are no-take zones where fishing (including artisanal fishing) is prohibited, but SCUBA diving and snorkelling are allowed.**

| FAMILY species   | Trophic Category | Twelve-mile Reef | Tubarão | Garoupa | Kingfish | Lighthouse Reef | Two-mile Reef | Camel's Hump | Amphitheatre |
|--|------------------|------------------|---------|---------|----------|-----------------|---------------|--------------|--------------|
| <b>Acanthuridae</b>  |                  |                  |         |         |          |                 |               |              |              |
| <i>Acanthurus dussumieri</i> Valenciennes, 1835                              | H                |                  | ●       | ●       |          | ●               | ●             | ●            | ●            |
| <i>Acanthurus leucocheilus</i> Herre, 1927                                   | H                |                  |         |         |          |                 | ●             |              |              |
| <i>Acanthurus leucosternon</i> Bennet, 1833                                  | H                | ●                |         | ●       | ●        | ●               | ●             | ●            | ●            |
| <i>Acanthurus lineatus</i> (Linnaeus, 1758)                                  | H                |                  |         |         |          | ●               | ●             |              |              |
| <i>Acanthurus mata</i> Russel in Cuvier, 1829                                | DP               |                  | ●       |         |          |                 | ●             |              |              |
| <i>Acanthurus nigrofuscus</i> (Forsskål, 1775)                               | H                | ●                | ●       | ●       |          | ●               | ●             | ●            | ●            |
| <i>Acanthurus tennentii</i> Günther, 1861                                    | H                | ●                |         | ●       | ●        | ●               | ●             | ●            |              |
| <i>Acanthurus thompsoni</i> Fowler, 1923                                     | H                |                  |         |         | ●        |                 |               |              | ●            |
| <i>Acanthurus triostegus triostegus</i> (Linnaeus, 1758)                     | H                |                  |         |         |          | ●               | ●             |              |              |
| <i>Ctenochaetus binotatus</i> Randall, 1955                                  | H                |                  |         |         |          | ●               | ●             |              |              |
| <i>Ctenochaetus strigosus</i> (Bennet, 1828)                                 | H                |                  |         | ●       |          | ●               | ●             |              |              |
| <i>Naso annulatus</i> (Quoy & Gaimard, 1825)                                 | H                |                  |         |         |          | ●               | ●             |              |              |
| <i>Naso brachycentron</i><br>(Valenciennes in Cuvier and Valenciennes, 1835) | H                |                  |         |         |          |                 | ●             |              |              |
| <i>Naso brevirostris</i> (Cuvier, 1829)                                      | H                |                  | ●       |         |          | ●               | ●             | ●            |              |
| <i>Naso hexacanthus</i> (Bleeker, 1855)                                      | H                | ●                |         |         | ●        |                 | ●             |              |              |
| <i>Naso lituratus</i> (Forster in Bloch & Schneider, 1801)                   | H                | ●                |         |         | ●        | ●               | ●             | ●            | ●            |
| <i>Naso unicornis</i> (Forsskål, 1775)                                       | H                | ●                |         |         |          | ●               | ●             |              |              |
| <i>Naso vlamingii</i> (Valenciennes, 1835)                                   | DP               | ●                |         |         |          | ●               | ●             |              |              |
| <i>Paracanthurus hepatus</i> (Linnaeus, 1766)                                | DP               | ●                |         |         |          | ●               | ●             |              |              |
| <i>Zebrasoma gemmatum</i> (Valenciennes, 1835)                               | H                |                  |         |         |          | ●               | ●             |              |              |
| <i>Zebrasoma scopas</i> (Cuvier, 1829)                                       | H                | ●                |         |         |          | ●               | ●             |              | ●            |
| <i>Zebrasoma desjardini</i> (Bennet, 1836)                                   | H                |                  |         |         |          | ●               | ●             |              |              |
| <b>APOGONIDAE</b>  |                  |                  |         |         |          |                 |               |              |              |
| <i>Apogon aureus</i> (Lacepède, 1802)  | DC               |                  |         |         | ●        |                 |               |              |              |
| <b>AULOSTOMIDAE</b>  |                  |                  |         |         |          |                 |               |              |              |
| <i>Aulostomus chinensis</i> (Linnaeus, 1766)                                 | PI               |                  |         |         | ●        |                 | ●             |              |              |
| <b>BALISTIDAE</b>  |                  |                  |         |         |          |                 |               |              |              |
| <i>Balistapus undulatus</i> (Mungo Park, 1797)                               | DC               |                  | ●       | ●       | ●        | ●               | ●             |              |              |
| <i>Balistoides conspicillum</i> (Bloch & Schneider, 1801)                    | DC               | ●                |         |         | ●        |                 | ●             |              | ●            |
| <i>Balistoides viridescens</i> (Bloch & Schneider, 1801)                     | DC               |                  |         |         |          | ●               | ●             |              |              |
| <i>Odonus niger</i> (Rüppel, 1836)   | DC               | ●                | ●       | ●       | ●        |                 |               | ●            | ●            |
| <i>Pseudobalistes fuscus</i> (Bloch & Schneider, 1801)                       | C                |                  |         |         | ●        |                 |               |              |              |
| <i>Rhinecanthus rectangulus</i> (Bloch & Schneider, 1801)                    | O                |                  |         |         | ●        | ●               |               |              |              |
| <i>Sufflamen chrysopterus</i> (Bloch & Schneider, 1801)                      | DC               |                  | ●       |         |          | ●               | ●             |              |              |
| <i>Sufflamen fraenatum</i> (Latreille, 1804)                                 | DC               | ●                | ●       | ●       | ●        | ●               | ●             |              |              |
| <b>BLENNIIDAE</b>  |                  |                  |         |         |          |                 |               |              |              |
| <i>Ecsenius midas</i> Stark, 1969  | H                |                  |         |         |          |                 | ●             | ●            |              |
| <i>Plagiotremus rhinorhynchus</i> (Bleeker, 1852)                            | NP               |                  |         | ●       |          | ●               |               |              |              |
| <i>Plagiotremus tapeinosoma</i> (Bleeker, 1857)                              | O                |                  |         |         | ●        |                 | ●             |              |              |



| <b>FAMILY species</b>  | Trophic Category | Twelve-mile Reef | Tubarão | Garoupa | Kingfish | Lighthouse Reef | Two-mile Reef | Camel's Hump | Amphitheatre |
|--|------------------|------------------|---------|---------|----------|-----------------|---------------|--------------|--------------|
| <b>CAESIONIDAE</b>   |                  |                  |         |         |          |                 |               |              |              |
| <i>Caesio caerulaurea</i> (Lacepède, 1801)                                     | DP               |                  |         |         |          | •               | •             |              |              |
| <i>Caesio lunaris</i> Cuvier, 1830   | DP               |                  |         |         |          | •               | •             |              |              |
| <i>Caesio sp.</i> Lacepède, 1801   | DP               |                  |         | •       |          |                 |               |              |              |
| <i>Caesio xanthonota</i> Bleeker, 1853   | DP               |                  | •       | •       | •        |                 | •             | •            |              |
| <i>Pterocaesio sp.</i> Bleeker, 1876   | DP               |                  |         | •       |          |                 |               |              |              |
| <i>Pterocaesio tile</i> (Cuvier, 1830)   | DP               |                  |         | •       |          |                 | •             |              |              |
| <b>CARANGIDAE</b>  |                  |                  |         |         |          |                 |               |              |              |
| <i>Carangoides fulvoguttatus</i> (Forsskål, 1775)                              | DC               |                  |         | •       |          | •               | •             | •            |              |
| <i>Caranx ignobilis</i> (Forsskål, 1775)                                       | DC               |                  |         | •       |          |                 |               |              |              |
| <i>Caranx melampygus</i> Cuvier & Valenciennes, 1833                           | DC               |                  |         | •       |          | •               | •             | •            |              |
| <i>Caranx papuensis</i> Alleyne & MacLeay, 1877                                | C                |                  |         |         | •        |                 |               |              |              |
| <i>Elagatis bipinnulata</i> (Quoy & Gaimard, 1825)                             | DC               |                  |         |         |          |                 | •             | •            |              |
| <i>Gnathanodon speciosus</i> (Forsskål, 1775)                                  | DC               |                  |         |         |          |                 | •             |              |              |
| <i>Scomberoides lysan</i> (Forsskål, 1775)                                     | PI               |                  |         |         |          |                 | •             |              |              |
| <b>CARCHARHINIDAE</b>  |                  |                  |         |         |          |                 |               |              |              |
| <i>Carcharhinus amblyrhynchus</i> (Bleeker, 1856)                              | PI               | •                |         |         |          |                 | •             |              |              |
| <i>Triaenodon obesus</i> (Rüppel, 1837)  | DC               |                  |         |         |          |                 |               |              | •            |
| <b>CHAETODONTIDAE</b>  |                  |                  |         |         |          |                 |               |              |              |
| <i>Chaetodon auriga</i> Forsskål, 1775   | BSI              | •                | •       | •       | •        | •               | •             | •            | •            |
| <i>Chaetodon blackburnii</i> Desjardins, 1836                                  | BSI              |                  |         | •       |          |                 |               |              |              |
| <i>Chaetodon dolosus</i> Ahl, 1923   | O                |                  | •       | •       |          |                 |               |              |              |
| <i>Chaetodon falcula</i> Bloch, 1793   | BSI              |                  |         |         |          |                 | •             |              |              |
| <i>Chaetodon guttatissimus</i> Bennet, 1832                                    | BSI              | •                | •       | •       | •        | •               | •             | •            | •            |
| <i>Chaetodon interruptus</i> Ahl, 1923   | BSI              | •                |         |         |          | •               | •             | •            | •            |
| <i>Chaetodon kleinii</i> Bloch, 1790   | BSI              | •                | •       | •       | •        | •               | •             | •            | •            |
| <i>Chaetodon lineolatus</i><br>(Quoy & Gaimard, 1831 in Cuvier & Valenciennes) | BSI              |                  |         |         |          | •               | •             |              |              |
| <i>Chaetodon lunula</i> (Lacepède, 1802)                                       | BSI              | •                |         | •       | •        | •               | •             | •            | •            |
| <i>Chaetodon madagaskariensis</i> Ahl, 1923                                    | BSI              | •                |         | •       | •        | •               | •             | •            | •            |
| <i>Chaetodon melannotus</i> (Bloch & Schneider, 1801)                          | BSI              | •                |         |         |          | •               | •             |              |              |
| <i>Chaetodon meyeri</i> (Bloch & Schneider, 1801)                              | BSI              | •                |         |         |          | •               | •             |              | •            |
| <i>Chaetodon trifasciatus</i> Quoy & Gaimard, 1825                             | BSI              |                  |         |         |          | •               | •             |              |              |
| <i>Chaetodon trifasciatus</i> (Mungo Park, 1797)                               | BSI              | •                |         | •       |          | •               | •             | •            | •            |
| <i>Chaetodon vagabundus</i> Linnaeus, 1758                                     | BSI              |                  | •       | •       | •        | •               | •             | •            | •            |
| <i>Chaetodon xanthocephalus</i> Bennet, 1832                                   | BSI              |                  |         |         |          | •               | •             |              | •            |
| <i>Chaetodon zanzibarensis</i><br>Playfair, in Playfair & Günther, 1867        | BSI              |                  |         |         |          |                 | •             |              |              |
| <i>Forcipiger flavissimus</i> Jordan & McGregor, 1898                          | BSI              |                  |         | •       |          |                 | •             | •            | •            |
| <i>Hemitaurichthys zoster</i> (Bennet, 1831)                                   | DP               |                  |         |         |          |                 | •             |              | •            |
| <i>Heniochus acuminatus</i> (Linnaeus, 1758)                                   | BSI              | •                | •       | •       | •        | •               | •             | •            | •            |
| <i>Heniochus diphreutes</i> Jordan, 1903                                       | PL               |                  |         |         |          |                 | •             |              |              |
| <i>Heniochus monoceros</i><br>Cuvier in Cuvier and Valenciennes, 1831          | BSI              |                  | •       |         |          |                 | •             |              |              |
| <b>CIRRHITIDAE</b>   |                  |                  |         |         |          |                 |               |              |              |
| <i>Cirrhitichthys oxycephalus</i> (Bleeker, 1855)                              | DC               | •                | •       | •       |          |                 | •             |              |              |
| <i>Paracirrhites arcatus</i><br>Cuvier in Cuvier and Valenciennes, 1829        | DC               | •                |         | •       | •        |                 | •             |              |              |
| <i>Paracirrhites forsteri</i> (Bloch & Schneider, 1801)                        | DC               |                  |         | •       |          | •               | •             |              | •            |

| <b>FAMILY species</b>                               | Trophic Category | Twelve-mile Reef | Tubarão | Garoupa | Kingfish | Lighthouse Reef | Two-mile Reef | Camel's Hump | Amphitheatre |
|---|------------------|------------------|---------|---------|----------|-----------------|---------------|--------------|--------------|
| <b>DASYATIDAE</b>                                   |                  |                  |         |         |          |                 |               |              |              |
| <i>Himantura gerrardi</i> (Gray, 1851)              | C                |                  | •       |         |          |                 |               |              |              |
| <i>Taeniura lymma</i> (Forsskål, 1775)              | NC               |                  |         |         |          |                 | •             |              |              |
| <b>DIODONTIDAE</b>                                  |                  |                  |         |         |          |                 |               |              |              |
| <i>Diodon liturosus</i> Shaw, 1804                  | NC               |                  |         |         |          |                 | •             |              |              |
| <b>ECHENEIDAE</b>                                   |                  |                  |         |         |          |                 |               |              |              |
| <i>Echeneis naucrates</i> Linnaeus, 1758            | NC               | •                |         | •       |          |                 | •             |              |              |
| <b>EPHIPPIDAE</b>                                   |                  |                  |         |         |          |                 |               |              |              |
| <i>Platax orbicularis</i> (Forsskål, 1775)          | O                | •                |         | •       |          |                 | •             |              | •            |
| <i>Platax teira</i> (Forsskål, 1775)                | O                |                  |         |         |          |                 | •             |              |              |
| <i>Tripteron orbis</i> Playfair, 1867               | C                |                  |         |         | •        |                 |               |              |              |
| <b>FISTULARIIDAE</b>                                |                  |                  |         |         |          |                 |               |              |              |
| <i>Fistularia commersonii</i> Rüppel, 1838          | DC               |                  |         |         |          | •               | •             |              | •            |
| <b>GOBIIDAE</b>                                     |                  |                  |         |         |          |                 |               |              |              |
| <i>Valenciennesa strigata</i> (Broussonet, 1782)    | DC               |                  |         | •       |          |                 | •             |              | •            |
| <b>HAEMULIDAE</b>                                   |                  |                  |         |         |          |                 |               |              |              |
| <i>Plectorhinchus chubbi</i> (Thunberg, 1792)       | DC               |                  |         | •       |          | •               | •             |              |              |
| <i>Plectorhinchus plagiodesmus</i> Fowler, 1935     | NC               |                  | •       |         | •        |                 |               |              |              |
| <i>Plectorhinchus flavomaculatus</i> (Cuvier, 1830) | NC               | •                | •       | •       | •        | •               | •             |              | •            |
| <i>Plectorhinchus gaterinus</i> (Forsskål, 1775)    | NC               |                  |         | •       | •        | •               | •             |              |              |
| <i>Plectorhinchus gibbosus</i> (Lacepède, 1802)     | C                |                  |         | •       |          |                 |               |              |              |
| <i>Plectorhinchus playfairi</i> (Pellegrin, 1914)   | DC               |                  | •       | •       | •        | •               | •             | •            | •            |
| <i>Plectorhinchus schotaf</i> (Forsskål, 1775)      | C                |                  |         |         | •        |                 |               |              |              |
| <b>HEMIRAMPHIDAE</b>                                |                  |                  |         |         |          |                 |               |              |              |
| <i>Hyporhamphus affinis</i> (Günther, 1866)         | O                |                  |         |         |          | •               | •             |              |              |
| <b>HOLOCENTRIDAE</b>                                |                  |                  |         |         |          |                 |               |              |              |
| <i>Myripristis botche</i> Cuvier, 1829              | NC               |                  |         |         | •        |                 |               |              |              |
| <i>Myripristis murdjan</i> Forsskål, 1775           | NP               |                  |         |         | •        | •               | •             |              |              |
| <i>Neoniphon argenteus</i> (Valenciennes, 1831)     | C                |                  |         |         |          | •               | •             |              |              |
| <i>Neoniphon sammara</i> Forsskål, 1775             | NC               |                  |         |         |          | •               | •             |              |              |
| <i>Sargocentron caudimaculatum</i> Rüppel, 1838     | NC               |                  |         |         | •        |                 | •             |              |              |
| <i>Sargocentron diadema</i> Lacepède, 1802          | NC               |                  |         | •       |          | •               | •             |              |              |
| <i>Sargocentron spiniferum</i> Forsskål, 1775       | NC               |                  |         |         |          |                 |               |              | •            |
| <b>KYPHOSIDAE</b>                                   |                  |                  |         |         |          |                 |               |              |              |
| <i>Kyphosus cinerascens</i> Forsskål, 1775          | H                |                  |         |         |          |                 | •             |              |              |
| <i>Kyphosus</i> sp. Lacepède, 1801                  | H                |                  |         |         |          | •               |               |              |              |
| <i>Kyphosus vaigiensis</i> (Quoy & Gaimard, 1825)   | O                |                  |         |         | •        |                 | •             |              |              |
| <b>LABRIDAE</b>                                     |                  |                  |         |         |          |                 |               |              |              |
| <i>Anampses caeruleopunctatus</i> Rüppel, 1829      | DC               |                  |         | •       | •        | •               | •             |              |              |
| <i>Anampses lineatus</i> Randall, 1972              | DC               |                  |         | •       |          |                 |               |              |              |
| <i>Anampses meleagrides</i> Valenciennes, 1840      | DC               |                  |         | •       | •        |                 | •             |              |              |
| <i>Anampses twistii</i> Bleeker, 1856               | DC               |                  |         |         | •        |                 | •             |              |              |
| <i>Bodianus anthioides</i> Bennet, 1832             | DC               |                  |         |         |          |                 | •             |              |              |
| <i>Bodianus axillaris</i> Bennet, 1832              | DC               |                  | •       |         | •        |                 | •             |              |              |
| <i>Bodianus bilunulatus</i> (Lacepède, 1801)        | DC               |                  |         |         | •        |                 | •             |              |              |

| <b>FAMILY species</b>  | <b>Trophic Category</b> | <b>Twelve-mile Reef</b> | <b>Tubarão</b> | <b>Garoupa</b> | <b>Kingfish</b> | <b>Lighthouse Reef</b> | <b>Two-mile Reef</b> | <b>Camel's Hump</b> | <b>Amphitheatre</b> |
|--|-------------------------|-------------------------|----------------|----------------|-----------------|------------------------|----------------------|---------------------|---------------------|
| <i>Bodianus diana</i> Lacepède, 1801                                   | DC                      | •                       | •              | •              | •               |                        | •                    |                     |                     |
| <i>Cheilinus fasciatus</i> (Bloch, 1791)                               | DC                      |                         |                |                |                 | •                      | •                    |                     |                     |
| <i>Cheilinus trilobatus</i> Lacepède, 1801                             | DC                      |                         |                |                |                 | •                      | •                    |                     |                     |
| <i>Cheilinus undulatus</i> Rüppel, 1835                                | DC                      | •                       |                |                |                 |                        | •                    |                     |                     |
| <i>Cheilio inermis</i> Forsskål, 1775                                  | DC                      |                         |                |                |                 |                        | •                    |                     |                     |
| <i>Cirrhilabrus exquisitus</i> Smith, 1957                             | DC                      |                         |                |                | •               |                        | •                    |                     |                     |
| <i>Coris aygula</i> Lacepède, 1801                                     | DC                      |                         |                |                | •               |                        | •                    |                     |                     |
| <i>Coris caudimacula</i> Quoy & Gaimard, 1834                          | DC                      |                         |                |                |                 |                        | •                    |                     |                     |
| <i>Coris cuvieri</i> (Bennett, 1831)                                   | DC                      |                         |                | •              |                 |                        | •                    |                     |                     |
| <i>Coris frerei</i> (Bennett, 1830)                                    | DC                      |                         |                | •              | •               | •                      | •                    |                     | •                   |
| <i>Gomphosus caeruleus</i> Lacepède, 1801                              | DC                      | •                       |                |                | •               | •                      | •                    |                     |                     |
| <i>Halichoeres cosmetus</i> Randall, & Smith, 1982                     | DC                      |                         |                | •              |                 |                        | •                    |                     |                     |
| <i>Halichoeres hortulanus</i> Lacepède, 1801                           | DC                      | •                       |                | •              |                 | •                      | •                    |                     |                     |
| <i>Halichoeres iridis</i> Randall, & Smith, 1982                       | DC                      |                         |                | •              |                 | •                      | •                    |                     |                     |
| <i>Halichoeres scapularis</i> Bennet, 1832                             | DC                      |                         |                |                |                 |                        | •                    |                     |                     |
| <i>Hemigymnus fasciatus</i> Bloch, 1792                                | DC                      |                         |                |                | •               | •                      | •                    |                     |                     |
| <i>Hologymnosus annulatus</i> Lacepède, 1801                           | DC                      |                         |                |                |                 |                        | •                    | •                   |                     |
| <i>Hologymnosus doliatus</i> Lacepède, 1801                            | DC                      |                         |                |                |                 |                        | •                    |                     |                     |
| <i>Labroides bicolor</i> Fowler & Bean, 1928                           | DC                      |                         |                | •              |                 |                        | •                    | •                   |                     |
| <i>Labroides dimidiatus</i> Valenciennes, 1839                         | DC                      | •                       | •              | •              | •               | •                      | •                    |                     | •                   |
| <i>Macropharyngodon bipartitus</i> Smith 1957                          | DC                      |                         |                | •              | •               |                        | •                    |                     |                     |
| <i>Macropharyngodon cyanoguttatus</i> Randall, 1978                    | DC                      |                         |                |                |                 |                        | •                    |                     |                     |
| <i>Novaculichthys taeniourus</i> (Lacepède, 1801)                      | DC                      |                         |                | •              |                 |                        | •                    |                     |                     |
| <i>Pseudocheilinus hexataenia</i> (Bleeker, 1857)                      | DC                      |                         |                |                |                 | •                      | •                    |                     |                     |
| <i>Pseudodax moluccanus</i> (Valenciennes, 1840)                       | O                       |                         |                | •              | •               |                        | •                    |                     |                     |
| <i>Stethojulis interrupta</i> (Bleeker, 1851)                          | DC                      |                         |                |                |                 |                        | •                    |                     |                     |
| <i>Thalassoma amblycephalum</i> Bleeker, 1856                          | DC                      |                         |                | •              | •               | •                      | •                    | •                   |                     |
| <i>Thalassoma hardwicke</i> Bennet, 1830                               | DC                      |                         |                |                |                 |                        | •                    |                     |                     |
| <i>Thalassoma hebraicum</i> Lacepède, 1801                             | DC                      | •                       | •              | •              | •               | •                      | •                    |                     |                     |
| <i>Thalassoma lunare</i> Linnaeus, 1758                                | DC                      | •                       |                | •              | •               | •                      | •                    | •                   |                     |
| <b>LETHRINIDAE</b>   |                         |                         |                |                |                 |                        |                      |                     |                     |
| <i>Gnathodentex aureolineatus</i> (Lacepède, 1802)                     | NC                      |                         |                |                | •               | •                      | •                    |                     |                     |
| <i>Lethrinus crocineus</i> Smith, 1959                                 | NC                      |                         |                |                |                 |                        | •                    |                     |                     |
| <i>Lethrinus harak</i> (Forsskål, 1775)                                | NC                      |                         |                |                |                 | •                      | •                    |                     |                     |
| <i>Lethrinus nebulosus</i> (Forsskål, 1775)                            | NC                      | •                       |                |                |                 |                        | •                    |                     |                     |
| <i>Lethrinus rubrioperculatus</i> Sato, 1978                           | NC                      | •                       |                |                |                 |                        | •                    |                     |                     |
| <i>Lethrinus mahsena</i> (Forsskål, 1775)                              | NC                      |                         |                |                |                 |                        | •                    |                     |                     |
| <i>Monotaxis grandoculis</i> (Forsskål, 1775)                          | NC                      |                         |                |                | •               | •                      | •                    |                     |                     |
| <b>LUTJANIDAE</b>  |                         |                         |                |                |                 |                        |                      |                     |                     |
| <i>Aphareus furca</i> (Lacepède, 1801)                                 | PI                      |                         |                |                |                 |                        | •                    |                     |                     |
| <i>Aprion virescens</i> Valenciennes, 1830                             | PI                      | •                       | •              | •              |                 |                        | •                    | •                   | •                   |
| <i>Lutjanus argentimaculatus</i> (Forsskål, 1775)                      | NC                      |                         |                | •              |                 |                        | •                    |                     |                     |
| <i>Lutjanus bohar</i> (Forsskål, 1775)                                 | NC                      | •                       | •              | •              |                 | •                      | •                    |                     | •                   |
| <i>Lutjanus fulviflamma</i> (Forsskål, 1775)                           | NC                      | •                       |                |                | •               | •                      | •                    |                     |                     |
| <i>Lutjanus gibbus</i> (Forsskål, 1775)                                | NC                      | •                       |                | •              |                 | •                      | •                    |                     |                     |
| <i>Lutjanus kasmira</i> (Forsskål, 1775)                               | NC                      |                         | •              | •              | •               | •                      | •                    | •                   | •                   |
| <i>Lutjanus lutjanus</i> Bloch, 1790                                   | NC                      |                         | •              | •              |                 |                        | •                    |                     |                     |
| <i>Lutjanus notatus</i> (Cuvier, 1828)                                 | NC                      |                         | •              | •              | •               |                        |                      |                     |                     |
| <i>Lutjanus rivulatus</i><br>(Cuvier in Cuvier and Valenciennes, 1828) | NC                      |                         |                | •              |                 |                        |                      |                     |                     |

| <b>FAMILY species</b>  | Trophic Category | Twelve-mile Reef Tubarão | Garoupa | Kingfish | Lighthouse Reef | Two-mile Reef | Camel's Hump Amphitheatre |
|--|------------------|--------------------------|---------|----------|-----------------|---------------|---------------------------|
| <i>Lutjanus sebae</i> (Cuvier, 1816)   | NC               |                          | •       |          |                 |               |                           |
| <i>Macolor niger</i> (Forsskål, 1775)  | NC               |                          |         |          |                 | •             | •                         |
| <b>MALACANTHIDAE</b>   |                  |                          |         |          |                 |               |                           |
| <i>Malacanthus brevirostris</i> Guichenot, 1848                                      | DC               |                          | •       |          |                 | •             |                           |
| <b>MICRODESMIDAE</b>   |                  |                          |         |          |                 |               |                           |
| <i>Ptereleotris evides</i> (Jordan & Hubbs, 1925)                                    | DP               |                          |         |          |                 | •             |                           |
| <i>Ptereleotris heteroptera</i> (Bleeker, 1855)                                      | DP               |                          |         |          |                 | •             |                           |
| <b>MOBULIDAE</b>   |                  |                          |         |          |                 |               |                           |
| <i>Manta birostris</i> (Walbaum, 1792)   | DP               |                          |         |          |                 |               | •                         |
| <b>MONACANTHIDAE</b>   |                  |                          |         |          |                 |               |                           |
| <i>Amanses scopas</i> (Cuvier, 1829)   | BSI              |                          |         |          |                 | •             |                           |
| <i>Cantherhines pardalis</i> (Rüppel, 1837)  | BSI              |                          |         |          |                 | •             |                           |
| <i>Pervagor janthinosoma</i> (Bleeker, 1854)   | BSI              |                          |         |          |                 | •             |                           |
| <b>MULLIDAE</b>  |                  |                          |         |          |                 |               |                           |
| <i>Mulloidichthys flavolineatus</i> (Lacepède, 1801)                                 | NC               |                          |         |          | •               | •             | •                         |
| <i>Mulloidichthys vanicolensis</i><br>Valenciennes in Cuvier and Valenciennes, 1831) | NC               |                          | •       |          | •               | •             |                           |
| <i>Parupeneus barberinus</i> (Lacepède, 1801)  | DC               |                          |         |          | •               |               |                           |
| <i>Parupeneus bifasciatus</i> (Lacepède, 1801)                                       | C                |                          |         |          | •               | •             |                           |
| <i>Parupeneus cyclostomus</i> (Lacepède, 1801)                                       | PI               |                          |         | •        |                 | •             |                           |
| <i>Parupeneus indicus</i> (Shaw, 1803)   | DC               | •                        |         |          |                 |               |                           |
| <i>Parupeneus macronema</i> (Lacepède, 1801)   | DC               |                          | •       | •        | •               | •             |                           |
| <b>MURAENIDAE</b>  |                  |                          |         |          |                 |               |                           |
| <i>Gymnothorax breedeni</i> McCosker and Randall, 1977                               | C                |                          | •       |          |                 |               |                           |
| <i>Gymnothorax favagineus</i> Bloch & Schneider, 1801                                | NC               |                          | •       |          |                 | •             | •                         |
| <i>Gymnothorax meleagris</i> (Shaw, 1795)  | DC               |                          | •       |          |                 | •             |                           |
| <b>MYLIOBATIDAE</b>  |                  |                          |         |          |                 |               |                           |
| <i>Aetobatus narinari</i> (Euphrasen, 1790)  | DC               |                          | •       |          |                 |               |                           |
| <b>NEMIPTERIDAE</b>  |                  |                          |         |          |                 |               |                           |
| <i>Scolopsis ghanam</i> (Forsskål, 1775)   | DC               |                          | •       | •        | •               | •             | •                         |
| <i>Scolopsis vosmeri</i> (Bloch, 1792)   | DC               | •                        |         | •        |                 |               |                           |
| <b>OSTRACIIDAE</b>   |                  |                          |         |          |                 |               |                           |
| <i>Ostracion cubicus</i> (Linnaeus, 1758)  | BSI              | •                        |         |          |                 | •             | •                         |
| <i>Ostracion meleagris</i> Shaw, 1796  | BSI              |                          |         | •        |                 | •             |                           |
| <b>PEMPHERIDAE</b>   |                  |                          |         |          |                 |               |                           |
| <i>Parapricanthus ransonneti</i> Steindachner, 1870                                  | NP               |                          |         | •        |                 |               |                           |
| <i>Pempheris adusta</i> Bleeker, 1877  | NP               |                          |         |          | •               | •             |                           |
| <b>PINGUIPEDIDAE</b>   |                  |                          |         |          |                 |               |                           |
| <i>Parapercis hexophthalma</i> (Cuvier, 1829)  | DC               |                          |         | •        | •               | •             |                           |
| <b>PLATYCEPHALIDAE</b>   |                  |                          |         |          |                 |               |                           |
| <i>Papilloculiceps longiceps</i> (Cuvier, 1829)                                      | DC               | •                        |         | •        |                 | •             |                           |
| <b>POMACANTHIDAE</b>   |                  |                          |         |          |                 |               |                           |
| <i>Apolemichthys trimaculatus</i> (Lacepède, 1831)                                   | O                | •                        | •       | •        | •               |               | •                         |

| <b>FAMILY species</b>  | Trophic Category | Twelve-mile Reef | Tubarão | Garoupa | Kingfish | Lighthouse Reef | Two-mile Reef | Camel's Hump | Amphitheatre |
|--|------------------|------------------|---------|---------|----------|-----------------|---------------|--------------|--------------|
| <i>Centropyge acanthops</i> (Norman, 1922)                                   | O                |                  |         |         |          |                 | •             |              |              |
| <i>Centropyge bispinosus</i> (Günther, 1860)                                 | O                |                  |         |         | •        |                 | •             |              |              |
| <i>Centropyge multispinis</i> (Playfair, 1867)                               | O                | •                | •       | •       | •        | •               | •             |              | •            |
| <i>Pomacanthus chrysurus</i> (Cuvier, 1831)                                  | O                |                  |         |         | •        |                 |               |              |              |
| <i>Pomacanthus imperator</i> (Bloch, 1787)                                   | BSI              | •                | •       | •       | •        | •               | •             | •            | •            |
| <i>Pomacanthus rhomboides</i><br>(Gilchrist and Thompson, 1908)              | C                |                  |         |         |          |                 | •             |              |              |
| <i>Pomacanthus semicirculatus</i><br>(Cuvier in Cuvier & Valenciennes, 1831) | BSI              | •                | •       | •       | •        | •               | •             | •            | •            |
| <b>POMACENTRIDAE</b>   |                  |                  |         |         |          |                 |               |              |              |
| <i>Abudefduf natalensis</i> Hensley & Randall, 1983                          | O                |                  |         |         |          |                 | •             |              |              |
| <i>Abudefduf notatus</i> Day, 1870   | O                |                  |         |         |          | •               |               |              |              |
| <i>Abudefduf sordidus</i> (Forsskål, 1775)                                   | O                |                  |         |         |          | •               |               |              |              |
| <i>Abudefduf sparoides</i> (Quoy & Gaimard, 1825)                            | O                |                  |         |         |          | •               | •             |              |              |
| <i>Abudefduf vaigiensis</i> (Quoy & Gaimard, 1825)                           | O                |                  |         |         |          | •               | •             |              |              |
| <i>Amphiprion akallopisos</i> Bleeker, 1853                                  | O                | •                |         |         | •        |                 | •             |              |              |
| <i>Amphiprion allardi</i> Klausewitz, 1970                                   | O                |                  |         |         | •        |                 | •             |              |              |
| <i>Chromis dimidiata</i> (Klunzinger, 1871)                                  | DP               | •                | •       | •       | •        | •               | •             |              |              |
| <i>Chromis opercularis</i> (Günther, 1867)                                   | DP               |                  | •       |         |          |                 | •             |              |              |
| <i>Chromis weberi</i> (Fowler & Bean, 1928)                                  | DP               |                  |         |         |          |                 | •             |              |              |
| <i>Chrysiptera unimaculata</i> (Cuvier, 1830)                                | O                |                  |         |         |          |                 | •             |              | •            |
| <i>Dascyllus carneus</i> Fischer, 1885                                       | O                |                  |         |         | •        | •               | •             |              |              |
| <i>Dascyllus trimaculatus</i> (Rüppel, 1829)                                 | DP               | •                | •       | •       | •        | •               | •             |              |              |
| <i>Plectroglyphidodon dickii</i> (Liènard, 1839)                             | O                |                  |         |         |          | •               | •             |              |              |
| <i>Plectroglyphidodon johnstonianus</i><br>Fowler & Ball, 1924               | O                |                  |         |         |          | •               | •             |              |              |
| <i>Plectroglyphidodon lacrymatus</i><br>(Quoy & Gaimard, 1825)               | O                |                  |         |         |          | •               | •             |              |              |
| <i>Pomacentrus caeruleus</i> (Quoy & Gaimard, 1825)                          | O                |                  |         | •       |          | •               | •             |              |              |
| <b>PRIACANTHIDAE</b>   |                  |                  |         |         |          |                 |               |              |              |
| <i>Priacanthus hamrur</i> (Forsskål, 1775)                                   | NC               |                  |         |         |          |                 | •             |              |              |
| <b>PSEUDOCHROMIDAE</b>   |                  |                  |         |         |          |                 |               |              |              |
| <i>Pseudochromis dutoiti</i> Smith, 1955                                     | DC               | •                |         | •       |          |                 | •             |              |              |
| <b>SCARIDAE</b>  |                  |                  |         |         |          |                 |               |              |              |
| <i>Scarus frenatus</i> (Lacepède, 1802)                                      | H                |                  |         |         |          | •               | •             |              |              |
| <i>Scarus ghobban</i> Forsskål, 1775   | H                | •                | •       | •       | •        | •               | •             | •            | •            |
| <i>Scarus rubroviolaceus</i> Bleeker, 1847                                   | H                | •                | •       | •       |          | •               | •             |              |              |
| <i>Scarus scaber</i><br>(Valenciennes in Cuvier and Valenciennes, 1840)      | H                |                  |         |         |          |                 | •             |              |              |
| <i>Scarus sordidus</i> (Forsskål, 1775)                                      | H                |                  |         |         |          | •               | •             |              |              |
| <i>Scarus tricolor</i> Bleeker, 1847   | H                |                  |         |         |          |                 | •             |              |              |
| <b>SCIAENIDAE</b>  |                  |                  |         |         |          |                 |               |              |              |
| <i>Umbrina robinsoni</i> Gilchrist and Thompson, 1908                        | C                |                  |         |         |          |                 |               |              | •            |
| <b>SCOMBRIDAE</b>  |                  |                  |         |         |          |                 |               |              |              |
| <i>Euthynnus affinis</i> (Cantor, 1849)                                      | C                |                  |         |         |          |                 | •             |              |              |
| <i>Scomberomorus commerson</i> (Lacepède, 1800)                              | PI               | •                |         | •       |          |                 |               | •            | •            |

| <b>FAMILY species</b>   | Trophic Category | Twelve-mile Reef Tubarão | Garoupa | Kingfish | Lighthouse Reef | Two-mile Reef | Camel's Hump | Amphitheatre |
|---|------------------|--------------------------|---------|----------|-----------------|---------------|--------------|--------------|
| <b>SCORPAENIDAE</b>   |                  |                          |         |          |                 |               |              |              |
| <i>Pterois miles</i> (Bennet, 1825)                                     | PI               | •                        |         | •        |                 |               | •            |              |
| <i>Scorpaenopsis venosa</i> (Cuvier, 1829)                              | C                |                          | •       |          |                 | •             |              |              |
| <b>SERRANIDAE</b>   |                  |                          |         |          |                 |               |              |              |
| <i>Aethaloperca rogae</i> (Forsskål, 1775)                              | NC               |                          | •       |          |                 | •             |              | •            |
| <i>Cephalopholis argus</i> Bloch & Schneider, 1801                      | PI               |                          | •       |          | •               | •             | •            | •            |
| <i>Cephalopholis miniata</i> (Forsskål, 1775)                           | NC               |                          | •       |          |                 | •             |              | •            |
| <i>Epinephelus fasciatus</i> (Forsskål, 1775)                           | NC               |                          |         |          |                 |               | •            |              |
| <i>Epinephelus flavocaeruleus</i> (Lacepède, 1801)                      | PI               |                          |         |          | •               |               |              |              |
| <i>Epinephelus lanceolatus</i> (Bloch, 1790)                            | NC               |                          | •       |          |                 |               |              |              |
| <i>Epinephelus macrospilos</i> (Bleeker, 1855)                          | C                |                          |         |          |                 | •             |              |              |
| <i>Epinephelus malabaricus</i> (Bloch & Schneider, 1801)                | NC               | •                        | •       |          |                 | •             |              |              |
| <i>Epinephelus tukula</i> Morgans, 1959                                 | NC               |                          | •       |          |                 | •             | •            | •            |
| <i>Nemanthias carberryi</i> Smith, 1954                                 | DP               |                          | •       |          |                 | •             |              |              |
| <i>Plectropomus punctatus</i> Quoy & Gaimard, 1824                      | PI               |                          |         |          |                 | •             |              |              |
| <i>Pseudanthias squamipinnis</i> Peters, 1855                           | DP               | •                        | •       | •        | •               | •             |              |              |
| <i>Variola louti</i> (Forsskål, 1775)                                   | PI               | •                        |         |          |                 | •             |              |              |
| <b>SIGANIDAE</b>  |                  |                          |         |          |                 |               |              |              |
| <i>Siganus luridus</i> (Rüppell, 1829)                                  | H                |                          |         |          |                 | •             |              |              |
| <i>Siganus sutor</i><br>(Valenciennes in Cuvier and Valenciennes, 1835) | H                | •                        | •       | •        | •               | •             | •            | •            |
| <b>SPARIDAE</b>   |                  |                          |         |          |                 |               |              |              |
| <i>Acanthopagrus bifasciatus</i> (Forsskål, 1775)                       | DC               |                          |         | •        |                 |               |              |              |
| <b>SPHYRAENIDAE</b>   |                  |                          |         |          |                 |               |              |              |
| <i>Sphyraena barracuda</i> (Walbaum, 1792)                              | DC               |                          |         |          |                 |               | •            |              |
| <i>Sphyraena jello</i> Cuvier in Cuvier and Valenciennes, 1829          | NC               |                          | •       | •        |                 |               |              |              |
| <i>Sphyraena putnamae</i> Jordan and Seale, 1905                        | NC               |                          |         | •        |                 |               |              |              |
| <b>SYNODONTIDAE</b>   |                  |                          |         |          |                 |               |              |              |
| <i>Synodus dermatogenys</i> Fowler, 1912                                | PI               |                          |         |          |                 |               |              | •            |
| <b>TETRAODONTIDAE</b>   |                  |                          |         |          |                 |               |              |              |
| <i>Arothron hispidus</i> (Linnaeus, 1758)                               | NC               |                          |         |          |                 | •             |              |              |
| <i>Arothron nigropunctatus</i> Bloch & Schneider 1801                   | NC               |                          |         |          |                 | •             |              |              |
| <i>Arothron stellatus</i> (Bloch & Schneider 1801)                      | NC               |                          |         |          |                 | •             |              |              |
| <i>Canthigaster smithae</i> Allen and Randall, 1977                     | O                |                          | •       |          |                 |               |              |              |
| <i>Canthigaster solandri</i> (Richardson, 1845)                         | O                |                          |         | •        |                 |               |              |              |
| <i>Canthigaster valentini</i> (Bleeker, 1853)                           | O                |                          | •       | •        |                 | •             |              |              |
| <b>ZANCLIDAE</b>  |                  |                          |         |          |                 |               |              |              |
| <i>Zanclus canescens</i> (Linnaeus, 1758)                               | BSI              | •                        | •       | •        | •               | •             | •            | •            |

58 families at Andavadoaka (south-west Madagascar) and van der Elst & Chater (2001), working at Bassas da India, recorded 305 species. It is not certain why Bazaruto's reefs have a lower species richness and, since the other studies in the region are not directly comparable, it is difficult to place Bazaruto in a spatial or temporal context.

Glorieuses Islands, Juan de Nova and Bassas da India are isolated coral atolls with no permanent human habitation, and consequently experience low to negligible fishing pressure, which has been reported to reduce species richness (McClanahan, 1994; Wantiez *et al.*, 1997). Their isolation from human disturbance and consequent lack of fishing would make them suitable candidates for control studies but their reefs are different from those at Bazaruto. Andavadoaka is more directly comparable with Bazaruto, being exposed to fishing pressure from a nearby fishing village and located on the mainland of Madagascar. Andavadoaka is also further south and therefore at a comparable latitude, negating the latitudinal effect on biodiversity. However, the study at Andavadoaka was conducted over one year compared to nine days at Bazaruto. A previous fish inventory of Bazaruto by van der Elst and Afonso (2008), based on a study done in the late 80s, yielded 269 species from 74 families, but their results included fishery-dependent data and are consequently

not comparable to our study, which used only UVC.

The five reef types sampled in this study comprised a submerged sandstone reef, sedimented rocky patch reefs, a fringing coral reef, a barrier coral reef and two submerged rocky massifs. In terms of reef damage, diver and anchor damage were evident in the coral-covered inner lagoon of Two-mile Reef where large areas of *Acropora* were dead. Corallivorous crown-of-thorns (*Acanthaster planci*) starfish were also observed on Two-mile Reef. These have been persistent, being first recorded at Bazaruto in 1994 (Schleyer, 1998), providing further ecological pressure. Nevertheless, Two-mile Reef had the highest species richness in our study. In other studies, a reduction in hard coral cover resulting from mechanical damage has been linked to recreational SCUBA diving (Hawkins *et al.*, 1999). This reduces reef complexity which correlates with species richness (Bell & Galzin 1984; Gratwicke & Speight 2005), but without long-term monitoring, it is uncertain whether such an effect is taking place on Two-mile Reef.

Garoupa (open to fishing) is a small, flat, sandy ledge with low coral cover and little physical complexity, yet its high fish species richness and abundance was comparable to that of Lighthouse Reef but with six more fish families. The reason for this rich diversity and abundance is unclear; however, it is remarkably similar to Stringer Reef, a small sandy ledge at

Sodwana Bay in South Africa (29° 31.784' S; 32° 40.969' E), which is closed to fishing. Both reefs are similar in terms of coral cover, structural simplicity and high fish abundance (pers. obs.). At Garoupa, the piscivorous lutjanid, *Aprion virescens*, and the scombrid (mackerel), *Scomberomorus commerson*, were prevalent in the mid-water, suggesting a predator-dominated environment, but further ecological investigation of this reef is warranted.

The fish diversity at Twelve-mile Reef (submerged sandstone reef) was expected to be high, given its physical complexity, high coral cover and relative inaccessibility, being furthest offshore. Since the greatest fishing pressure on the Bazaruto reefs is believed to be caused by artisanal fishermen using non-motorised dhows, Twelve-mile Reef should experience relatively little fishing pressure because of its remoteness. High fish diversity may be associated with low fishing effort (McClanahan, 1994; Wantiez *et al.*, 1997) but did not prove the case on Twelve-mile Reef.

The submerged rocky massifs; Camel's Hump and Amphitheatre (both open to fishing) have high vertical relief but are not structurally complex and have minimal coral cover, probably due to high turbidity (Rogers, 1990). Their relatively low fish diversity may be explained by these attributes, which may also be the reason why the top five families (labrids, acanthurids, chaetodonts, pomacentrids and serranids) were underrepresented.

### Trophic Structure

Kulbicki (1988) suggested that trophic structure is usually constant within a region and this has been confirmed in other studies in the WIO (Table 3). Reef disturbances in the form of over-fishing, pollution or coral bleaching have been reported to cause a reduction in the number of carnivores and an increase in herbivores (Chabanet, 2002; Harmelin-Vivien, 1992). While large carnivores are targeted by fishers (Chabanet & Durville 2005), a reduction in coral cover caused by pollution (Rogers, 1990) and

**Table 3. Comparison of trophic structure (%) in reef fish communities in the Western Indian Ocean (WIO).**

| Location         | Reference                       | Carnivores | Omnivores | Herbivores |
|------------------|---------------------------------|------------|-----------|------------|
| Tuléar           | Harmelin-Vivien, 1979           | 74         | 13.5      | 12.5       |
| Réunion          | Chabanet, 1994                  | 51         | 24        | 25         |
| Mayotte          | Chabanet, 2002                  | 69         | 12.5      | 18.5       |
| Geyser and Zéléé | Chabanet <i>et al.</i> , 2002   | 69         | 16        | 15         |
| Glorieuses       | Durville <i>et al.</i> , 2003   | 73         | 12        | 15         |
| Juan de Nova     | Chabanet & Durville 2005        | 73         | 11        | 16         |
| Andavadoaka      | Gillibrand <i>et al.</i> , 2007 | 76         | 11        | 13         |
| Bazaruto         | This study                      | 76         | 12        | 12         |



coral bleaching encourages the rapid growth of filamentous algae, which provides increased food for herbivores (Chabanet, 2002).

Reefs that are considered healthy usually have carnivore levels of between 60-80% (Harmelin-Vivien, 1979), as found on the Bazaruto and other reefs in the WIO (Table 3). Bazaruto had a high proportion of carnivores compared to these other studies. If one compares the Bazaruto fish communities with those in isolated environments with little or no human interference (Glorieuses – Durville *et al.*, 2003; Juan de Nova – Chabanet & Durville 2005), they appear, superficially, to be in a healthy state. However, amalgamation of the carnivorous groups in this study and in many others yields an oversimplification of the situation. Andavadoaka also has an abundance of carnivores (Gillibrand *et al.*, 2007) similar to the Bazaruto reefs, yet experiences fishing pressure and the reefs are reported to be in a degraded state following broad-scale coral bleaching (Gillibrand *et al.*, 2007). It is therefore difficult to link reef health to a simple index such as carnivore abundance.

Future studies should focus on the true trophic hierarchy in the fish communities on the Bazaruto reefs, differentiating between the higher and lower carnivores, rather than between diel preferences. Small carnivorous species (e.g., labrids, chaetodons) are the most abundant and are unlikely

to be affected by fishing pressure. These smaller carnivores may even benefit from fishing because of reduced predation, giving a false impression of reef health. Although a thorough analysis of the carnivore hierarchy was lacking, the proportion of herbivores on Bazaruto's reefs was low, consistent with other reef environments in the region. This indicates a relative measure of reef health where pollution and bleaching are concerned.

## CONCLUSION

The Bazaruto reefs are exposed to fishing pressure, diving and anchor damage, and crown-of-thorns (*Acanthaster planci*) starfish, yet have fish communities rich in diversity and a trophic structure similar to that of other reefs in the WIO which are considered healthy. They endured the 1998 bleaching event without substantial die-off (Schleyer & Celliers 2005). However, without a long-term quantitative monitoring programme, it is difficult to place the health of the Bazaruto reefs in context. This study, like others in the WIO, presents a representative, updated inventory of the fish communities, providing a baseline for more detailed studies. However, further studies should analyse the trophic hierarchy and include abundance measurements. A lack of directly comparable results became evident during our study, highlighting a significant gap in the regional understanding of the reef

fish populations. It is suggested that a long-term monitoring programme (Chabanet & Durville 2005), specifically adapted for Bazaruto's multiple reef-types, would be suitable for conservation planning in the area.

*Acknowledgements:* Sasol provided funding for this research. Eduardo Videira participated in data collection in the first survey. Bruce Mann is thanked for advice in planning the surveys, Fiona Mackay for assistance with data analysis, Taryn Winson and Derick Young for preparation of the map and Pierre Pradervand and Sean Fennessy for their general comments. Ben Thompson, Vicky Page, John Cranswick and Nicolene Rossouw provided logistical support in the field. Anonymous referees are gratefully acknowledged for valuable input to the manuscript.

## REFERENCES

- Bell, J.D. & Galzin, R. (1984) Influence of Live Coral Cover on Coral Reef Fish Communities. *Mar. Ecol. Prog. Ser.* **15**: 265-274.
- Bellwood, D.R. & Hughes, T.P. (2001) Regional-scale Assembly Rules and Biodiversity of Coral Reefs. *Science* **292**:1532-1534.
- Benayahu, Y. & Schleyer, M.H. (1996) Corals of the south-west Indian Ocean III. Alcyonacea (Octocorallia) of Bazaruto Island, Mozambique, with a Redescription of *Cladiella australis* (Macfayden, 1963) and a Description of *Cladiella kashmani* spec. nov. Oceanographic Research Institute, Durban. Investigational Report, 69. 22pp.
- Chabanet, P. (1994) Etude des relations entre les peuplements benthiques et les peuplements ichtyologiques sur le complexe récifal de St-Gilles La Saline à l'île de La Réunion. Thèse Environ. Marin, Univ. Aix-Marseille III, 235pp. + annexes
- Chabanet, P. (2002) Coral Reef Fish Communities of Mayotte (Western Indian Ocean) Two Years After the Bleaching Event. *Mar. Fresh. Wat. Res.* **53**: 107-113.
- Chabanet, P. & Durville, P. (2005) Reef fish Inventory of Juan de Nova's Natural Park (Western Indian Ocean). *Western Indian Ocean Journal of Marine Science* **4**: 145-162.
- Chabanet, P., Tessier, E., Durville, P., Mulochau, T., Rene, F. (2002) Fish Communities of Geysers and Zéléé Coral Banks (Western Indian Ocean). *Cybium* **26**: 11-26.
- Chater, S.A., Beckley, L.E., Garrat, P.A., Ballard, J.A. & van der Elst, R.P. (1993) Fishes from Offshore Reefs in the St Lucia and Maputaland marine reserves, South Africa. *Lammergeyer* **42**: 1-17.
- Chater, S.A., Beckley, L.E., van der Elst, R.P. & Garratt, P.A. (1995) Underwater Visual Census of Fishes in the St Lucia Marine Reserve, South Africa. *Lammergeyer* **43**: 15-23.
- Durville, P., Chabanet, P. & Quod, J.P. (2003) Visual Census of the Reef Fishes in the Natural Reserve of the Glorieuses Islands (Western Indian Ocean). *Western Indian Ocean Journal of Marine Science* **2**: 95-104.
- Everett, B.I., van der Elst, R.P. & Schleyer, M.H. (eds) (2008) *A natural history of Bazaruto Archipelago, Mozambique*. Oceanographic Research Institute, Durban. Special Publication, 8. 118pp.
- Fowler, J. (1987) The Development of Sampling Strategies for Population Studies of Coastal Reef Fishes. A Case Study. *Coral Reefs* **6**: 49-58.

- Froese, R. & Pauly, D. (eds) (2009) FishBase. World Wide Web Electronic Publication. [www.fishbase.org](http://www.fishbase.org). Version (09/2009), November 2009.
- Gillibrand, C.J., Harris, A.R., Mara, E. (2007) Inventory and Spatial Assemblage Study of Reef Fish in the Area of Andavadoaka, south-west Madagascar (Western Indian Ocean). *Western Indian Ocean Journal of Marine Science* **6**: 183-197.
- Gratwicke, B. & Speight, M.R. (2005) The Relationship Between Fish Species Richness, Abundance and Habitat Complexity in a Range of Shallow Tropical Marine Habitats. *Journal of Fish Biology* **66**: 650-667.
- Harmelin-Vivien, M.L. (1979) Ichtyofaune des récifs coralliens en France Outre-Mer. ICRI. Doc. Secrétariat d'Etat à l'Outre-Mer et Ministère de l'Aménagement du Territoire et de l'Environnement. 136pp.
- Harmelin-Vivien, M.L. (1992) Impact des activités humaines sur les peuplements ichthyologiques des récifs coralliens de Polynésie française. *Cybium* **16**: 279-289.
- Harmelin-Vivien, M.L., Harmelin, J., Chauvet, C., Duval, C., Galzin, R., Lejeune, P., Barnabé, G., Blanc, F., Chevalier, R., Duclerc, J. & Lasserre, G. (1985) Evaluation visuelle des peuplements et populations de poissons: méthodes et problèmes. *Revue Ecologie (Terre Vie)* **40**: 467-539.
- Hawkins, J.P., Roberts, C.M., Van't Hoff, T., De Meyer, K., Tratalos, J. & Aldam, C. (1999) Effects of Recreational Scuba Diving on Caribbean Coral and Fish Communities. *Conservation Biology* **13**: 888-897.
- Heemstra, P.C. & Heemstra, E. (2004) *Coastal sea Fishes of southern Africa*. National Inquiry Services Centre, Grahamstown. 512pp.
- Hiatt, W.R. & Strasberg, D.W. (1960) Ecological Relationship of the Fish Fauna on Coral Reefs of the Marshall Islands. *Ecological Monograph* **30**: 65-127.
- Hobson, E.S. (1974) Feeding Relationships of Teleostean Fish on Coral Reefs in Kona, Hawaii. *Fish Bulletin* **72**: 915-1031.
- King, D. (1996) *Reef Fishes and Corals: East Coast of southern Africa*. Struik Publishers, Cape Town. 128pp.
- King, D. & Fraser, V. (2001) *More Reef Fishes and Nudibranchs: East and South Coast of southern Africa*. Struik Publishers, Cape Town. 136pp.
- Kulbicki, M. (1988) Main Variation of the Trophic Structure of Fish Populations in the SW Lagoon of New Caledonia. *Proc. 6th Coral Reef Symp.* Townsville, Australia (August 8-12). **2**: 305-312.
- Lieske, E. & Myers, R. (1999) *Coral Reef Fishes: Caribbean, Indian Ocean, and Pacific Ocean including the Red Sea*. Princeton University Press, Princeton. 400pp.
- Lindén, O., Souter, D., Wilhelmsson, D. & Obura, D.O. (eds) (2002). Coral Reef Degradation in the Indian Ocean: Status Report 2002. CORDIO, Kalmar, Sweden. 284pp.
- McClanahan, T. (1994) Kenyan Coral Reef Lagoon Fish: Effects of Fishing, Substrate Complexity and Sea Urchins. *Coral Reefs* **13**: 231-241.
- Motta, H., Pereira, M.A.M. & Schleyer, M.H. (2002) Coral Reef Degradation in Mozambique, Results of Monitoring 1999-2000. In: Lindén, O., Souter, D., Wilhelmsson, D. & Obura, D.O. (eds) (2002). *Coral Reef Degradation in the Indian Ocean. Status Report 2002*. CORDIO, Kalmar, Sweden. pp55-60.

- Myers, R.F. (1999) *Micronesian Reef Fishes*. Coral Graphics, Barrigada, Guam, 298pp.
- Obura, D.O., Wanyonyi, I.N., Mwaura, J.M. (2002) Participatory Monitoring of an Artisanal Fishery in Kenya. *In*: Lindén, O., Souter, D., Wilhelmsson, D. & Obura, D.O. (eds), *Coral Reef Degradation in the Indian Ocean: Status Report 2000*, CORDIO, Stockholm. pp 70-82.
- Pereira, M.A.M. (2000) *Preliminary Checklist of Reef-Associated Fishes of Mozambique*. MICOA, Maputo. 21pp.
- Reina, A. (1998) Bazaruto Archipelago: Protected Area Development and Management. *Proceedings of International Tropical Marine Ecosystems Management Symposium (ITMEMS)*, Townsville, November, 1998. pp343–353.
- Rodrigues, M.J., Motta, H., Pereira, M.A.M., Gonçalves, M., Carvalho, M. & Schleyer, M.H. (2000) *Coral Reef Monitoring in Mozambique: The Monitoring Programme and 1999 Report*. MICOA-ORI-IIP, Maputo, 64 pp.
- Rogers, C.S. (1990) Responses of Coral Reefs and Reef Organisms to Sedimentation. *Mar. Ecol. Prog. Ser.* **62**: 185-202.
- Samoilys, M.A. (1997) Underwater Visual Census Surveys. *In*: Samoilys MA (ed) *Manual for Assessing Fish Stocks on Pacific coral reefs*. Department of Primary Industries, Training Series QE97009, Queensland, pp 16-29.
- Schleyer, M.H. (1998) Observations on the Incidence Of Crown-of-thorns Starfish in the Western Indian Ocean. *Reef Encounter* **23**: 25-27.
- Schleyer, M.H. & Celliers, L. (2005) The Coral Reefs of Bazaruto Island, Mozambique, with recommendations for their management. *Western Indian Ocean Journal of Marine Science* **4** (2): 227-236.
- Schleyer, M.H. & Maggs, J.Q. (2008) Surveys of Reef Benthos Conducted in the Bazaruto Archipelago on Behalf of Sasol in 2007. Oceanographic Research Institute, Durban. Unpublished Report, 257. 9pp.
- Smith, M.M. & Heemstra, P.C. (1986) *Smith's Sea Fishes*. Struik Publishers, Cape Town: 1047p.
- Van der Elst, R.P. & Afonso, P.S. (2008) Fish and fisheries. *In*: Everett, B.I., van der Elst, R.P. & Schleyer, M.H. (eds) *A Natural History of Bazaruto Archipelago, Mozambique*. Oceanographic Research Institute, Durban. Special Publication, 8. pp 93-109.
- Van der Elst, R.P. & Chater, S. (2001) The Ichthyofauna of Bassas da India atoll in the Mozambique Channel. *6th Fish Indo-Pacific Conf.* Durban, South Africa (May 20-25) (abstract).
- Wantiez, L., Thollot, P., Kulbicki, M. (1997) Effects of Marine Reserves on Coral Reef Fish Communities from Five Islands in New Caledonia. *Coral Reefs* **16**: 215-224.