

The effect of short-term Cu exposure on the oxygen consumption and Cu accumulation of mudfish (*Labeo capensis*) and the largemouth bass (*Micropterus salmoides*) in hard water

WJ van Aardt* and M Hough

School of Environmental Sciences and Development, N.W.U, Potchefstroom Campus, Private Bag X6001, Potchefstroom 2025, South Africa

Abstract

Sediment samples from 3 dams near the gold-mining area in the Mooi River catchment, South Africa, and fish tissue from the mudfish (*Labeo capensis*) and largemouth bass (*Micropterus salmoides*) were analysed for Cu to assess environmental pollution. Copper concentrations of sediment samples in 50 mm deep profiles at Klerkskraal Dam (22.2 mg Cu·kg⁻¹), Boskop Dam (14.1 mg Cu·kg⁻¹) and Potchefstroom Dam (21.7 mg Cu·kg⁻¹) and profiles 100 mm and 150 mm deep were above the risk assessment values for Cu, as implemented by the US EPA. Lowest Cu concentrations were found in gonads and blood samples in fish from both species in the 3 dams, but accumulated 3 to 5 times more, to 110.1±17.8 mg Cu·kg⁻¹ dry mass in the liver. After 120 min Cu exposure at 20°C to 10 mg Cu·l⁻¹ (157.3 mmol Cu·l⁻¹) and a 96 h Cu exposure to 1 mg Cu·l⁻¹ (15.73 mmol Cu·l⁻¹) Cu accumulated mainly in liver tissue and gills. For the mudfish, upon exposure to 10 mg Cu·l⁻¹ (157.3 mmol Cu·l⁻¹), the opercular frequency increased significantly from 80 (± 5.7) cycles·min⁻¹ to above 100 (± 5.8) cycles·min⁻¹ after 90 min, but thereafter decreased to zero cycles·min⁻¹. For largemouth bass the same increase in opercular frequency was found during 10 mg Cu·l⁻¹ exposure, but this Cu level did not stop opercular frequency. For *L. capensis* the oxygen consumption rate MO₂ for the two hour exposure period at 10 mg Cu·l⁻¹ decrease significantly from 5.17 (± 0.32) mmol O₂·l⁻¹·kg⁻¹·h⁻¹ for the controls to 4.5 (± 0.37) mmol O₂·l⁻¹·kg⁻¹·h⁻¹ and for experimental *M. salmoides* from 4.91(± 0.45) mmol O₂·l⁻¹·kg⁻¹·h⁻¹ to 3.13 (±0.74) mmol O₂·l⁻¹·g⁻¹·h⁻¹. For the exposure period of 96 h at 1 mg Cu·l⁻¹, MO₂ for both fish species, decreased to 2.9 (± 0.3) mmol O₂·l⁻¹·kg⁻¹·h⁻¹. It is concluded that:

- The imported *M. salmoides* from the USA is biologically more tolerant to acute Cu exposure compared to the endemic mudfish, *Labeo capensis*
- For the 2 fish species Cu accumulates mainly in the liver, followed by the gills and kidney
- [Cu] above 20 mg Cu·kg⁻¹ dry sediment may be released in the water column if the pH value decreases below 5 and, together with the physical disturbance of the sediment layer, acute Cu pollution will be the result
- Copper is about 10 times more toxic for the 2 fish species studied compared to Pb and Cd in hard water as found in previous studies.

Keywords: Cu, sediment, fish, MO₂, opercular frequency, hard water

Introduction

Effluents from eroded or disused slimes dams at the Anglo-Gold Mine at Carletonville, previously known as the West Wits Goldfields, contain 84.97 μmol Cu·l⁻¹ (5.4 mg·l⁻¹) and 3.97 mmol Zn·l⁻¹ (26.0 mg Zn·l⁻¹) while the dry pelitic sediments contain 484 mg Cu·kg⁻¹ (7616.2 μmol Cu·kg⁻¹) and 6440 mg Zn·kg⁻¹ (98501 μmol Zn·kg⁻¹). The slimes dams receive discharges of high acidity (pH 1.7) containing large amounts of soluble ions (Wittmann and Förstner, 1977). About 139 Ml·d⁻¹ (Winde, 2007) of underground water from these mines, mainly dolomitic in origin (Midgley et al., 1990), are pumped into the Wonderfonteinspruit, part of the upper Mooi River system. Since that study the polluted status of the slimes dams and the mining activities has not changed appreciably (Wade et al., 2002; Coetzee et al., 2006). Van Aardt and Erdman (2004) found a mean value of 35 (± 7.5) mg Cu·kg⁻¹

and 60 (±10.4) mg Zn·kg⁻¹ in the dried clay fraction of sediment samples from the Boskop Dam about 38 km downstream from the slimes dams. The [Cu] in the water of the Wonderfonteinspruit is between 2 and 6 μg Cu·l⁻¹ but in sediment from a dam located in the Wonderfonteinspruit 305 mg Cu·kg⁻¹ was found (Coetzee et al., 2006). The average [Cu] in soils world-wide is 20 mg·kg⁻¹ (Nriagu, 1979). In *Labeo capensis*, a mudfish from the Mooi River Dams, Van Aardt and Erdmann (2004) found 150 (±17.8) mg Cu·kg⁻¹ and 130 (± 23.1) mg Zn·kg⁻¹ in dried liver. Copper concentrations for relatively unpolluted water include a mean of 5 μg Cu·l⁻¹ for 'average river water' while the Amazon and its tributaries range from 0.3 μg·Cu·l⁻¹ to 2.3 μg Cu·l⁻¹ (Boyle, 1976; Kelly, 1988). According to the South African Water Quality Guidelines (DWAF, 1996) for aquatic ecosystems the acute effect value for Cu is 12 μg Cu·l⁻¹ in very hard water.

Generally for aquatic animals such as tubificids, amphipods, crayfish, mud snails and mussels, (Hodson et al., 1979) and in tilapia fish (Van Aardt and Hough, 2006) Cu decreases the oxygen consumption rates. Water hardness affects acute Cu toxicity because Cu-complexes, especially from carbonates such as CuCO₃⁰ and Cu (CO₃)₂²⁻, are less toxic than cupric or

* To whom all correspondence should be addressed.

☎ +27 18 299-2503; fax: +27 18 299-2511;

e-mail: willie.vanaardt@nwu.ac.za

Received 21 July 2006; accepted in revised form 19 July 2007.

Cu hydroxyl ions (Andrew et al., 1977). In hard water systems Cu tends to accumulate in sediments but, if dissolved in water, can still be as high as 25 µg Cu·ℓ⁻¹ (Smith et al., 1996). Toxic Cu levels found in water systems originate mainly from mining processes, the leaching of Cu from minerals due to acid mine drainage and, thirdly, effluents from heavily urbanised and industrialised areas. Generally Cu in aquatic systems is 10 times more toxic compared to Pb, Ni and Zn (Kelly, 1988).

Rainbow trout, *Oncorhynchus mykiss*, at a 96 h exposure to 311 µg Cu·ℓ⁻¹ in pH 7.9 water with only dissolved calcium, undergoes severe iono-regulatory failure combined with a progressive systemic hypoxia and massive haemoglobin concentration (Wilson and Taylor, 1993). This is usually accompanied by excessive production of mucus by the mucus cells on the gill epithelium. The mucus largely stays on the gill surface probably as a protective layer (Tkatcheva et al., 2004). The 96 h LC₅₀ Cu concentrations for fish start from 10 µg Cu·ℓ⁻¹ and end at 10.0 mg Cu·ℓ⁻¹ (McKim et al., 1978). Water hardness, high pH values, or water alkalinity reduce the lethality of Cu to fish (Zitko and Carson, 1976; Howarth and Sprague, 1978; Erickson et al., 1996) but at a pH of lower than 6.5 toxic Cu hydrides are formed (Stouthart et al., 1996). Compared to blue gill and fathead minnow, LC₅₀ for Cu is about 10 times lower in the salmon family (Hodson et al., 1979).

A change in respiration rate (MO₂) is one of the common physiological responses to metal toxicants (Connell et al., 1999) and is easily detectable through changes in the oxygen consumption rates. Generally the MO₂ in fish is reduced when exposed to sub-lethal levels of Cu (Beaumont et al., 2003). However, at lethal Cu concentrations for blue gill (O'Hara, 1971) and *O. mykiss* (Wilson and Taylor, 1993) MO₂ first increases above normal values before it declines before death. At a water pH of lower than 4.0, standard oxygen consumption in fish is reduced and the values of critical oxygen tension (Herried, 1980) increase, with the result that fish are less able to handle environmental hypoxia at low pH values (Ultsch et al., 1978; Van Dijk et al., 1993).

This paper reports the acute effects of Cu on the MO₂ and gill frequency in hard water of an indigenous mudfish, *L. capensis* and the largemouth bass, *Micropterus salmoides*, imported from the USA. Measurements were also made of the Cu concentrations in the water and sediments of the dams in the Mooi River that harbours these fish and also receives water from the Carletonville mining area. Copper levels in the fish tissue for the 2 fish species, before and after exposure to Cu in the laboratory, were also analysed.

Material and methods

Collection and preparation of sediment samples

Six core samples were collected during the summer and winter season of 1998 from each of the 3 dams (Klerkskraal Dam: 26.3.745/27.09.063; Boskop Dam: 26.31.531/27.07.348; Potchefstroom Dam: 16.39.714/27.05.328) in the Mooi River catchment. Sampling localities were situated in the water near the dam inflow and were between 50 and 100 m apart. The core samples were collected using a specially made stainless steel core sampler with a mechanical valve (to be able to salvage the sediment sample), a cutting face and an integral hammer device to be able to drive the core sampler deeply into the dam sediment. The sampler was operated from a boat and took sediment samples from about 2 m below the water surface. Each sample, with an average mass of 900 g, was taken from the core sampler,

enclosed in its polyvinyl jacket, and air dried in a vertical position for 24 h (Van Aardt and Erdmann, 2004).

The top 50 mm sample layer was removed and dried for 12 h at 80°C. With the aid of a Wolfram-ring swing mill (Siebtechnik, Mulheim, Germany) the samples were pulverised for 15 s. One gram from each sediment sample was weighed, 1 ml de-ionised water, 2 ml HNO₃ (70% *pro analysi*) and 1 ml HClO₄ (65% *pro analysi*) from Merck were added and digested at 80°C for 12 h. The digested samples were filled up with de-ionised water to a total volume of 10 ml.

Copper analysis of the sediment and Mooi River water was performed by flame atomic absorption spectrometry at 324.8 nm using standard reference materials (MESS-2) from The National Research Council, Canada, in quality assurance protocols. The calibration graph, with an accuracy of 10 µg Cu·ℓ⁻¹ was linear between 0.01 mg Cu·ℓ⁻¹ to 8.88 mg Cu·ℓ⁻¹. One gram samples from the pulverised cores were also used for energy dispersive X-ray spectrometry analysis (EDAX), using a Phillips EDAX-analyser (EDAX, CDU LEAP Detector) and an electron microscope (XL 30 Phillips DZI). The sample buttons for each sample were prepared using 'sticking film'. The sample was slightly pressed onto the film and covered with a layer of carbon (Emscope TB 500).

Fish collection and the fish-keeping plant

Labeo capensis were collected by gill-netting from the 3 dams in the Mooi River during the summer and winter of 1998. During the same period *Micropterus salmoides* were caught by line fishing. Klerkskraal Dam was used as a control dam compared to Boskop and Potchefstroom Dams (Fig. 1a). The latter 2 dams receive water from mining effluents via the Mooi River (Wade et al., 2002). The fish were transported in a 400 ℓ container filled with aerated dam water and spiked with 12 g NaCl·ℓ⁻¹ water (Walsh, 1984). At the fish plant the fish were treated with 2% formaldehyde and 33 mg malachite green·ℓ⁻¹ for 10 s and transferred to the 5 000 ℓ fish-holding tanks (Van Aardt and Booysen, 2004). *M. salmoides* were fed earthworms or fish fry (*Tilapia sparrmanii*) from Boskop Dam. All fish were kept for at least 3 weeks in the fish-holding tanks at 20°C ±0.5°C before experiments were undertaken at this temperature range.

Chemical analysis of Mooi River water

Water samples were analysed for hardness and macro-elements (Midvaal Water Co., accredited laboratory number T0132).

Copper levels in fish before experimental Cu exposure

From 20 fish caught per dam a 1 ml blood sample was immediately taken by heart puncture whereupon the fish were transported to the laboratory. One gram fish tissue (gonads, the jejunum part of the intestines, gills, liver, kidney and muscle) was dissected out. In a glass vial 2 ml HNO₃ (70% *pro analysi*, Merck) and 1 ml HClO₄ (65% *pro analysi* (Merck) were added and each type of tissue was processed separately and digested as described above for sediment samples. The analysis protocols for Cu were the same as for the sediments except that the standard reference material used was (Dorm-2) from the National Research Council, Canada. To express the metal concentrations obtained from *L. capensis* per gram dried tissue, the percentage of water for each of the seven tissue types was determined as described by van Aardt and Erdmann (2004).

Because of the low numbers of *S. micropterus* caught in Boskop and Klerkskraal Dams, only the tissues from *L. capensis* were analysed for Cu (Fig. 5).

Copper levels in fish after copper exposure

After the Cu exposure experiments, and MO_2 determinations for both *S. micropterus* and *L. capensis* to different Cu concentrations, the 7 types of fish tissue were also harvested from the experimental fish and digested and analysed as described above for control fish.

Oxygen consumption rate measurements (MO_2)

Measurements were done on individual fish after exposure for 120 min to $10 \text{ mg Cu}\cdot\text{l}^{-1}$ ($157.3 \text{ }\mu\text{mol Cu}\cdot\text{l}^{-1}$) as $(\text{NO}_3)_2\cdot 3\text{H}_2\text{O}$. Individual fish were also exposed for 96 h to $1 \text{ mg Cu}\cdot\text{l}^{-1}$ ($15.7 \text{ }\mu\text{mol Cu}\cdot\text{l}^{-1}$) as $\text{Cu}(\text{NO}_3)_2\cdot 3\text{H}_2\text{O}$. Control fish, without Cu exposure, were measured for MO_2 using the same batch caught during the same season as for the experimental fish. All MO_2 measurements for Cu exposed fish were made 2 h after handling of the fish and 24 h after feeding (Jobling and Davies, 1980; Jobling, 1981). To determine the effects of handling on the MO_2 oxygen consumption measurements were also made 0, 2, 4, 6, and 12 h after handling of the fish (Fig. 1a). The mean live fish mass for both control and experimental fish was $864.1 (\pm 85.1) \text{ g}$ for *S. micropterus* and $992 (\pm 112.0) \text{ g}$ for *L. capensis*.

MO_2 and opercular frequency was determined after $10 \text{ mg Cu}\cdot\text{l}^{-1}$ exposure using individual fish in a large 23.03 l Perspex respirometer (Fig. 2) set up as a closed system respirometer (Bridges and Butler, 1989) with continuous PO_2 monitoring of the water. MO_2 calculations and MO_2 corrections for the different fish masses were done according to Van Aardt (1990) and Van Aardt and Booysen (2004). The PO_2 values to calculate MO_2 for each fish were taken at a starting PO_2 value of 120 mm Hg and, after about a 20 min measuring time, again at a PO_2 of 80 mm Hg . In another experiment the progressive decrease of the PO_2 in the respiration water was determined for both fish species at 10 min intervals for 120 min (Fig. 3 a, b). At the same 10 min intervals the opercular frequency was also visually counted (Fig. 4 a, b). The MO_2 determined in 10 control fish (no Cu) and 10 experimental fish exposed to $1 \text{ mg Cu}\cdot\text{l}^{-1}$ for 96 h at 20°C ($\pm 0.5^\circ\text{C}$), was done in a smaller volume (5.15 l) respirometer without PO_2 and opercular monitoring. This was done by exposing the free swimming fish for 84 h in a $1\,000 \text{ l}$ exposure tank. The fish were then caught and placed individually in open respirometers (Van Aardt and Booysen, 2004) in the same exposure tank for a further 12 h in order to eliminate handling stress on MO_2 . After this period the water-tight lids were screwed on for 20 min. The ΔPO_2 was then determined to enable the calculation of MO_2 (Van Aardt, 1990; Van Aardt and Booysen, 2004) (Fig. 8a, b).




Figure 1a
Map of the Mooi River catchment indicating the three dams and the associated towns and cities




Figure 1b
The effects of fish handling on the oxygen consumption rate of the mudfish (*Labeo capensis*, $n=10$) and bass (*Micropterus salmoides*, $n=10$). Vertical bars denote the standard deviation from the mean.




Figure 2
A diagrammatic representation (not to scale) of the large size closed system respirometer. During measurements the respirometer chamber was submerged in a thermostatically controlled water bath. The water turbine prevent the development of partial oxygen tension gradients between the fish and the oxygen electrode.





Figure 3
The progressive decrease of the mean PO_2 in the water measured for (a) *L. capensis* and (b) *M. salmoides* in the large respirometer. Each fish was monitored for 2 h, exposed to $10 \text{ mg Cu} \cdot \text{l}^{-1}$ ($n=10$). Vertical bars denote the standard deviation from the mean





Figure 4
The mean opercular frequency during a 2 h exposure period of (a) *L. capensis* and (b) *M. salmoides* to $10 \text{ mg Cu} \cdot \text{l}^{-1}$. ($n = 10$). Vertical bars denote the standard deviation from the mean.

Results

The water in the 3 dams of the Mooi River can be classified as very hard (Cooney, 1995) with total alkalinity (as CaCO_3) of $252 \text{ mg} \cdot \text{l}^{-1}$; Ca: $67 \text{ mg} \cdot \text{l}^{-1}$; Cl: $36 \text{ mg} \cdot \text{l}^{-1}$; Mg: $50 \text{ mg} \cdot \text{l}^{-1}$; K: $4 \text{ mg} \cdot \text{l}^{-1}$; Na: $28 \text{ mg} \cdot \text{l}^{-1}$; SO_4 : $115 \text{ mg} \cdot \text{l}^{-1}$; pH 8.2. Copper levels in dam sediment (Fig. 5) were $21.1 \pm 6.1 \mu\text{g Cu} \cdot \text{g}^{-1}$, $14.0 \pm 3.3 \mu\text{g Cu} \cdot \text{g}^{-1}$ and




Figure 5
The mean Cu concentration in the upper 50 mm sediment fractions ($n = 6$ per dam) in the three dams situated in the Mooi River system. Vertical bars denote the standard deviation from the mean,




Figure 6
The mean Cu concentration ($\mu\text{g Cu} \cdot \text{g}^{-1}$) in 1 g dried tissue from *L. capensis* collected from the three dams during 1998. Twenty fish per dam were analysed. Vertical bars denote the standard deviation from the mean.

$21.2 \pm 4.7 \mu\text{g Cu} \cdot \text{g}^{-1}$ respectively for Klerkskraal-, Boskop- and Potchefstroom Dam. The percentage Cu in sediment samples from Boskop Dam, analysed by energy dispersive X-ray analysis, (EDAX) was $0.55 \pm 0.06\%$. For Potchefstroom Dam it was $0.54 \pm 0.05\%$ and for Klerkskraal Dam it was $0.51 \pm 0.1\%$. This value represents the percentage occurrence of Cu on the surface of the scanned film compared to the other metal elements.

For the 3 dams Cu in mudfish accumulates 3 to 5 times more in the liver compared to other tissues (Fig. 6). Lowest Cu concentrations were found in gonads and blood samples. The Cu concentration for mudfish liver in the Klerkskraal Dam was $42.10 \pm 9.8 \text{ mg Cu} \cdot \text{kg}^{-1}$ dry tissue. This was less than half the value found for liver tissue in Boskop Dam ($110.1 \pm 17.8 \text{ mg Cu} \cdot \text{kg}^{-1}$) or Potchefstroom Dam ($90.5 \pm 12.2 \text{ mg Cu} \cdot \text{kg}^{-1}$) dried liver.

Copious secretion and production of mucus on the gill lamellae were observed during the $10 \text{ mg Cu} \cdot \text{l}^{-1}$ exposure for both *L. capensis* and *M. salmoides* but not for the $1 \text{ mg Cu} \cdot \text{l}^{-1}$ exposure experiment. In both fish species the cough-reflex was observed when exposed to $10 \text{ mg Cu} \cdot \text{l}^{-1}$. Presumably the inside surface of the oral and opercular cavity was irritated or blocked with mucus (Heath, 1995). For both fish species, Cu accumulated mainly in the liver during the exposure of $1 \text{ mg Cu} \cdot \text{l}^{-1}$ for 96 h or $10 \text{ mg Cu} \cdot \text{l}^{-1}$ for 2 h (Figs. 6 and 7). Copper concentrations in fish tissue, after a 96 h exposure to $1 \text{ mg Cu} \cdot \text{l}^{-1}$ or after 2 h to $10 \text{ mg Cu} \cdot \text{l}^{-1}$, increased more than twofold in liver, kidney and gills for both species. The data were almost the same obtained for *M. salmoides* exposed to $1.0 \text{ mg Cu} \cdot \text{l}^{-1}$ or $10 \text{ mg Cu} \cdot \text{l}^{-1}$ (Fig. 6).

The PO_2 depletion capacity by both *M. salmoides* and *L. capensis* of the water medium was significantly reduced when




Figure 7

The mean Cu concentration ($\mu\text{g Cu}\cdot\text{g}^{-1}$) in 1g dried tissue from *M. salmoides* collected after exposure in the laboratory to $10\text{ mg Cu}\cdot\text{L}^{-1}$ for 2h. The results for 1 mg Cu exposure and the 10 mg Cu exposure for both species were the same when compared with data expressed in Fig. 7. Vertical bars denote the standard deviation from the mean.

fish were exposed to $10\text{ mg Cu}\cdot\text{L}^{-1}$ compared to the controls (Fig. 3a, b). However, when the opercular frequencies between the two fish species were compared, *L. capensis* showed an increase in both control and experimental fish before a traumatic collapse in the opercular frequency for experimental mudfish occurred (Fig. 4 a, b). This happened at PO_2 values of the water below 30 mm Hg and after about 80 min exposure to Cu. All *L. capensis* specimens died before the end of the 120 min exposure time to $10\text{ mg Cu}\cdot\text{L}^{-1}$. For the largemouth bass an initial increase in opercular frequency was found in acute Cu exposure, but it levelled off after about 60 min exposure time (Fig. 4). No deaths or collapse of the opercular frequency were found for largemouth bass. During progressive hypoxia (Fig. 3) it was visually observed that the opercular stroke volume for bass was increased by lowering the lower jaw and the floor of the oral cavity during each opercular opening. This reaction of the gill ventilation to Cu was not observed for mudfish. The MO_2 values obtained for *M. salmoides* using the closed respirometry system compares favourably with the open system operated by Beamish (1970) and Beamish (1974) on the same fish species.

The MO_2 of largemouth bass and mudfish decreased nearly twofold from $5.9\text{ mmol O}_2\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$ to $2.12\text{ mmol O}_2\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$, 6 h after handling (Fig. 1). After this period the oxygen consumption rate stabilised, most probably as resting (standard) MO_2 . Beamish (1970) found for *M. salmoides* at 20°C a standard MO_2 of $3.43\text{ mmol O}_2\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$ for a 150 g fish. In our experiments a statistically significant decrease in the MO_2 between control fish and fish exposed for 2 h to $10\text{ mg Cu}\cdot\text{L}^{-1}$ was found. For the 96 h exposure period to $1\text{ mg Cu}\cdot\text{L}^{-1}$ for both the mudfish and largemouth bass the MO_2 for experimental fish was significantly lower compared with the controls.

Discussion

The hard water in the Mooi River system probably played an important role in the precipitation of Cu compounds before the water from the mines reached the Potchefstroom and Boskop dams. The Cu concentration in the water decreased by more than half, 2h after the pH of the water was increased from a value of below 7 to above 8, in agreement with the findings reported by Shaw and Brown (1974). The result of this pH effect is that a relatively low Cu level is found in Mooi River water. Furthermore the high pH values of the water in the 3 dams increased the adsorption of Cu onto the small clay particles (Kishk and Hassan, 1973; Farrah and Pickering 1977; Daoust et al. 2006).





Figure 8

The oxygen consumption rate from control ($n=10$) and experimental fish ($n=10$) for both species exposed to (a) $10\text{ mg Cu}\cdot\text{L}^{-1}$ and (b) to $1\text{ mg Cu}\cdot\text{L}^{-1}$. Vertical bars denote the standard deviation from the mean.

An average of $18.7 (\pm 7.5)\text{ mg Cu}\cdot\text{kg}^{-1}$ dried sediment was found in sediment samples collected for the 3 dams in the Mooi River. Most of the Cu is contained in the clay fractions (Van Aardt and Erdmann, 2004).

The results of the closed system respirometry used in this study compare favourably with the elaborate open system respirometry used by other workers (Beamish, 1970; Ultsch et al., 1980; Van Aardt and Frey, 1985; De Boeck et al., 1995). To circumvent the possible effects of accumulated excretory products in the closed system respirometry that may influence the respiration rate, the volume of the respiration water should be at least 100 times larger than the volume of an individual fish.

From the opercular frequency results it is clear that the largemouth bass (*Micropterus salmoides*) is physiologically better adapted to acute exposure to Cu in hard water compared to the indigenous *L. capensis*. Why the opercular frequency and MO_2 in *M. salmoides* does not collapse during acute Cu exposure, compared to *L. capensis*, is difficult to explain. It is expected that the much larger gill surface area of largemouth bass, compared to that of the mudfish, should result in a higher Cu exposure per unit surface of the gills resulting in an increased uptake of this metal. McKim et al. (1978) found for *M. salmoides* hatchlings a LC_{50} of $6.97\text{ mg Cu}\cdot\text{L}^{-1}$. This is the most tolerant fish species to Cu toxicity when compared with 7 other freshwater species (McKim et al., 1978). Furthermore the Salmonidae are more sensitive to Cu exposure compared to the Percidae (Spear and Pierce, 1979; Taylor et al., 2003). In this study, opercular frequency measurements indicated that, apart from electrolyte loss, *Labeo capensis*, as a member of the Cyprinidae, is much more sensitive to Cu than largemouth bass, belonging to the Centrarchidae. However, if the oxygen consumption rates of mudfish and largemouth bass are compared on a mass-specific basis, the metabolic rates are nearly the same. Furthermore, the amount of Cu entering the liver, gills and kidney at a 96 h and 2 h exposure periods, respectively, is the same in both fish species. From these data we suggest that *M. salmoides*, being a highly active predator (Rice, 1990), may have better developed neuro-muscular

functions which enables it to operate its opercular muscles during acute Cu exposure, than the relatively sedentary mud fish.

Generally the oxygen consumption rate in fish exposed to Cu decreases. However, an initial stimulation of MO_2 was found for blue gill (O'Hara, 1971), although this was not found by De Boeck et al. (1995) who found an immediate MO_2 decrease in carp. Higher than normal MO_2 values were also noted in our experiments but it could be ascribed to irritation of the gill membranes as well as the epithelium of the oral cavity causing the fish to move around more frequently in the large volume (23.03 l) respirometer compared to the small volume (5.15 l) respirometers. The standard MO_2 of 1.8 mmol $O_2 \cdot kg^{-1} \cdot h^{-1}$ in acidified water at 15°C was found for 1.960 kg carp, *Cyprinus carpio*, (Ultsch et al., 1980) while De Boeck et al. (1995) measured standard MO_2 as 5.3 mmol $O_2 \cdot kg^{-1} \cdot h^{-1}$ in their small 25 g carp. However, for 25 g carp exposed to 53.37 $\mu g \text{ Cu} \cdot l^{-1}$ (0.84 $\mu mol \text{ Cu} \cdot l^{-1}$) the MO_2 decreased substantially in hard water at 20°C to well below 3 mmol $O_2 \cdot kg^{-1} \cdot h^{-1}$ (De Boeck et al., 1995).

In general, the ability of fish, exposed to Cu, to deplete the oxygen content in the water is impaired, compared with controls. This points to a disruption of the branchial structure in fish, probably due to apoptotic death (Wendelaar-Bonga and Lock, 1992; Dang et al., 2000) and not due to damage of the oxygen receptors in the gill arteries that govern gill frequency and ventilation volume (Hughes and Shelton, 1958; Hughes, 1966). According to Heath (1995) metabolic depression observed in fish during Cu exposure could be explained as inhibition of enzymatic reactions related to respiration, and also a decrease in muscle tone resulting in a decrease in spontaneous swimming activity.

The high Ca^{2+} levels found in Mooi River hard water may reduce the permeability of cell membranes including the gill epithelium to water and ions as well as ionic Cu species. High levels of calcium in the water modify the impact of most pollutants by changing membrane permeability to substances such as Cu (Wendelaar-Bonga, 1997). In diluted seawater it was found that ionoregulatory disturbances are caused by increased gill permeability to water and ions by Cu that displaces gill surface bound calcium (Wilson and Taylor, 1993).

Liver Cu concentrations in *M. salmoides* living in Mooi River water are $41.4 \pm 3.7 \mu g \text{ Cu} \cdot g^{-1}$. This is nearly the same concentration ($47.7 \mu g \text{ Cu} \cdot g^{-1}$) found for *M. salmoides* living in a reactor cooling reservoir (Pinder and Giesy, 1981). These values are not above normal values found in fish (Pelgrom et al., 1995). However, the 10 mg $\text{Cu} \cdot l^{-1}$ and 1 mg $\text{Cu} \cdot l^{-1}$ exposed fish in our study had Cu levels of above $75.0 \mu g \cdot \text{Cu}^{-1}$ in liver tissue and gills, indicating that the liver (Felts and Heath, 1984) and the branchial epithelium (Van Heerden et al., 2004) are actively sequestering Cu through metallothionein synthesis (Dang et al., 1999). According to Collvin (1985) the liver in fish implies an increased maintenance cost to detoxify excess Cu. Contrary to other heavy metals and organic pollutants such as organophosphates, Cu exposure of fish does not seem to cause liver histopathology (Baker, 1969). Fish species differences in the pathology to Cu have been recorded (Leland, 1983).

Although chronic and acute stress are mostly associated with increased metabolic rate in fish (Wendelaar-Bonga, 1997) the metabolic rates, as reflected by MO_2 in *M. salmoides* and *L. capensis*, were decreased during short-term exposure to Cu. This suggests a direct toxic effect of Cu on the gills because in both species examined Cu accumulation was significantly higher in gill tissue and also to a degree in kidneys.

The dolomitic origin (Midgley et al., 1990) of the hard water in the Mooi River system in South Africa, with a hardness of 252 $\text{mg} \cdot \text{l}^{-1}$ (as CO_3), should result in the speciation of cupric Cu

to $\text{Cu}(\text{CO}_3)_2^{-2}$; CuCO_3 ; or $\text{Cu}(\text{OH})_2$ in a ratio, respectively, of 0.5; 0.8 and 1.0 (Leckie and Davis, 1979; Tao et al., 2002). It was found (Wagemann and Barica, 1979; Stouthart et al., 1996) that $\text{Cu}(\text{OH})_2$ is the most toxic Cu species.

It is known that acute as well as prolonged Cu exposure to freshwater fish disrupts the osmoregulatory mechanisms in fish with wide differences in sensitivity and tolerance between species to Cu exposure (Taylor et al., 2003). Generally a decrease in K, Na and Cl uptake was found as well as a substantial loss of these electrolytes from the blood plasma, causing massive haemoglobin concentration, increases in the arterial blood pressure and heart failure (Lewis and Lewis, 1971; Courtois and Meyerhoff, 1975; Heath, 1984; Lauren and McDonald, 1985; Wilson and Taylor, 1993). The mechanism of the effect of Cu toxicity on osmoregulation is an inhibition of Na and Cl uptake by the gills caused by Cu that suppresses the activity of Na^+ / K^+ ATP-ase in gill tissue (Heath, 1995; Li et al., 1996; Dang et al., 2000).

Pioneering experiments by Lauren and McDonald (1985) showed that differences in Cu toxicity between hard water and soft water are related to the affinity for complex carbonate, and not to water hardness *per se*. Thus water hardness (e.g. calcium) does not affect Cu speciation but may influence or modify the effects of Cu on physiological functions such as oxygen consumption rate and bio-membrane function in fish. Pagenkopf (1983) suggested that calcium in the water medium may compete with metals for binding sites on fish gill epithelium. Lauren and McDonald (1985) proved that the physiological basis for the ameliorative effects of alkalinity lies in the reduction of toxic effects of Cu on the fish gill capacity to regulate ions. Several physiologists (Potts and Fleming, 1971; Cuthbert and Maetz, 1972) have shown that calcium reduces gill permeability in fish and thus, in freshwater, reduces the diffusion loss of ions across the epithelial layer.

Acute exposure to Cu causes swelling and curling of the gill lamellae and tends to increase the blood/water distance for diffusion, resulting in internal hypoxia in fish (Hughes and Perry, 1976; Satchell, 1984; Mallat, 1985; Poleksić and Mittrivčić-Tutundžić, 1994) and in a reduction of the oxygen consumption rate.

Because of its flexible redox state, Cu in fish plays a vital role in cellular respiration, with cytochrome-c oxidase as the most important Cu protein (Bury et al., 2003). Although Cu is an essential element, with a daily requirement of between 15 to 60 $\mu mol \text{ Cu} \cdot l^{-1} \cdot kg^{-1}$ dry mass (Lanno et al., 1985), copper is toxic to salmonid fish exposed at concentrations as low as 11.95 $\mu mol \cdot l^{-1}$ (0.76 $\text{mg} \cdot \text{l}^{-1}$) at pH 8.6 in hard water (De et al., 1976) and 0.157 $\mu mol \cdot l^{-1}$ (0.1 $\text{mg} \cdot \text{l}^{-1}$) in soft water (Anderson et al., 1989). This metal also causes the immuno suppression of antibody-producing cells (Anderson et al., 1989), and an increase in the number of leucocytes in tilapia (Nusse et al., 1995). Mature, fully functional chloride cells decrease substantially and necrotic apoptotic chloride cell numbers increase in tilapia exposed to copper (*Oreochromis mossambicus*) fish (Dang et al., 2000).

In Mooi River hard water, copper is about 10 times more toxic for tilapia fish compared to cadmium (Van Aardt and Booysen, 2004) or lead (Van Aardt and Venter, 2004). These metals are known for their high toxicity to fish in soft water but in high alkalinity hard water, Cd and Pb completely precipitate from the water column compared to Cu.

Conclusions

It is concluded that:

- The imported *M. salmoides* is physiologically more tolerant to acute Cu exposure compared to the endemic mudfish.

- The high [Cu] found in the sediments in the Mooi River catchment, if released in the water through pH and physical changes would result in acute Cu environmental pollution.
- Copper is about 10 times more toxic for the 2 fish species studied compared to Pb and Cd in hard water.
- The PO₂ depletion capacity for *M. salmoides* exposed to Cu decreases significantly after 15 min compared with the controls. For *L. capensis* PO₂ depletion capacity decreases significantly after 45 min.
- Copper toxicity decreases in hard water but is still toxic if tested at between 1 mg and 10 mg Cu·ℓ⁻¹.
- Copper as toxicant accumulates mainly in fish liver, gills and kidney. This metal can be evaluated, in short-term exposure regimes, using oxygen consumption rate and opercular frequency as physiological parameters.
- Experimental data from the literature (Wendelaar Bonga and Lock, 1992; Tkatcheva et al., 2004) advocate that electrolyte concentrations of Cl, Na and K and the osmotic value of blood plasma should also be measured to evaluate Cu uptake during Cu exposure studies.

Acknowledgements

We wish to thank the Research Focus Area: Environmental Sciences and Management; the National Research Foundation, for financial assistance and the School of Environmental Sciences and Development, NWU, Potchefstroom Campus, Potchefstroom, South Africa, for providing the research facilities.

References

- ANDERSON DP, DIXSON OW, BODAMMER JE and LIZZIO EF (1989) Suppression of antibody-producing cells in rainbow trout spleen sections exposed to copper *in vitro*. *J. Aquat. Anim. Health* **1** 57-61.
- ANDREW RW, BIESINGER KE and GLASS GE (1977) Effects of inorganic complexing on the toxicity of copper to *Daphnia magna*. *Water Res.* **11** 309-315.
- BAKER JTP (1969) Histological and electron microscopic observations on copper poisoning in the winter flounder, *Pseudopleuronectes americanus*. *J. Fish. Res. Bd. Can.* **26** 2785- 2790.
- BEAMISH FWH (1970) Oxygen consumption of largemouth bass, *Micropterus salmoides* in relation to swimming speed and temperature. *Can. J. Zool.* **48** 1221-1228.
- BEAMISH FWH (1974) Apparent specific dynamic action of largemouth bass *Micropterus salmoides*. *J. Fish. Res. Board Can.* **31** 1763-1769.
- BEAUMONT MW, BUTLER PJ and TAYLOR EW (2003) Exposure of brown trout to a sublethal concentration of copper in soft acidic water: effects upon gas exchange and ammonia accumulation. *J. Exp. Biol.* **206** 153-162.
- BRIDGES CR and BUTLER PJ (1989) *Techniques in Comparative Respiratory Physiology. An Experimental Approach*. Cambridge University Press, Cambridge.
- BOYLE EA (1976) Copper in natural waters. In: Nriagu JO (ed.) *Copper in the Environment. Part 1: Ecological Cycling*. John Wiley and Sons, New York. 77-88.
- BURY NR, WALKER PA and GLOVER CN (2003) Nutritive metal uptake in teleost fish. *J. Exp. Biol.* **206** 11-23.
- COETZEE H, WINDE F and WADE PW (2006) An Assessment of Sources, Pathways, Mechanisms and Risks of Current and Potential Future Pollution of Water and Sediments in Gold Mining Areas of the Wonderfonteinpruit Catchment. WRC Report No. 1214/1/06. Water Research Commission, Pretoria, South Africa. 266 pp.
- COLLVIN L (1985) The effect of copper on growth, food consumption and food conversion of perch *Perca fluviatilis* L. offered maximal food rations. *Aquat. Toxicol.* **6** 105-109.
- CONNELL D, LAM P, RICHARDSON B. and WU R (1999) *Introduc-*

- tion to Ecotoxicology*. Blackwell Science, Australia.
- COONEY JD (1995) Fresh water tests. In: Rand GM (ed.) *Fundamentals of Aquatic Toxicology: Effects, Environmental Fate, and Risk Assessment*. Taylor and Francis, Washington, DC. 71-102.
- COURTOIS LA and MEYERHOFF RD (1975) Effects of copper exposure on water balance. *Bull. Environ. Contam. Toxicol.* **14** 211-219.
- CUTHBERT AW and MAETZ M (1972) The effects of calcium and magnesium on sodium fluxes through the gills of *Carassius auratus*. *L. J. Physiol.* **221** 633-643.
- DANG ZC, LOCK RAC, FLIK G and WENDELAAR-BONGA SE (1999) The metallothionein response in gills of *Oreochromis mossambicus* exposed to copper in fresh water. *Am. J. Physiol.* **277** R320-R331.
- DANG Z, LOCK R AC, FLIK G and WENDELAAR-BONGA SE (2000) Na⁺/ K⁺ ATP-ase immunoreactivity in branchial chloride cells of *Oreochromis mossambicus* chronically exposed to copper. *J. Exp. Biol.* **203** 379-387.
- DAOUST CM, BASTIENC and DESCHÊNES L (2006) Influence of soil properties and aging on the toxicity of copper on compost worm and barley. *J. Environ. Qual.* **35** 558-567.
- DE JF, SOLBE JF and COPPER VA (1976) Studies on the toxicology of copper sulphate to Stone Loach *Noenacheilus barbatulus* (L) in hard water. *Water Res.* **19** 523-527.
- DE BOECK G, SMET H and BLUST R (1995) The effect of sublethal levels of copper on oxygen consumption and ammonia excretion in the common carp, *Cyprinus carpio*. *Aquat. Toxicol.* **32** 27-141.
- DWAF (Department of Water Affairs and Forestry) (1996) *South African Water Quality Guidelines. Vol.7: Aquatic Ecosystems*. Pretoria.
- ERIKCSON RJ, BENOIT DA, MATTSON VR, NELSON HP and LEONARD NL (1996) The effects of water chemistry on the toxicity of copper to fathead minnows. *Environ. Toxicol. Chem.* **15** 181-193.
- FARRAH, H. and PICKERING, W.F. (1977). Influence of clay-solute interactions on aqueous heavy metal ion levels. *Water Air Soil Pollut.* **8** 189-197.
- FELTS PA and HEATH AG (1984) Interactions of temperature and sublethal environmental copper exposure on the energy metabolism of bluegill, *Lepomis macrochirus* Rafinesque. *J. Fish Biol.* **25** 445-453.
- HEATH AG (1984) Changes in tissue adenylates and water content of bluegill, *Lepomis macrochirus*, exposed to copper. *J. Fish. Biol.* **24** 299-309.
- HEATH AG (1995) *Water Pollution and Fish Physiology*. CRC Press, Boca Raton, Florida, USA.
- HERRIED CF (1980) Hypoxia in invertebrates. *Comp. Biochem. Physiol.* **67A** 311-320.
- HODSON V, BORGMAN U and SHEAR H (1979) Toxicity of copper to aquatic biota. In: Nriagu JO (ed.) *Copper in the Environment Part II Environmental and Occupational Exposure to Copper*. John Wiley & Sons, New York. 307-372.
- HOWARTH RS and SPRAGUE JB (1978) Copper lethality to rainbow trout in water of various hardness and pH. *Water Res.* **12** 455-462.
- HUGHES GM (1966) The dimensions of fish gills in relation to their function. *J. Exp. Biol.* **45** 177-195.
- HUGHES GM and PERRY SF (1976) A morphometric study of trout gills: a light-microscopic method for the evaluation of pollutant action. *J. Exp. Biol.* **64** 447-460.
- HUGHES GM and SHELTON G (1958) The mechanism of gill ventilation in three freshwater teleosts. *J. Exp. Biol.* **35** 807-823.
- HUGHES GM and UMEZAWA S-I (1968) On respiration in the dragonet *Callinymus lyra*. *J. Exp. Biol.* **49** 565-571.
- JOBLING M (1981) The influences of feeding on the metabolic rate of fishes: a short review. *J. Fish Biol.* **18** 385-400.
- JOBLING M and DAVIES PS (1980) Effects of feeding on metabolic rate, and the specific dynamic action in Plaice, *Pleuronectes platessa* L. *J. Fish Biol.* **16** 629-638.
- KISHK FM and HASSAN MN (1973) Sorption and desorption of copper by and from clay minerals. *Plant Soil* **39** 497-505.
- KELLY M (1988) *Mining and the Freshwater Environment*. Elsevier, Applied Science, London.
- LANNO RP, SLINGER SJ and HILTON JW (1985) Maximum tolerable and toxicity levels of dietary copper in rainbow-trout (*Salmo gairdneri* Richardson). *Aquacult.* **49** 257-268.

- LAURÉN DJ and MCDONALD DG (1985) Effects of copper on branchial iono-regulation in the rainbow trout, *Salmo gairdneri* Richardson – modulation by water hardness and pH. *J. Comp. Physiol.* **155** 635-644.
- LECKIE JO and DAVIS JA (1979) Aqueous environmental chemistry of copper. In: Nriagu JO (ed.) *Copper in the Environment. Part I: Ecological Cycling*. John Wiley & Sons, New York. 89-121.
- LELAND HV (1983) Ultrastructural changes in the hepatocytes of juvenile rainbow trout and mature brown trout exposed to copper or zinc. *Environ. Toxicol. Chem.* **2** 353-359.
- LI J, LOCK RA, KLAREN PHM, SWARTS HGP, SCHUURMANS-STEKHOVEN FMA and WENDELAAR-BONGA SE (1996) Kinetics of copper inhibition of Na⁺/K⁺-ATPase. *Toxicol. Lett.* **87** 31-38.
- LEWIS SD and LEWIS WM (1971) The effect of zinc and copper on the osmolality of blood serum of the channel catfish, *Ictalurus punctatus* Rafinesque, and golden shiner, *Notemigonus crysoleucas* Mitchill. *Trans. Am. Fish Soc.* **100** 639-643.
- MALLAT J (1985) Fish gill structural changes induced by toxicants and other irritants: a statistical review. *Can. J. Fish. Aquat. Sci.* **42** 630-648.
- McKIM JM, EATON JG and HOLCOMBE GW (1978) Metal toxicity to embryos and larvae of eight species of freshwater fish. II: Copper. *Bull. Environ. Contam. Toxicol.* **19** 608-616.
- MIDGLEY DC, PITMANN WV and MIDDLETON BJ (1994) Surface Water Resources of South Africa 1990. WRC Report No 298/1.1 through to 6.2. Water Research Commission, Pretoria, South Africa.
- NRIAGU JO (1979) The global copper cycle. In: JO Nriagu (ed.) *Copper in the Environment*. John Wiley and Sons, New York. 1-17.
- NRIAGU J O and PACYNA JM (1988) Quantitative assessment of worldwide contamination of air, water and soils by trace metals. *Nature* **333** 134-139.
- NUSSEY G, VAN VUREN JHJ and DU PREEZ HH (1995) Effect of copper on the differential white blood cell counts of the Mozambique tilapia (*Oreochromis mossambicus*). *Comp. Biochem. Physiol.* **111C** 381-388.
- O'HARA J (1971) Alterations in oxygen consumption by bluegills exposed to sublethal treatment with copper. *Water Res.* **5** 321-329.
- PAGENKOPF GK (1983) Gill surface interaction model for trace-metal toxicity to fishes: role of complexation, pH and water hardness. *Environ. Sci. Technol.* **17** 342-347.
- PELGROM SMGJ, LAMMERS LPM, LOCK RAC, BALM PHM and WENDELAAR-BONGA SE (1995) Interactions between copper and cadmium modify metal organ distribution in mature tilapia, *Oreochromis mossambicus*. *Environ. Pollut.* **90** 415-423.
- PINDER JE and GIESY JP (1981) Frequency distributions of the concentrations of essential and nonessential elements in largemouth bass, *Micropterus salmoides*. *Ecol.* **62** 456-468.
- POLEKSIĆ Vand MITROVIĆ-TUTUNDŽIĆ V (1994) Fish gills as a monitor of sublethal and chronic effects of pollution. In: Muller R and Lloyd R (eds.) *Sublethal and Chronic Effects of Pollutants on Freshwater Fish*. Cambridge University Press, Cambridge. 339-352.
- POTTS WTW and FLEMINGWR (1971) The effects of environmental calcium and ovine prolactin on sodium balance in *Fundulus kansae*. *J. Exp. Biol.* **116** 411-433.
- RICE JA (1990) Bioenergetics modeling approaches to evaluation of stress in fish. *Am. Fish. Soc. Symp.* **8** 80-92.
- SATCHELL GH (1984) Respiratory toxicology of fishes. In: Weber LJ (ed.) *Aquatic Toxicology*. Raven Press, New York. 23-45.
- SHAW TL and BROWN VM (1974) The toxicity of some forms of copper to rainbow trout. *Water Res.* **8** 377-382.
- SMITH S, CHEN M-H, BAILEY RG and WILLIAMS WP (1996) Concentration and distribution of copper and cadmium in water, sediments, detritus, plants and animals in a hardwater lowland river. *Hydrobiol.* **341** 71-80.
- SPEAR PA and PIERCE RC (1979) Copper in the Aquatic Environment: Chemistry, Distribution and Toxicology. National Research Council of Canada, (NRCC 16454), Ottawa.
- STOUGHTART AH, HAANS JLM, LOCK RAC and WENDELAAR-BONGA SE (1996) Effects of water pH on copper toxicity to early life stages of the common carp (*Cyprinus carpio*). *Environ. Toxicol. Chem.* **15** 376-383.
- TAO, S, LONG A, XU M and DAWSON RW (2002). Copper Speciation in the Gill Microenvironment of Carp (*Cyprinus carpio*) at Various Levels of pH. *Ecotox. Environ. Safety* **52** 211-226.
- TAYLOR, LN WOOD CM and MCDONALD DG (2003) An evaluation of sodium loss and gill metal binding properties in rainbow trout and yellow perch to explain species differences in copper tolerance. *Environ. Toxicol. Chem.* **22** 2159-2166.
- TKATCHEVA V, HYVÄRINEN H, KUKKONEN J, RYZHKOV LP and HOLOPAINEN I J (2004) Toxic effects of mining effluents on fish gills in a subarctic lake system in NW Russia. *Ecotox. Environ. Safety* **57** 278-289.
- ULTSCH G (1978) Oxygen consumption as a function of pH in three species of freshwater fishes. *Copeia* 272-283.
- ULTSCH G, OTT M and HEISLER N (1980) Acid-base and electrolyte status in carp (*Cyprinus carpio*) exposed to low environmental pH. *J. Exp. Biol.* **93** 65-80.
- VAN AARDT WJ (1990) Respirometrie soos toegepas op terreestriële en akwatiese invertebrate. *S. Afr. Tydskr. Natuurwet.Tegnot.* **9** 149-161.
- VAN AARDT WJ and ERDMANN R (2004) Heavy metals (Cd, Pb, Cu, Zn) in mudfish and sediments from three hard-water dams of the Mooi River catchment, South Africa. *Water SA* **30** 211-218. <http://www.wrc.org.za/archives/watersa%20archive/2004/Apr-04/9.pdf>
- VAN AARDT WJ and BOOYSEN J (2004) Water hardness and the effects of Cd on oxygen consumption, plasma chlorides and bioaccumulation in *Tilapia sparrmanii*. *Water SA* **30** 57- 64. <http://www.wrc.org.za/archives/watersa%20archive/2004/Jan-04/9.pdf>
- VAN AARDT WJ and FREY BJ (1985) Effects of temperature and hypoxic stress on the oxygen consumption rates of the mudsucker fish *Labeo capensis* (Smith). *S. Afr. J. Zool.* **12** 156-160.
- VAN AARDT WJ and HOUGH M (2006) Acute effects of Cu on oxygen consumption and 96 hr-LC₅₀ values in the fresh water fish *Tilapia sparrmanii* (Teleostei:Cichlidae) in Mooi River hard water, South Africa. *Afr. J. Aquat. Sci.* **31** (2) 305-311.
- VAN AARDT WJ and VENTER LCR (2004) Toxicity of high levels of lead in hard water: biological effects on freshwater fish, *Tilapia sparrmanii*, in the laboratory. *Afr. J. Aquat. Sci.* **29** 37-46.
- VAN DIJK P, VAN DEN THILLART G and WENDELAAR-BONGA SE (1993) Is there a synergistic effect between steady-state exercise and water acidification in carp? *J. Fish Biol.* **42** 673-680.
- VAN HEERDEN DALÉNE, VOSLOO A and NIKINMAA M (2004) Effects of short-term copper exposure on gill structure, metallothionein and hypoxia-inducible factor-1 α (HIF-1 α) levels in rainbow trout (*Oncorhynchus mykiss*). *Aquat. Toxicol.* **69** 271-280.
- VAN RENSBURG WCJ and PRETORIUS DA (1977) *South Africa's Strategic Minerals – Pieces on a Continental Chess Board*. Valiant Publishers, Johannesburg.
- WADE PW, WOORDBOURNE S, MORRIS WM, VOS P and JARVIS NV (2002): Tier 1 Risk Assessment of Radionuclides in Selected Sediments of the Mooi River. WRC Report No 1095/1/02. Water Research Commission, Pretoria, South Africa.
- WAGEMANN R and BARICA J (1979) Speciation and rate of loss of copper from lake water with implications to toxicity. *Water Res.* **13** 515-523.
- WALSH AH (1984) Biology and diseases of fish. In: Fox JG (ed.) *Laboratory Animal Medicine*. Academic Press, London. 477-541.
- WENDELAAR-BONGA SE (1997) The stress response in fish. *Physiol. Rev* **77** 591-625.
- WENDELAAR-BONGA SE and LOCK RCA (1992) Toxicants and osmoregulation in fish. *Neth. J. Zool.* **42** 478-493.
- WINDE, F (2007) Personal communication, Dept. Geography, North West University, Potchefstroom
- WILSON RW and TAYLOR EW (1993) The physiological responses of freshwater rainbow trout, *Oncorhynchus mykiss*, during acutely lethal copper exposure. *J. Comp. Physiol.* **163** 38-47.
- WITTMANN GTW and FÖRSTNER U (1977) Heavy metal enrichment in mine drainage: III The Klerksdorp, West Wits and Evander Goldfields. *S. Afr. J. Sci.* **73** 53-57.
- ZITKO V and CARSON WG (1976) A mechanism of the effects of water hardness on the lethality of heavy metals to fish. *Chemosphere* **5** 299-303.